



# THE PRINCIPLE OF RELATIVITY

A COLLECTION OF ORIGINAL MEMOIRS ON THE  
SPECIAL AND GENERAL THEORY OF RELATIVITY

BY

H. A. LORENTZ, A. EINSTEIN  
H. MINKOWSKI AND H. WEYL


WITH NOTES BY

A. SOMMERFELD

TRANSLATED BY

W. PERRETT AND G. B. JEFFERY

WITH SEVEN DIAGRAMMS



DOVER PUBLICATIONS, INC.

IIA LIB

Published in Canada by General Publishing Company, Ltd., 30 Lesmill Road, Don Mills, Toronto, Ontario.

Published in the United Kingdom by Constable and Company, Ltd., 10 Orange Street, London WC 2.

This Dover edition, first published in 1952, is an unabridged and unaltered republication of the 1923 translation first published by Methuen and Company, Ltd. in 1923. This edition is reprinted through special arrangement with Methuen and Company and Albert Einstein.

*Standard Book Number: 486-60081-5*

*Library of Congress Catalog Card Number: A52-9845*

Manufactured in the United States of America  
Dover Publications, Inc.  
180 Varick Street  
New York, N.Y. 10014

## TRANSLATORS' PREFACE

THE Theory of Relativity is at the moment the subject of two main lines of inquiry : there is an endeavour to express its principles in logical and concise form, and there is the struggle with analytical difficulties which stand in the way of further progress. In the midst of such problems it is easy to forget the way in which the theory gradually grew under the stimulus of physical experiment, and thus to miss much of its meaning. It is this growth which the present collection of papers is designed chiefly to exhibit. In the earlier papers there are some things which the authors would no doubt now express differently ; the later papers deal with problems which are not by any means yet fully solved. At the end we must confess that Relativity is still very much of a problem—and therefore worthy of our study.

The authors of the papers are still actively at work on the subject—all save Minkowski. His paper on "Space and Time" is a measure of the loss which mathematical physics suffered by his untimely death.

The translations have been made from the text, as published in a German collection, under the title "Des Relativitätsprinzip" (Teubner, 4th ed., 1922).



## vi THE PRINCIPLE OF RELATIVITY

The second paper by Lorentz is an exception to this. It is reprinted from the original English version in the Proceedings of the Amsterdam Academy. Some minor changes have been made, and the notation has been brought more nearly into conformity with that employed in the other papers.

W. P.  
G. B. J.

# TABLE OF CONTENTS

	PAGES
I. MICHELSON'S INTERFERENCE EXPERIMENT. By H. A. Lorentz .	1-7
§ 1. The experiment. § 2. The contraction hypothesis. §§ 3-4.	
The contraction in relation to molecular forces.	
II. ELECTROMAGNETIC PHENOMENA IN A SYSTEM MOVING WITH ANY VELOCITY LESS THAN THAT OF LIGHT. By H. A. Lorentz .	9-34
§ 1. Experimental evidence. § 2. Poincaré's criticism of the contraction hypothesis. § 3. Maxwell's equations for moving axes. § 4. The modified vectors. § 5. Retarded potentials. § 6. Electrostatic fields. § 7. A polarized particle. § 8. Corresponding states. § 9. Momentum of an electron. § 10. The influence of the earth's motion on optical phenomena. § 11. Applications. § 12. Molecular motions. § 13. Kaufmann's experiments.	
III. ON THE ELECTRODYNAMICS OF MOVING BODIES. By A. Einstein.	35-65
KINEMATICAL PART. § 1. Definition of simultaneity. § 2. On the relativity of lengths and times. § 3. The transforma- tion of co-ordinates and times. § 4. Physical meaning of the equations. § 5. The composition of velocities.	
ELECTRODYNAMICAL PART. § 6. Transformation of the Max- well-Hertz equations. § 7. Doppler's principle and aber- ration. § 8. The energy of light rays and the pressure of radiation. § 9. Transformation of the equations with convection currents. § 10. Dynamics of the slowly accel- erated electron.	
IV. DOES THE INERTIA OF A BODY DEPEND UPON ITS ENERGY- CONTENT? By A. Einstein . . . . .	67-71
V. SPACE AND TIME. By H. Minkowski . . . . .	73-91
I. The invariance of the Newtonian equations and its represen- tation in four dimensional space. II. The world-postulate.	
III. The representation of motion in the continuum. IV. The new mechanics. V. The motion of one and two electrons.	
<i>Notes on this paper.</i> By A. Sommerfeld . . . . .	92-96
VI. ON THE INFLUENCE OF GRAVITATION ON THE PROPAGATION OF LIGHT. By A. Einstein . . . . .	97-108
§ 1. The physical nature of gravitation. § 2. The gravitation of energy. § 3. The velocity of light. § 4. Bending of light-rays.	

# viii THE PRINCIPLE OF RELATIVITY

PAGES

VII. THE FOUNDATION OF THE GENERAL THEORY OF RELATIVITY. By A. Einstein . . . . .	109-164
A. FUNDAMENTAL CONSIDERATIONS ON THE POSTULATE OF RELATIVITY. § 1. Observations on the special theory. § 2. The need for an extension of the postulate of relativity. § 3. The space-time continuum; general co-variance. § 4. Measurement in Space and Time.	
B. MATHEMATICAL AIDS TO THE FORMULATION OF GENERALLY COVARIANT EQUATIONS. § 5. Contravariant and covariant four-vectors. § 6. Tensors of the second and higher ranks. § 7. Multiplication of tensors. § 8. The fundamental tensor $g_{\mu\nu}$ . § 9. The equation of the geodetic line. § 10. The formation of tensors by differentiation. § 11. Some cases of special importance. § 12. The Riemann-Christoffel tensor.	
C. THEORY OF THE GRAVITATIONAL FIELD. § 13. Equations of motion of a material point. § 14. The field equations of gravitation in the absence of matter. § 15. The Hamiltonian function for the gravitational field. Laws of momentum and energy. § 16. The general form of the field equations. § 17. The laws of conservation. § 18. The laws of momentum and energy.	
D. MATERIAL PHENOMENA. § 19. Euler's equations for a fluid. § 20. Maxwell's equations for free space.	
E. APPLICATIONS OF THE THEORY. § 21. Newton's theory as a first approximation. § 22. Behaviour of rods and clocks in a static gravitational field. Bending of light rays. Motion of the perihelion of a planetary orbit.	
VIII. HAMILTON'S PRINCIPLE AND THE GENERAL THEORY OF RELATIVITY. By A. Einstein . . . . .	165-178
§ 1. The principle of variation and the field-equations. § 2. Separate existence of the gravitational field. § 3. Properties of the field equations conditioned by the theory of invariants.	
IX. COSMOLOGICAL CONSIDERATIONS ON THE GENERAL THEORY OF RELATIVITY. By A. Einstein . . . . .	175-188
§ 1. The Newtonian theory. § 2. The boundary conditions according to the general theory of relativity. § 3. The spatially finite universe. § 4. On an additional term for the field equations of gravitation. § 5. Calculation and result.	
X. DO GRAVITATIONAL FIELDS PLAY AN ESSENTIAL PART IN THE STRUCTURE OF THE ELEMENTARY PARTICLES OF MATTER? By A. Einstein . . . . .	189-198
§ 1. Defects of the present view. § 2. The field equations freed of scalars. § 3. On the cosmological question. § 4. Concluding remarks.	
XI. GRAVITATION AND ELECTRICITY. By H. Weyl . . . . .	200-216

# MICHELSON'S INTERFERENCE EXPERIMENT

BY

H. A. LORENTZ

*translated from "Versuch einer Theorie der elektrischen  
und optischen Erscheinungen in bewegten Körpern,"  
Leiden, 1895, §§ 89-92.*



# MICHELSON'S INTERFERENCE EXPERIMENT

By H. A. LORENTZ

1. **A**S Maxwell first remarked and as follows from a very simple calculation, the time required by a ray of light to travel from a point A to a point B and back to A must vary when the two points together undergo a displacement without carrying the ether with them. The difference is, certainly, a magnitude of second order; but it is sufficiently great to be detected by a sensitive interference method.

The experiment was carried out by Michelson in 1881.\* His apparatus, a kind of interferometer, had two horizontal arms, P and Q, of equal length and at right angles one to the other. Of the two mutually interfering rays of light the one passed along the arm P and back, the other along the arm Q and back. The whole instrument, including the source of light and the arrangement for taking observations, could be revolved about a vertical axis; and those two positions come especially under consideration in which the arm P or the arm Q lay as nearly as possible in the direction of the Earth's motion. On the basis of Fresnel's theory it was anticipated that when the apparatus was revolved from one of these *principal positions* into the other there would be a displacement of the interference fringes.

But of such a displacement—for the sake of brevity we will call it the Maxwell displacement—conditioned by the change in the times of propagation, no trace was discovered, and accordingly Michelson thought himself justified in concluding that while the Earth is moving, the ether does not remain at rest. The correctness of this inference was soon brought into question, for by an oversight Michelson had

\* Michelson, American Journal of Science, 22, 1881, p. 120.

taken the change in the phase difference, which was to be expected in accordance with the theory, at twice its proper value. If we make the necessary correction, we arrive at displacements no greater than might be masked by errors of observation.

Subsequently Michelson \* took up the investigation anew in collaboration with Morley, enhancing the delicacy of the experiment by causing each pencil to be reflected to and fro between a number of mirrors, thereby obtaining the same advantage as if the arms of the earlier apparatus had been considerably lengthened. The mirrors were mounted on a massive stone disc, floating on mercury, and therefore easily revolved. Each pencil now had to travel a total distance of 22 meters, and on Fresnel's theory the displacement to be expected in passing from the one principal position to the other would be 0.4 of the distance between the interference fringes. Nevertheless the rotation produced displacements not exceeding 0.02 of this distance, and these might well be ascribed to errors of observation.

Now, does this result entitle us to assume that the ether takes part in the motion of the Earth, and therefore that the theory of aberration given by Stokes is the correct one? The difficulties which this theory encounters in explaining aberration seem too great for me to share this opinion, and I would rather try to remove the contradiction between Fresnel's theory and Michelson's result. An hypothesis which I brought forward some time ago,† and which, as I subsequently learned, has also occurred to Fitzgerald,‡ enables us to do this. The next paragraph will set out this hypothesis.

2. To simplify matters we will assume that we are working with apparatus as employed in the first experiments, and that in the one principal position the arm P lies exactly in

\* Michelson and Morley, *American Journal of Science*, 34, 1887, p. 333; *Phil. Mag.*, 24, 1887, p. 449.

† Lorentz, *Zittingsverslagen der Akad. v. Wet. te Amsterdam*, 1892-98, p. 74.

‡ As Fitzgerald, kindly tells me, he has for a long time dealt with his hypothesis in his lectures. The only published reference which I can find to the hypothesis is by Lodge, "Aberration Problems," *Phil. Trans. R.S.*, 184 A, 1893.

the direction of the motion of the Earth. Let  $v$  be the velocity of this motion,  $L$  the length of either arm, and hence  $2L$  the path traversed by the rays of light. According to the theory,\* the turning of the apparatus through  $90^\circ$  causes the time in which the one pencil travels along  $P$  and back to be longer than the time which the other pencil takes to complete its journey by

$$\frac{Lv^2}{c^3}.$$

There would be this same difference if the translation had no influence and the arm  $P$  were longer than the arm  $Q$  by  $\frac{1}{2}Lv^2/c^2$ . Similarly with the second principal position.

Thus we see that the phase differences expected by the theory might also arise if, when the apparatus is revolved, first the one arm and then the other arm were the longer. It follows that the phase differences can be compensated by contrary changes of the dimensions.

If we assume the arm which lies in the direction of the Earth's motion to be shorter than the other by  $\frac{1}{2}Lv^2/c^2$ , and, at the same time, that the translation has the influence which Fresnel's theory allows it, then the result of the Michelson experiment is explained completely.

Thus one would have to imagine that the motion of a solid body (such as a brass rod or the stone disc employed in the later experiments) through the resting ether exerts upon the dimensions of that body an influence which varies according to the orientation of the body with respect to the direction of motion. If, for example, the dimensions parallel to this direction were changed in the proportion of 1 to  $1 + \delta$ , and those perpendicular in the proportion of 1 to  $1 + \epsilon$ , then we should have the equation

$$\epsilon - \delta = \frac{1}{2} \frac{v^2}{c^2} \quad . \quad . \quad . \quad . \quad (1)$$

in which the value of one of the quantities  $\delta$  and  $\epsilon$  would remain undetermined. It might be that  $\epsilon = 0$ ,  $\delta = -\frac{1}{2}v^2/c^2$ , but also  $\epsilon = \frac{1}{2}v^2/c^2$ ,  $\delta = 0$ , or  $\epsilon = \frac{1}{4}v^2/c^2$ , and  $\delta = -\frac{1}{4}v^2/c^2$ .

3. Surprising as this hypothesis may appear at first sight,

\* Cf. Lorentz. Arch. Néerl. 2. 1887. pp. 168-176.



yet we shall have to admit that it is by no means far-fetched, as soon as we assume that molecular forces are also transmitted through the ether, like the electric and magnetic forces of which we are able at the present time to make this assertion definitely. If they are so transmitted, the translation will very probably affect the action between two molecules or atoms in a manner resembling the attraction or repulsion between charged particles. Now, since the form and dimensions of a solid body are ultimately conditioned by the intensity of molecular actions, there cannot fail to be a change of dimensions as well.

From the theoretical side, therefore, there would be no objection to the hypothesis. As regards its experimental proof, we must first of all note that the lengthenings and shortenings in question are extraordinarily small. We have  $v^2/c^2 = 10^{-8}$ , and thus, if  $\epsilon = 0$ , the shortening of the one diameter of the Earth would amount to about 6.5 cm. The length of a meter rod would change, when moved from one principal position into the other, by about  $\frac{1}{200}$  micron. One could hardly hope for success in trying to perceive such small quantities except by means of an interference method. We should have to operate with two perpendicular rods, and with two mutually interfering pencils of light, allowing the one to travel to and fro along the first rod, and the other along the second rod. But in this way we should come back once more to the Michelson experiment, and revolving the apparatus we should perceive no displacement of the fringes. Reversing a previous remark, we might now say that the displacement produced by the alterations of length is compensated by the Maxwell displacement.

4. It is worth noticing that we are led to just the same changes of dimensions as have been presumed above if we, *firstly*, without taking molecular movement into consideration, assume that in a solid body left to itself the forces, attractions or repulsions, acting upon any molecule maintain one another in equilibrium, and, *secondly*—though to be sure, there is no reason for doing so—if we apply to these molecular forces the law which in another place\* we deduced for

\* Viz., § 23 of the book, "Versuch einer Theorie der elektrischen und optischen Erscheinungen in bewegten Körpern."

electrostatic actions. For if we now understand by  $S_1$  and  $S_2$ , not, as formerly, two systems of charged particles, but two systems of molecules—the second at rest and the first moving with a velocity  $v$  in the direction of the axis of  $x$ —between the dimensions of which the relationship subsists as previously stated; and if we assume that in both systems the  $x$  components of the forces are the same, while the  $y$  and  $z$  components differ from one another by the factor  $\sqrt{1 - v^2/c^2}$ , then it is clear that the forces in  $S_1$  will be in equilibrium whenever they are so in  $S_2$ . If therefore  $S_2$  is the state of equilibrium of a solid body at rest, then the molecules in  $S_1$  have precisely those positions in which they can persist under the influence of translation. The displacement would naturally bring about this disposition of the molecules of its own accord, and thus effect a shortening in the direction of motion in the proportion of 1 to  $\sqrt{1 - v^2/c^2}$ , in accordance with the formulæ given in the above-mentioned paragraph. This leads to the values

$$\delta = -\frac{1}{2} \frac{v^2}{c^2}, \quad \epsilon = 0$$

in agreement with (1).

In reality the molecules of a body are not at rest, but in every "state of equilibrium" there is a stationary movement. What influence this circumstance may have in the phenomenon which we have been considering is a question which we do not here touch upon; in any case the experiments of Michelson and Morley, in consequence of unavoidable errors of observation, afford considerable latitude for the values of  $\delta$  and  $\epsilon$ .



**ELECTROMAGNETIC PHENOMENA IN  
A SYSTEM MOVING WITH ANY VELO-  
CITY LESS THAN THAT OF LIGHT**

**BY**

**H. A. LORENTZ**

*Reprinted from the English version in Proceedings of the  
Academy of Sciences of Amsterdam, 6, 1904.*



# ELECTROMAGNETIC PHENOMENA IN A SYSTEM MOVING WITH ANY VELOCITY LESS THAN THAT OF LIGHT

By H. A. LORENTZ

§ 1. **T**HE problem of determining the influence exerted on electric and optical phenomena by a translation, such as all systems have in virtue of the Earth's annual motion, admits of a comparatively simple solution, so long as only those terms need be taken into account, which are proportional to the first power of the ratio between the velocity of translation  $v$  and the velocity of light  $c$ . Cases in which quantities of the second order, i.e. of the order  $v^2/c^2$ , may be perceptible, present more difficulties. The first example of this kind is Michelson's well-known interference-experiment, the negative result of which has led Fitzgerald and myself to the conclusion that the dimensions of solid bodies are slightly altered by their motion through the ether.

Some new experiments, in which a second order effect was sought for, have recently been published. Rayleigh\* and Brace† have examined the question whether the Earth's motion may cause a body to become doubly refracting. At first sight this might be expected, if the just mentioned change of dimensions is admitted. Both physicists, however, have obtained a negative result.

In the second place Trouton and Noble ‡ have endeavoured to detect a turning couple acting on a charged condenser, the plates of which make a certain angle with the direction of translation. The theory of electrons, unless it be modified by some new hypothesis, would undoubtedly require the

\* Rayleigh, Phil. Mag. (6), 4, 1902, p. 678.

† Brace, Phil. Mag. (6), 7, 1904, p. 317.

‡ Trouton and Noble, Phil. Trans. Roy. Soc. Lond., A 202, 1903, p. 165.

## 12 ELECTROMAGNETIC PHENOMENA

existence of such a couple. In order to see this, it will suffice to consider a condenser with ether as dielectric. It may be shown that in every electrostatic system, moving with a velocity  $v$ ,\* there is a certain amount of "electromagnetic momentum." If we represent this, in direction and magnitude, by a vector  $\mathbf{G}$ , the couple in question will be determined by the vector product †

$$[\mathbf{G} \cdot \mathbf{v}] \quad . \quad . \quad . \quad . \quad . \quad (1)$$

Now, if the axis of  $z$  is chosen perpendicular to the condenser plates, the velocity  $\mathbf{v}$  having any direction we like; and if  $U$  is the energy of the condenser, calculated in the ordinary way, the components of  $\mathbf{G}$  are given ‡ by the following formulæ, which are exact up to the first order,

$$G_x = \frac{2U}{c^2} v_x, \quad G_y = \frac{2U}{c^2} v_y, \quad G_z = 0.$$

Substituting these values in (1), we get for the components of the couple, up to terms of the second order,

$$\frac{2U}{c^2} v_y v_z, \quad - \frac{2U}{c^2} v_x v_z, \quad 0.$$

These expressions show that the axis of the couple lies in the plane of the plates, perpendicular to the translation. If  $\alpha$  is the angle between the velocity and the normal to the plates, the moment of the couple will be  $U(v/c)^2 \sin 2\alpha$ ; it tends to turn the condenser into such a position that the plates are parallel to the Earth's motion.

In the apparatus of Trouton and Noble the condenser was fixed to the beam of a torsion-balance, sufficiently delicate to be deflected by a couple of the above order of magnitude. No effect could however be observed.

§ 2. The experiments of which I have spoken are not the only reason for which a new examination of the problems connected with the motion of the Earth is desirable. Poin-

\* A vector will be denoted by a Clarendon letter, its magnitude by the corresponding Latin letter.

† See my article: "Weiterbildung der Maxwell'schen Theorie. Electronentheorie," *Mathem. Encyclopädie*, V, 14, § 21, a. (This article will be quoted as "M.E.")

‡ "M.E.," § 56, c.

caré \* has objected to the existing theory of electric and optical phenomena in moving bodies that, in order to explain Michelson's negative result, the introduction of a new hypothesis has been required, and that the same necessity may occur each time new facts will be brought to light. Surely this course of inventing special hypotheses for each new experimental result is somewhat artificial. It would be more satisfactory if it were possible to show by means of certain fundamental assumptions and without neglecting terms of one order of magnitude or another, that many electromagnetic actions are entirely independent of the motion of the system. Some years ago, I already sought to frame a theory of this kind.† I believe it is now possible to treat the subject with a better result. The only restriction as regards the velocity will be that it be less than that of light.

§ 3. I shall start from the fundamental equations of the theory of electrons.‡ Let  $\mathbf{D}$  be the dielectric displacement in the ether,  $\mathbf{H}$  the magnetic force,  $\rho$  the volume-density of the charge of an electron,  $\mathbf{v}$  the velocity of a point of such a particle, and  $\mathbf{F}$  the ponderomotive force, i.e. the force, reckoned per unit charge, which is exerted by the ether on a volume-element of an electron. Then, if we use a fixed system of co-ordinates,

$$\left. \begin{aligned} \operatorname{div} \mathbf{D} &= \rho, \quad \operatorname{div} \mathbf{H} = 0, \\ \operatorname{curl} \mathbf{H} &= \frac{1}{c} \left( \frac{\partial \mathbf{D}}{\partial t} + \rho \mathbf{v} \right), \\ \operatorname{curl} \mathbf{D} &= - \frac{1}{c} \frac{\partial \mathbf{H}}{\partial t}, \\ \mathbf{F} &= \mathbf{D} + \frac{1}{c} [\mathbf{v} \cdot \mathbf{H}]. \end{aligned} \right\} \quad (2)$$

I shall now suppose that the system as a whole moves in the direction of  $x$  with a constant velocity  $v$ , and I shall denote by  $\mathbf{u}$  any velocity which a point of an electron may have in addition to this, so that

$$v_x = v + u_x, \quad v_y = u_y, \quad v_z = u_z.$$

\* Poincaré, *Rapports du Congrès de physique de 1900*, Paris, 1, pp. 22, 23.

† Lorentz, *Zittingsverslag Akad. v. Wet.*, 7, 1899, p. 507; *Amsterdam Proc.*, 1898-99, p. 427.

‡ "M.E.," § 2.



## 14 ELECTROMAGNETIC PHENOMENA

If the equations (2) are at the same time referred to axes moving with the system, they become

$$\operatorname{div} \mathbf{D} = \rho, \quad \operatorname{div} \mathbf{H} = 0,$$

$$\frac{\partial H_x}{\partial y} - \frac{\partial H_y}{\partial z} = \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) D_x + \frac{1}{c} \rho (v + u_x),$$

$$\frac{\partial H_x}{\partial z} - \frac{\partial H_z}{\partial x} = \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) D_y + \frac{1}{c} \rho u_y,$$

$$\frac{\partial H_y}{\partial x} - \frac{\partial H_x}{\partial y} = \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) D_z + \frac{1}{c} \rho u_z,$$

$$\frac{\partial D_x}{\partial y} - \frac{\partial D_y}{\partial z} = - \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) H_x,$$

$$\frac{\partial D_x}{\partial z} - \frac{\partial D_z}{\partial x} = - \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) H_y,$$

$$\frac{\partial D_y}{\partial x} - \frac{\partial D_x}{\partial y} = - \frac{1}{c} \left( \frac{\partial}{\partial t} - v \frac{\partial}{\partial x} \right) H_z,$$

$$F_x = D_x + \frac{1}{c} (u_y H_z - u_z H_y),$$

$$F_y = D_y - \frac{1}{c} v H_z + \frac{1}{c} (u_z H_x - u_x H_z),$$

$$F_z = D_z + \frac{1}{c} v H_y + \frac{1}{c} (u_x H_y - u_y H_x).$$

§ 4. We shall further transform these formulæ by a change of variables. Putting

$$\frac{c^2}{c^2 - v^2} = \beta^2, \quad . \quad . \quad . \quad (3)$$

and understanding by  $l$  another numerical quantity, to be determined further on, I take as new independent variables

$$x' = \beta l x, \quad y' = l y, \quad z' = l z, \quad . \quad . \quad . \quad (4)$$

$$t' = \frac{l}{\beta} t - \beta \frac{v}{c^2} x, \quad . \quad . \quad . \quad (5)$$

and I define two new vectors  $\mathbf{D}'$  and  $\mathbf{H}'$  by the formulæ

Here  $dS'$  is an element of the space  $S'$ ,  $r'$  its distance from  $P'$ , and the brackets serve to denote the quantity  $\rho'$  and the vector  $\rho' \mathbf{u}'$  such as they are in the element  $dS'$ , for the value  $t' - r'/c$  of the fourth independent variable.

Instead of (15) and (16) we may also write, taking into account (4) and (7),

$$\phi' = \frac{1}{4\pi} \int \frac{[\rho]}{r} dS \quad . \quad . \quad . \quad (17)$$

$$\mathbf{A}' = \frac{1}{4\pi c} \int \frac{[\rho \mathbf{u}]}{r} dS, \quad . \quad . \quad . \quad (18)$$

the integrations now extending over the electromagnetic system itself. It should be kept in mind that in these formulæ  $r'$  does not denote the distance between the element  $dS$  and the point  $(x, y, z)$  for which the calculation is to be performed. If the element lies at the point  $(x_1, y_1, z_1)$ , we must take

$$r' = l \sqrt{\beta^2 (x - x_1)^2 + (y - y_1)^2 + (z - z_1)^2}.$$

It is also to be remembered that, if we wish to determine  $\phi'$  and  $\mathbf{A}'$  for the instant at which the local time in  $P$  is  $t'$ , we must take  $\rho$  and  $\rho \mathbf{u}'$ , such as they are in the element  $dS$  at the instant at which the local time of that element is  $t' - r'/c$ .

§ 6. It will suffice for our purpose to consider two special cases. The first is that of an electrostatic system, i.e. a system having no other motion but the translation with the velocity  $v$ . In this case  $\mathbf{u}' = 0$ , and therefore, by (12),  $\mathbf{A}' = 0$ . Also,  $\phi'$  is independent of  $t'$ , so that the equations (11), (13), and (14) reduce to

$$\left. \begin{aligned} \nabla'^2 \phi' &= -\rho', \\ \mathbf{D}' &= -\text{grad}' \phi', \\ \mathbf{H}' &= 0 \end{aligned} \right\} \quad . \quad . \quad . \quad (19)$$

After having determined the vector  $\mathbf{D}'$  by means of these equations, we know also the ponderomotive force acting on electrons that belong to the system. For these the formulæ (10) become, since  $\mathbf{u}' = 0$ ,

$$F_x = l^2 D'_x, \quad F_y = \frac{l^2}{\beta} D'_y, \quad F_z = \frac{l^2}{\beta} D'_z \quad . \quad . \quad (20)$$

## 18 ELECTROMAGNETIC PHENOMENA

The result may be put in a simple form if we compare the moving system  $\Sigma$ , with which we are concerned, to another electrostatic system  $\Sigma'$  which remains at rest, and into which  $\Sigma$  is changed if the dimensions parallel to the axis of  $x$  are multiplied by  $\beta l$ , and the dimensions which have the direction of  $y$  or that of  $z$ , by  $l$ —a deformation for which  $(\beta l, l, l)$  is an appropriate symbol. In this new system, which we may suppose to be placed in the above-mentioned space  $S'$ , we shall give to the density the value  $\rho'$ , determined by (7), so that the charges of corresponding elements of volume and of corresponding electrons are the same in  $\Sigma$  and  $\Sigma'$ . Then we shall obtain the forces acting on the electrons of the moving system  $\Sigma$ , if we first determine the corresponding forces in  $\Sigma'$ , and next multiply their components in the direction of the axis of  $x$  by  $l^2$ , and their components perpendicular to that axis by  $\frac{l^2}{\beta}$ . This is conveniently expressed by the formula

$$F(\Sigma) = \left( l^2, \frac{l^2}{\beta}, \frac{l^2}{\beta} \right) F(\Sigma') \quad . \quad . \quad . \quad (21)$$

It is further to be remarked that, after having found  $D'$  by (19), we can easily calculate the electromagnetic momentum in the moving system, or rather its component in the direction of the motion. Indeed, the formula

$$\mathbf{G} = \frac{1}{c} \int [\mathbf{D} \cdot \mathbf{H}] dS$$

shows that

$$G_x = \frac{1}{c} \int (D_y H_z - D_z H_y) dS.$$

Therefore, by (6), since  $\mathbf{H}' = 0$

$$G_x = \frac{\beta^2 l^4 v}{c^2} \int (D_y'^2 + D_z'^2) dS = \frac{\beta l v}{c^2} \int (D_y'^2 + D_z'^2) dS'. \quad (22)$$

§ 7. Our second special case is that of a particle having an electric moment, i.e. a small space  $S$ , with a total charge  $\int \rho dS = 0$ , but with such a distribution of density that the

integrals  $\int \rho x dS$ ,  $\int \rho y dS$ ,  $\int \rho z dS$  have values differing from 0.

Let  $\xi$ ,  $\mu$ ,  $\zeta$  be the co-ordinates, taken relatively to a fixed point A of the particle, which may be called its centre, and let the electric moment be defined as a vector  $\mathbf{P}$  whose components are

$$P_x = \int \rho \xi dS, \quad P_y = \int \rho \eta dS, \quad P_z = \int \rho \zeta dS. \quad (23)$$

Then

$$\frac{dP_x}{dt} = \int \rho u_x dS, \quad \frac{dP_y}{dt} = \int \rho u_y dS, \quad \frac{dP_z}{dt} = \int \rho u_z dS. \quad (24)$$

Of course, if  $\xi$ ,  $\eta$ ,  $\zeta$  are treated as infinitely small,  $u_x$ ,  $u_y$ ,  $u_z$  must be so likewise. We shall neglect squares and products of these six quantities.

We shall now apply the equation (17) to the determination of the scalar potential  $\phi'$  for an exterior point P ( $x, y, z$ ), at a finite distance from the polarized particle, and for the instant at which the local time of this point has some definite value  $t'$ . In doing so, we shall give the symbol  $[\rho]$ , which, in (17), relates to the instant at which the local time in  $dS$  is  $t' - r'/c$ , a slightly different meaning. Distinguishing by  $r'_0$  the value of  $r'$  for the centre A, we shall understand by  $[\rho]$  the value of the density existing in the element  $dS$  at the point ( $\xi, \eta, \zeta$ ), at the instant  $t_0$  at which the local time of A is  $t' - r'_0/c$ .

It may be seen from (5) that this instant precedes that for which we have to take the numerator in (17) by

$$\beta^2 \frac{v\xi}{c^2} + \frac{\beta(r'_0 - r')}{lc} = \beta^2 \frac{v\xi}{c^2} + \frac{\beta}{lc} \left( \xi \frac{\partial r'}{\partial x} + \eta \frac{\partial r'}{\partial y} + \zeta \frac{\partial r'}{\partial z} \right)$$

units of time. In this last expression we may put for the differential coefficients their values at the point A.

In (17) we have now to replace  $[\rho]$  by

$$[\rho] + \beta^2 \frac{v\xi}{c^2} \left[ \frac{\partial \rho}{\partial t} \right] + \frac{\beta}{lc} \left( \xi \frac{\partial r'}{\partial x} + \eta \frac{\partial r'}{\partial y} + \zeta \frac{\partial r'}{\partial z} \right) \left[ \frac{\partial \rho}{\partial t} \right] \quad (25)$$

where  $\left[ \frac{\partial \rho}{\partial t} \right]$  relates again to the time  $t_0$ . Now, the value of  $t'$  for which the calculations are to be performed having been

## 20 ELECTROMAGNETIC PHENOMENA

chosen, this time  $t_0$  will be a function of the co-ordinates  $x, y, z$  of the exterior point P. The value of  $[\rho]$  will therefore depend on these co-ordinates in such a way that

$$\frac{\partial[\rho]}{\partial x} = -\frac{\beta}{lc} \frac{\partial r'}{\partial x} \left[ \frac{\partial \rho}{\partial t} \right], \text{ etc.}$$

by which (25) becomes

$$[\rho] + \beta^2 \frac{v^2}{c^2} \xi \left[ \frac{\partial \rho}{\partial t} \right] - \left( \xi \frac{\partial[\rho]}{\partial x} + \eta \frac{\partial[\rho]}{\partial y} + \zeta \frac{\partial[\rho]}{\partial z} \right).$$

Again, if henceforth we understand by  $r'$  what has above been called  $r'_0$ , the factor  $\frac{1}{r'}$  must be replaced by

$$\frac{1}{r'} - \xi \frac{\partial}{\partial x} \left( \frac{1}{r'} \right) - \eta \frac{\partial}{\partial y} \left( \frac{1}{r'} \right) - \zeta \frac{\partial}{\partial z} \left( \frac{1}{r'} \right),$$

so that after all, in the integral (17), the element  $dS$  is multiplied by

$$\frac{[\rho]}{r'} + \frac{\beta^2 v^2}{c^2 r'^2} \xi \left[ \frac{\partial \rho}{\partial t} \right] - \frac{\partial}{\partial x} \frac{\xi[\rho]}{r'} - \frac{\partial}{\partial y} \frac{\eta[\rho]}{r'} - \frac{\partial}{\partial z} \frac{\zeta[\rho]}{r'}.$$

This is simpler than the primitive form, because neither  $r'$ , nor the time for which the quantities enclosed in brackets are to be taken, depend on  $x, y, z$ . Using (23) and remembering that  $\int \rho dS = 0$ , we get

$$\phi' = \frac{\beta^2 v}{4pc^2 r'} \left[ \frac{\partial P_x}{\partial t} \right] - \frac{1}{4p} \left\{ \frac{\partial}{\partial x} \frac{[P_x]}{r'} + \frac{\partial}{\partial y} \frac{[P_y]}{r'} + \frac{\partial}{\partial z} \frac{[P_z]}{r'} \right\},$$

a formula in which all the enclosed quantities are to be taken for the instant at which the local time of the centre of the particle is  $t' = r'/c$ .

We shall conclude these calculations by introducing a new vector  $P'$ , whose components are

$$P'_x = \beta l P_x, \quad P'_y = l P_y, \quad P'_z = l P_z, \quad . \quad . \quad . \quad (26)$$

passing at the same time to  $x', y', z', t'$  as independent variables. The final result is

$$\phi' = \frac{v}{4pc^2 r'} \frac{\partial [P'_x]}{\partial t'} - \frac{1}{4p} \left\{ \frac{\partial}{\partial x'} \frac{[P'_x]}{r'} + \frac{\partial}{\partial y'} \frac{[P'_y]}{r'} + \frac{\partial}{\partial z'} \frac{[P'_z]}{r'} \right\}.$$

As to the formula (18) for the vector potential, its transformation is less complicated, because it contains the infinitely small vector  $\mathbf{u}'$ . Having regard to (8), (24), (26), and (5), I find

$$\mathbf{A}' = \frac{1}{4\pi cr'} \frac{\partial [\mathbf{P}']}{\partial t'}.$$

The field produced by the polarized particle is now wholly determined. The formula (13) leads to

$$\mathbf{D}' = -\frac{1}{4\pi c^2} \frac{\partial^2 [\mathbf{P}']}{\partial t'^2} + \frac{1}{4\pi} \text{grad}' \left\{ \frac{\partial [\mathbf{P}'_x]}{\partial x'} \frac{1}{r'} + \frac{\partial [\mathbf{P}'_y]}{\partial y'} \frac{1}{r'} + \frac{\partial [\mathbf{P}'_z]}{\partial z'} \frac{1}{r'} \right\} \quad (27)$$

and the vector  $\mathbf{H}'$  is given by (14). We may further use the equations (20), instead of the original formulæ (10), if we wish to consider the forces exerted by the polarized particle on a similar one placed at some distance. Indeed, in the second particle, as well as in the first, the velocities  $\mathbf{u}$  may be held to be infinitely small.

It is to be remarked that the formulæ for a system without translation are implied in what precedes. For such a system the quantities with accents become identical to the corresponding ones without accents; also  $\beta = 1$  and  $l = 1$ . The components of (27) are at the same time those of the electric force which is exerted by one polarized particle on another.

§ 8. Thus far we have used only the fundamental equations without any new assumptions. I shall now suppose *that the electrons, which I take to be spheres of radius  $R$  in the state of rest, have their dimensions changed by the effect of a translation, the dimensions in the direction of motion becoming  $\beta l$  times and those in perpendicular directions  $l$  times smaller.*

In this deformation, which may be represented by  $\left( \frac{1}{\beta l}, \frac{1}{l}, \frac{1}{l} \right)$ , each element of volume is understood to preserve its charge.

Our assumption amounts to saying that in an electrostatic system  $\Sigma$ , moving with a velocity  $v$ , all electrons are flattened ellipsoids with their smaller axes in the direction of

## 22 ELECTROMAGNETIC PHENOMENA

motion. If now, in order to apply the theorem of § 6, we subject the system to the deformation  $(\beta l, l, l)$ , we shall have again spherical electrons of radius  $R$ . Hence, if we alter the relative position of the centres of the electrons in  $\Sigma$  by applying the deformation  $(\beta l, l, l)$ , and if, in the points thus obtained, we place the centres of electrons that remain at rest, we shall get a system, identical to the imaginary system  $\Sigma'$ , of which we have spoken in § 6. The forces in this system and those in  $\Sigma$  will bear to each other the relation expressed by (21).

In the second place I shall suppose *that the forces between uncharged particles, as well as those between such particles and electrons, are influenced by a translation in quite the same way as the electric forces in an electrostatic system*. In other terms, whatever be the nature of the particles composing a ponderable body, so long as they do not move relatively to each other, we shall have between the forces acting in a system  $(\Sigma')$  without, and the same system  $(\Sigma)$  with a translation, the relation specified in (21), if, as regards the relative position of the particles,  $\Sigma'$  is got from  $\Sigma$  by the deformation  $(\beta l, l, l)$ , or  $\Sigma$  from  $\Sigma'$  by the deformation  $\left(\frac{1}{\beta l}, \frac{1}{l}, \frac{1}{l}\right)$ .

We see by this that, as soon as the resulting force is zero for a particle in  $\Sigma'$ , the same must be true for the corresponding particle in  $\Sigma$ . Consequently, if, neglecting the effects of molecular motion, we suppose each particle of a solid body to be in equilibrium under the action of the attractions and repulsions exerted by its neighbours, and if we take for granted that there is but one configuration of equilibrium, we may draw the conclusion that the system  $\Sigma'$ , if the velocity  $v$  is imparted to it, will *of itself* change into the system  $\Sigma$ . In other terms, the translation will *produce* the deformation  $\left(\frac{1}{\beta l}, \frac{1}{l}, \frac{1}{l}\right)$ .

The case of molecular motion will be considered in § 12.

It will easily be seen that the hypothesis which was formerly advanced in connexion with Michelson's experiment, is implied in what has now been said. However, the present hypothesis is more general, because the only

limitation imposed on the motion is that its velocity be less than that of light.

§ 9. We are now in a position to calculate the electromagnetic momentum of a single electron. For simplicity's sake I shall suppose the charge  $e$  to be uniformly distributed over the surface, so long as the electron remains at rest. Then a distribution of the same kind will exist in the system  $\Sigma'$  with which we are concerned in the last integral of (22). Hence

$$\int (D'^2_y + D'^2_z) dS' = \frac{2}{3} \int D'^2 dS' = \frac{e^2}{6\pi} \int_R \frac{dr}{r^2} = \frac{e^2}{6\pi R},$$

and

$$G_x = \frac{e^2}{6\pi c^2 R} \beta l v.$$

It must be observed that the product  $\beta l$  is a function of  $v$  and that, for reasons of symmetry, the vector  $\mathbf{G}$  has the direction of the translation. In general, representing by  $\mathbf{v}$  the velocity of this motion, we have the vector equation

$$\mathbf{G} = \frac{e^2}{6\pi c^2 R} \beta l \mathbf{v} \quad . \quad . \quad . \quad (28)$$

Now, every change in the motion of a system will entail a corresponding change in the electromagnetic momentum and will therefore require a certain force, which is given in direction and magnitude by

$$\mathbf{F} = \frac{d\mathbf{G}}{dt} \quad . \quad . \quad . \quad (29)$$

Strictly speaking, the formula (28) may only be applied in the case of a uniform rectilinear translation. On account of this circumstance—though (29) is always true—the theory of rapidly varying motions of an electron becomes very complicated, the more so, because the hypothesis of § 8 would imply that the direction and amount of the deformation are continually changing. It is, indeed, hardly probable that the form of the electron will be determined solely by the velocity existing at the moment considered.

Nevertheless, provided the changes in the state of motion



## 24 ELECTROMAGNETIC PHENOMENA

be sufficiently slow, we shall get a satisfactory approximation by using (28) at every instant. The application of (29) to such a *quasi-stationary* translation, as it has been called by Abraham,\* is a very simple matter. Let, at a certain instant,  $\mathbf{a}_1$  be the acceleration in the direction of the path, and  $\mathbf{a}_2$  the acceleration perpendicular to it. Then the force  $\mathbf{F}$  will consist of two components, having the directions of these accelerations and which are given by

$$\mathbf{F}_1 = m_1 \mathbf{a}_1 \text{ and } \mathbf{F}_2 = m_2 \mathbf{a}_2,$$

if

$$m_1 = \frac{e^2}{6\pi c^2 R} \frac{d(\beta l v)}{dv} \text{ and } m_2 = \frac{e^2}{6\pi c^2 R} \beta l. \quad (30)$$

Hence, in phenomena in which there is an acceleration in the direction of motion, the electron behaves as if it had a mass  $m_1$ ; in those in which the acceleration is normal to the path, as if the mass were  $m_2$ . These quantities  $m_1$  and  $m_2$  may therefore properly be called the "longitudinal" and "transverse" electromagnetic masses of the electron. I shall suppose *that there is no other, no "true" or "material" mass.*

Since  $\beta$  and  $l$  differ from unity by quantities of the order  $v^2/c^2$ , we find for very small velocities

$$m_1 = m_2 = \frac{e^2}{6\pi c^2 R}.$$

This is the mass with which we are concerned, if there are small vibratory motions of the electrons in a system without translation. If, on the contrary, motions of this kind are going on in a body moving with the velocity  $v$  in the direction of the axis of  $x$ , we shall have to reckon with the mass  $m_1$ , as given by (30), if we consider the vibrations parallel to that axis, and with the mass  $m_2$ , if we treat of those that are parallel to OY or OZ. Therefore, in short terms, referring by the index  $\Sigma$  to a moving system and by  $\Sigma'$  to one that remains at rest,

$$m(\Sigma) = \left( \frac{d(\beta l v)}{dv}, \beta l, \beta l \right) m(\Sigma') \quad (31)$$

\* Abraham, Wied. Ann., 10, 1908, p. 105.

§ 10. We can now proceed to examine the influence of the Earth's motion on optical phenomena in a system of transparent bodies. In discussing this problem we shall fix our attention on the variable electric moments in the particles or "atoms" of the system. To these moments we may apply what has been said in § 7. For the sake of simplicity we shall suppose that, in each particle, the charge is concentrated in a certain number of separate electrons, and that the "elastic" forces that act on one of these, and, conjointly with the electric forces, determine its motion, have their origin within the bounds of the *same* atom.

I shall show that, if we start from any given state of motion in a system without translation, we may deduce from it a corresponding state that can exist in the same system after a translation has been imparted to it, the kind of correspondence being as specified in what follows.

(a) Let  $A'_1, A'_2, A'_3$ , etc., be the centres of the particles in the system without translation ( $\Sigma'$ ); neglecting molecular motions we shall assume these points to remain at rest. The system of points  $A_1, A_2, A_3$ , etc., formed by the centres of the particles in the moving system  $\Sigma$ , is obtained from  $A'_1, A'_2, A'_3$ , etc., by means of a deformation  $\left(\frac{1}{\beta l}, \frac{1}{l}, \frac{1}{l}\right)$ . According to what has been said in § 8, the centres will of themselves take these positions  $A'_1, A'_2, A'_3$ , etc., if originally, before there was a translation, they occupied the positions  $A_1, A_2, A_3$ , etc.

We may conceive any point  $P'$  in the space of the system  $\Sigma'$  to be displaced by the above deformation, so that a definite point  $P$  of  $\Sigma$  corresponds to it. For two corresponding points  $P'$  and  $P$  we shall define corresponding instants, the one belonging to  $P'$ , the other to  $P$ , by stating that the true time at the first instant is equal to the local time, as determined by (5) for the point  $P$ , at the second instant. By corresponding times for two corresponding *particles* we shall understand times that may be said to correspond, if we fix our attention on the *centres*  $A'$  and  $A$  of these particles.

(b) As regards the interior state of the atoms, we shall assume that the configuration of a particle  $A$  in  $\Sigma$  at a certain

## 26 ELECTROMAGNETIC PHENOMENA

time may be derived by means of the deformation  $\left(\frac{1}{\beta t}, \frac{1}{t}, \frac{1}{t}\right)$  from the configuration of the corresponding particle in  $\Sigma'$ , such as it is at the corresponding instant. In so far as this assumption relates to the form of the electrons themselves, it is implied in the first hypothesis of § 8.

Obviously, if we start from a state really existing in the system  $\Sigma'$ , we have now completely defined a state of the moving system  $\Sigma$ . The question remains, however, whether this state will likewise be a possible one.

In order to judge of this, we may remark in the first place that the electric moments which we have supposed to exist in the moving system and which we shall denote by  $\mathbf{P}$ , will be certain definite functions of the co-ordinates  $x, y, z$  of the centres  $A$  of the particles, or, as we shall say, of the co-ordinates of the particles themselves, and of the time  $t$ . The equations which express the relations between  $\mathbf{P}$  on one hand and  $x, y, z, t$  on the other, may be replaced by other equations containing the vectors  $\mathbf{P}'$  defined by (26) and the quantities  $x', y', z', t'$  defined by (4) and (5). Now, by the above assumptions  $a$  and  $b$ , if in a particle  $A$  of the moving system, whose co-ordinates are  $x, y, z$ , we find an electric moment  $\mathbf{P}$  at the time  $t$ , or at the local time  $t'$ , the vector  $\mathbf{P}'$  given by (26) will be the moment which exists in the other system at the true time  $t'$  in a particle whose co-ordinates are  $x', y', z'$ . It appears in this way that the equations between  $\mathbf{P}', x', y', z', t'$  are the same for both systems, the difference being only this, that for the system  $\Sigma'$  without translation these symbols indicate the moment, the co-ordinates, and the true time, whereas their meaning is different for the moving system,  $\mathbf{P}', x', y', z', t'$  being here related to the moment  $\mathbf{P}$ , the co-ordinates  $x, y, z$  and the general time  $t$  in the manner expressed by (26), (4), and (5).

It has already been stated that the equation (27) applies to both systems. The vector  $\mathbf{D}'$  will therefore be the same in  $\Sigma'$  and  $\Sigma$ , provided we always compare corresponding places and times. However, this vector has not the same meaning in the two cases. In  $\Sigma'$  it represents the electric force, in  $\Sigma$  it is related to this force in the way expressed by (20). We may therefore conclude that the ponderomotive

forces acting, in  $\Sigma$  and in  $\Sigma'$ , on corresponding particles at corresponding instants, bear to each other the relation determined by (21). In virtue of our assumption (b), taken in connexion with the second hypothesis of § 8, the same relation will exist between the "elastic" forces; consequently, the formula (21) may also be regarded as indicating the relation between the total forces, acting on corresponding electrons, at corresponding instants.

It is clear that the state we have supposed to exist in the moving system will really be possible if, in  $\Sigma$  and  $\Sigma'$ , the products of the mass  $m$  and the acceleration of an electron are to each other in the same relation as the forces, i.e. if

$$ma(\Sigma) = \left( l^2, \frac{l^2}{\beta}, \frac{l^2}{\beta} \right) ma(\Sigma'). \quad (32)$$

Now, we have for the accelerations

$$a(\Sigma) = \left( \frac{l}{\beta^3}, \frac{l}{\beta^3}, \frac{l}{\beta^3} \right) a(\Sigma'). \quad (33)$$

as may be deduced from (4) and (5), and combining this with (32), we find for the masses

$$m(\Sigma) = (\beta^3 l, \beta l, \beta l) m(\Sigma').$$

If this is compared with (31), it appears that, whatever be the value of  $l$ , the condition is always satisfied, as regards the masses with which we have to reckon when we consider vibrations perpendicular to the translation. The only condition we have to impose on  $l$  is therefore

$$\frac{d(\beta l v)}{dv} = \beta^3 l.$$

But, on account of (3),

$$\frac{d(\beta v)}{dv} = \beta^3,$$

so that we must put

$$\frac{dl}{dv} = 0, \quad l = \text{const.}$$

The value of the constant must be unity, because we know already that, for  $v = 0$ ,  $l = 1$ .

## 28 ELECTROMAGNETIC PHENOMENA

We are therefore led to suppose *that the influence of a translation on the dimensions (of the separate electrons and of a ponderable body as a whole) is confined to those that have the direction of the motion, these becoming  $\beta$  times smaller than they are in the state of rest.* If this hypothesis is added to those we have already made, we may be sure that two states, the one in the moving system, the other in the same system while at rest, corresponding as stated above, may both be possible. Moreover, this correspondence is not limited to the electric moments of the particles. In corresponding points that are situated either in the ether between the particles, or in that surrounding the ponderable bodies, we shall find at corresponding times the same vector  $\mathbf{D}'$  and, as is easily shown, the same vector  $\mathbf{H}'$ . We may sum up by saying: If, in the system without translation, there is a state of motion in which, at a definite place, the components of  $\mathbf{P}$ ,  $\mathbf{D}$ , and  $\mathbf{H}$  are certain functions of the time, then the same system after it has been put in motion (and thereby deformed) can be the seat of a state of motion in which, at the corresponding place, the components of  $\mathbf{P}'$ ,  $\mathbf{D}'$ , and  $\mathbf{H}'$  are the same functions of the local time.

There is one point which requires further consideration. The values of the masses  $m_1$  and  $m_2$  having been deduced from the theory of quasi-stationary motion, the question arises, whether we are justified in reckoning with them in the case of the rapid vibrations of light. Now it is found on closer examination that the motion of an electron may be treated as quasi-stationary if it changes very little during the time a light-wave takes to travel over a distance equal to the diameter. This condition is fulfilled in optical phenomena, because the diameter of an electron is extremely small in comparison with the wave-length.

§ 11. It is easily seen that the proposed theory can account for a large number of facts.

Let us take in the first place the case of a system without translation, in some parts of which we have continually  $\mathbf{P} = 0$ ,  $\mathbf{D} = 0$ ,  $\mathbf{H} = 0$ . Then, in the corresponding state for the moving system, we shall have in corresponding parts (or, as we may say, in the same parts of the deformed system)  $\mathbf{P}' = 0$ ,  $\mathbf{D}' = 0$ ,  $\mathbf{H}' = 0$ . These equations implying  $\mathbf{P} = 0$ ,

$D = 0$ ,  $H = 0$ , as is seen by (26) and (6), it appears that those parts which are dark while the system is at rest, will remain so after it has been put in motion. It will therefore be impossible to detect an influence of the Earth's motion on any optical experiment, made with a terrestrial source of light, in which the geometrical distribution of light and darkness is observed. Many experiments on interference and diffraction belong to this class.

In the second place, if, in two points of a system, rays of light of the same state of polarization are propagated in the same direction, the ratio between the amplitudes in these points may be shown not to be altered by a translation. The latter remark applies to those experiments in which the intensities in adjacent parts of the field of view are compared.

The above conclusions confirm the results which I formerly obtained by a similar train of reasoning, in which, however, the terms of the second order were neglected. They also contain an explanation of Michelson's negative result, more general than the one previously given, and of a somewhat different form; and they show why Rayleigh and Brace could find no signs of double refraction produced by the motion of the Earth.

As to the experiments of Trouton and Noble, their negative result becomes at once clear, if we admit the hypotheses of § 8. It may be inferred from these and from our last assumption (§ 10) that the only effect of the translation must have been a contraction of the whole system of electrons and other particles constituting the charged condenser and the beam and thread of the torsion-balance. Such a contraction does not give rise to a sensible change of direction.

It need hardly be said that the present theory is put forward with all due reserve. Though it seems to me that it can account for all well-established facts, it leads to some consequences that cannot as yet be put to the test of experiment. One of these is that the result of Michelson's experiment must remain negative, if the interfering rays of light are made to travel through some ponderable transparent body.

Our assumption about the contraction of the electrons

## 30 ELECTROMAGNETIC PHENOMENA

cannot in itself be pronounced to be either plausible or inadmissible. What we know about the nature of electrons is very little, and the only means of pushing our way farther will be to test such hypotheses as I have here made. Of course, there will be difficulties, e.g. as soon as we come to consider the rotation of electrons. Perhaps we shall have to suppose that in those phenomena in which, if there is no translation, spherical electrons rotate about a diameter, the points of the electrons in the moving system will describe elliptic paths, corresponding, in the manner specified in § 10, to the circular paths described in the other case.

§ 12. There remain to be said a few words about molecular motion. We may conceive that bodies in which this has a sensible influence or even predominates, undergo the same deformation as the systems of particles of constant relative position of which alone we have spoken till now. Indeed, in two systems of molecules  $\Sigma'$  and  $\Sigma$ , the first without and the second with a translation, we may imagine molecular motions corresponding to each other in such a way that, if a particle in  $\Sigma'$  has a certain position at a definite instant, a particle in  $\Sigma$  occupies at the corresponding instant the corresponding position. This being assumed, we may use the relation (33) between the accelerations in all those cases in which the velocity of molecular motion is very small as compared with  $v$ . In these cases the molecular forces may be taken to be determined by the relative positions, independently of the velocities of molecular motion. If, finally, we suppose these forces to be limited to such small distances that, for particles acting on each other, the difference of local times may be neglected, one of the particles, together with those which lie in its sphere of attraction or repulsion, will form a system which undergoes the often mentioned deformation. In virtue of the second hypothesis of § 8 we may therefore apply to the resulting molecular force acting on a particle, the equation (21). Consequently, the proper relation between the forces and the accelerations will exist in the two cases, if we suppose *that the masses of all particles are influenced by a translation to the same degree as the electromagnetic masses of the electrons.*

§ 13. The values (30), which I have found for the longi-

tudinal and transverse masses of an electron, expressed in terms of its velocity, are not the same as those that had been previously obtained by Abraham. The ground for this difference is to be sought solely in the circumstance that, in his theory, the electrons are treated as spheres of invariable dimensions. Now, as regards the transverse mass, the results of Abraham have been confirmed in a most remarkable way by Kaufmann's measurements of the deflexion of radium-rays in electric and magnetic fields. Therefore, if there is not to be a most serious objection to the theory I have now proposed, it must be possible to show that those measurements agree with my values nearly as well as with those of Abraham.

I shall begin by discussing two of the series of measurements published by Kaufmann\* in 1902. From each series he has deduced two quantities  $\eta$  and  $\zeta$ , the "reduced" electric and magnetic deflexions, which are related as follows to the ratio  $\gamma = v/c$  :—

$$\gamma = k_1 \frac{\zeta}{\eta}, \quad \psi(\gamma) = \frac{\eta}{k_2 \zeta^2} . \quad . \quad . \quad (34)$$

Here  $\psi(\gamma)$  is such a function, that the transverse mass is given by

$$m_2 = \frac{3}{4} \cdot \frac{e^2}{6\pi c^2 R} \psi(\gamma), \quad . \quad . \quad . \quad (35)$$

whereas  $k_1$  and  $k_2$  are constant in each series.

It appears from the second of the formulæ (30) that my theory leads likewise to an equation of the form (35); only Abraham's function  $\psi(\gamma)$  must be replaced by

$$\frac{4}{3}\beta = \frac{4}{3}(1 - \gamma^2)^{-1/2}.$$

Hence, my theory requires that, if we substitute this value for  $\psi(\gamma)$  in (34), these equations shall still hold. Of course, in seeking to obtain a good agreement, we shall be justified in giving to  $k_1$  and  $k_2$  other values than those of Kaufmann, and in taking for every measurement a proper value of the velocity  $v$ , or of the ratio  $\gamma$ . Writing  $sk_1, \frac{3}{4}k_2$ ,

\* Kaufmann, Physik. Zeitschr., 4, 1902, p. 55.



## 32 ELECTROMAGNETIC PHENOMENA

and  $\gamma'$  for the new values, we may put (34) in the form

$$\gamma' = s k_1 \frac{\zeta}{\eta} \quad . \quad . \quad . \quad (36)$$

and

$$(1 - \gamma'^2)^{-1/2} = \frac{\eta}{k_2' \zeta^2} \quad . \quad . \quad . \quad (37)$$

Kaufmann has tested his equations by choosing for  $k_1$  such a value that, calculating  $\gamma$  and  $k_2$  by means of (34), he obtained values for this latter number which, as well as might be, remained constant in each series. This constancy was the proof of a sufficient agreement.

I have followed a similar method, using, however, some of the numbers calculated by Kaufmann. I have computed for each measurement the value of the expression

$$k_2' = (1 - \gamma'^2)^{1/2} \psi(\gamma) k_2, \quad . \quad . \quad . \quad (38)$$

that may be got from (37) combined with the second of the equations (34). The values of  $\psi(\gamma)$  and  $k_2$  have been taken from Kaufmann's tables, and for  $\gamma'$  I have substituted the value he has found for  $\gamma$ , multiplied by  $s$ , the latter coefficient being chosen with a view to obtaining a good constancy of (38). The results are contained in the tables on opposite page, corresponding to the Tables III and IV in Kaufmann's paper.

The constancy of  $k_2'$  is seen to come out no less satisfactorily than that of  $k_2$ , the more so as in each case the value of  $s$  has been determined by means of only two measurements. The coefficient has been so chosen that for these two observations, which were in Table III the first and the last but one, and in Table IV the first and the last, the values of  $k_2'$  should be proportional to those of  $k_2$ .

I shall next consider two series from a later publication by Kaufmann,\* which have been calculated by Runge† by means of the method of least squares, the coefficients  $k_1$  and  $k_2$  having been determined in such a way that the values of  $\eta$ , calculated, for each observed  $\zeta$ , from Kaufmann's equations (34), agree as closely as may be with the observed values of  $\eta$ .

\* Kaufmann, Gött. Nachr. Math. phys. Kl., 1903, p. 90.

† Runge, *ibid.*, p. 326.

III.  $s = 0.983$ .

$\gamma$ .	$\psi(\gamma)$ .	$k_2$ .	$\gamma'$ .	$k'_2$ .
0.851	2.147	1.721	0.794	2.246
0.766	1.86	1.786	0.715	2.258
0.727	1.78	1.725	0.678	2.256
0.6615	1.66	1.727	0.617	2.256
0.6075	1.595	1.655	0.567	2.175

IV.  $s = 0.954$ .

$\gamma$ .	$\psi(\gamma)$ .	$k_2$ .	$\gamma'$ .	$k'_2$ .
0.963	3.28	8.12	0.919	10.86
0.949	2.86	7.99	0.905	9.70
0.933	2.73	7.46	0.890	9.28
0.883	2.31	8.32	0.842	10.86
0.860	2.195	8.09	0.820	10.15
0.880	2.06	8.18	0.792	10.28
0.801	1.96	8.18	0.764	10.28
0.777	1.89	8.04	0.741	10.20
0.752	1.83	8.02	0.717	10.22
0.732	1.785	7.97	0.698	10.18

I have determined by the same condition, likewise using the method of least squares, the constants  $a$  and  $b$  in the formula

$$\eta^2 = a\zeta^2 + b\zeta^4,$$

which may be deduced from my equations (36) and (37). Knowing  $a$  and  $b$ , I find  $\gamma$  for each measurement by means of the relation

$$\gamma = \sqrt{a} \frac{\zeta}{\eta}.$$

For two plates on which Kaufmann had measured the electric and magnetic deflexions, the results are as follows (p. 34), the deflexions being given in centimetres.

I have not found time for calculating the other tables in Kaufmann's paper. As they begin, like the table for Plate 15 (next page) with a rather large negative difference between the values of  $\eta$  which have been deduced from the observations and calculated by Runge, we may expect a satisfactory agreement with my formulæ.

# 84 ELECTROMAGNETIC PHENOMENA

Plate No. 15.  $\alpha = 0.06489$ ,  $b = 0.8039$ .

$\zeta$	$\eta$					$\gamma$	
	Observed.	Calculated by R.	Diff.	Calculated by L.	Diff.	Calculated by	
						R.	L.
0.1495	0.0388	0.0404	- 16	0.0400	- 12	0.987	0.951
0.199	0.0548	0.0550	- 2	0.0552	- 4	0.984	0.918
0.2475	0.0716	0.0710	+ 6	0.0715	+ 1	0.980	0.881
0.296	0.0896	0.0887	+ 9	0.0895	+ 1	0.889	0.842
0.3485	0.1080	0.1081	- 1	0.1090	- 10	0.847	0.803
0.391	0.1290	0.1297	- 7	0.1305	- 15	0.804	0.763
0.437	0.1524	0.1527	- 3	0.1532	- 8	0.768	0.727
0.4 25	0.1788	0.1777	+ 11	0.1777	+ 11	0.724	0.692
0.5265	0.2089	0.2089	- 6	0.2089	0	0.688	0.660

Plate No. 19.  $\alpha = 0.05867$ ,  $b = 0.2591$ .

$\zeta$	$\eta$					$\gamma$	
	Observed.	Calculated by R.	Diff.	Calculated by L.	Diff.	Calculated by	
						R.	L.
0.1495	0.0404	0.0388	+ 16	0.0379	+ 25	0.990	0.954
0.199	0.0529	0.0527	+ 2	0.0522	+ 7	0.969	0.923
0.247	0.0678	0.0675	+ 3	0.0674	+ 4	0.989	0.888
0.296	0.0834	0.0842	- 8	0.0844	- 10	0.902	0.849
0.3485	0.1019	0.1022	- 3	0.1026	- 7	0.862	0.811
0.391	0.1219	0.1222	- 3	0.1226	- 7	0.822	0.773
0.437	0.1429	0.1434	- 5	0.1437	- 8	0.782	0.736
0.4825	0.1660	0.1665	- 5	0.1664	- 4	0.744	0.702
0.5265	0.1916	0.1906	+ 10	0.1902	+ 14	0.709	0.671

# ON THE ELECTRODYNAMICS OF MOVING BODIES

BY

A. EINSTEIN

*Translated from "Zur Elektrodynamik bewegter Körper,"  
Annalen der Physik, 17, 1905.*



## ON THE ELECTRODYNAMICS OF MOVING BODIES

By A. EINSTEIN

IT is known that Maxwell's electrodynamics—as usually understood at the present time—when applied to moving bodies, leads to asymmetries which do not appear to be inherent in the phenomena. Take, for example, the reciprocal electrodynamic action of a magnet and a conductor. The observable phenomenon here depends only on the relative motion of the conductor and the magnet, whereas the customary view draws a sharp distinction between the two cases in which either the one or the other of these bodies is in motion. For if the magnet is in motion and the conductor at rest, there arises in the neighbourhood of the magnet an electric field with a certain definite energy, producing a current at the places where parts of the conductor are situated. But if the magnet is stationary and the conductor in motion, no electric field arises in the neighbourhood of the magnet. In the conductor, however, we find an electromotive force, to which in itself there is no corresponding energy, but which gives rise—assuming equality of relative motion in the two cases discussed—to electric currents of the same path and intensity as those produced by the electric forces in the former case.

Examples of this sort, together with the unsuccessful attempts to discover any motion of the earth relatively to the "light medium," suggest that the phenomena of electrodynamics as well as of mechanics possess no properties corresponding to the idea of absolute rest. They suggest rather that, as has already been shown to the first order of small quantities, the same laws of electrodynamics and optics will be valid for all frames of reference for which the equations of

mechanics hold good.\* We will raise this conjecture (the purport of which will hereafter be called the "Principle of Relativity") to the status of a postulate, and also introduce another postulate, which is only apparently irreconcilable with the former, namely, that light is always propagated in empty space with a definite velocity  $c$  which is independent of the state of motion of the emitting body. These two postulates suffice for the attainment of a simple and consistent theory of the electrodynamics of moving bodies based on Maxwell's theory for stationary bodies. The introduction of a "luminiferous ether" will prove to be superfluous inasmuch as the view here to be developed will not require an "absolutely stationary space" provided with special properties, nor assign a velocity-vector to a point of the empty space in which electromagnetic processes take place.

The theory to be developed is based—like all electrodynamics—on the kinematics of the rigid body, since the assertions of any such theory have to do with the relationships between rigid bodies (systems of co-ordinates), clocks, and electromagnetic processes. Insufficient consideration of this circumstance lies at the root of the difficulties which the electrodynamics of moving bodies at present encounters.

## I. KINEMATICAL PART

### § 1. Definition of Simultaneity

Let us take a system of co-ordinates in which the equations of Newtonian mechanics hold good.† In order to render our presentation more precise and to distinguish this system of co-ordinates verbally from others which will be introduced hereafter, we call it the "stationary system."

If a material point is at rest relatively to this system of co-ordinates, its position can be defined relatively thereto by the employment of rigid standards of measurement and the methods of Euclidean geometry, and can be expressed in Cartesian co-ordinates.

If we wish to describe the *motion* of a material point, we

\* The preceding memoir by Lorentz was not at this time known to the author.

† i.e. to the first approximation.

give the values of its co-ordinates as functions of the time. Now we must bear carefully in mind that a mathematical description of this kind has no physical meaning unless we are quite clear as to what we understand by "time." We have to take into account that all our judgments in which time plays a part are always judgments of *simultaneous events*. If, for instance, I say, "That train arrives here at 7 o'clock," I mean something like this: "The pointing of the small hand of my watch to 7 and the arrival of the train are simultaneous events." \*

It might appear possible to overcome all the difficulties attending the definition of "time" by substituting "the position of the small hand of my watch" for "time." And in fact such a definition is satisfactory when we are concerned with defining a time exclusively for the place where the watch is located; but it is no longer satisfactory when we have to connect in time series of events occurring at different places, or—what comes to the same thing—to evaluate the times of events occurring at places remote from the watch.

We might, of course, content ourselves with time values determined by an observer stationed together with the watch at the origin of the co-ordinates, and co-ordinating the corresponding positions of the hands with light signals, given out by every event to be timed, and reaching him through empty space. But this co-ordination has the disadvantage that it is not independent of the standpoint of the observer with the watch or clock, as we know from experience. We arrive at a much more practical determination along the following line of thought.

If at the point A of space there is a clock, an observer at A can determine the time values of events in the immediate proximity of A by finding the positions of the hands which are simultaneous with these events. If there is at the point B of space another clock in all respects resembling the one at A, it is possible for an observer at B to determine the time values of events in the immediate neighbourhood of B. But it is not possible without further assumption to compare, in

\* We shall not here discuss the inexactitude which lurks in the concept of simultaneity of two events at approximately the same place, which can only be removed by an abstraction.



respect of time, an event at A with an event at B. We have so far defined only an "A time" and a "B time." We have not defined a common "time" for A and B, for the latter cannot be defined at all unless we establish *by definition* that the "time" required by light to travel from A to B equals the "time" it requires to travel from B to A. Let a ray of light start at the "A time"  $t_A$  from A towards B, let it at the "B time"  $t_B$  be reflected at B in the direction of A, and arrive again at A at the "A time"  $t'_A$ .

In accordance with definition the two clocks synchronize if

$$t_B - t_A = t'_A - t_B.$$

We assume that this definition of synchronism is free from contradictions, and possible for any number of points; and that the following relations are universally valid:—

1. If the clock at B synchronizes with the clock at A, the clock at A synchronizes with the clock at B.

2. If the clock at A synchronizes with the clock at B and also with the clock at C, the clocks at B and C also synchronize with each other.

Thus with the help of certain imaginary physical experiments we have settled what is to be understood by synchronous stationary clocks located at different places, and have evidently obtained a definition of "simultaneous," or "synchronous," and of "time." The "time" of an event is that which is given simultaneously with the event by a stationary clock located at the place of the event, this clock being synchronous, and indeed synchronous for all time determinations, with a specified stationary clock.

In agreement with experience we further assume the quantity

$$\frac{2AB}{t'_A - t_A} = c,$$

to be a universal constant—the velocity of light in empty space.

It is essential to have time defined by means of stationary clocks in the stationary system, and the time now defined being appropriate to the stationary system we call it "the time of the stationary system."

## § 2. On the Relativity of Lengths and Times

The following reflexions are based on the principle of relativity and on the principle of the constancy of the velocity of light. These two principles we define as follows:—

1. The laws by which the states of physical systems undergo change are not affected, whether these changes of state be referred to the one or the other of two systems of co-ordinates in uniform translatory motion.

2. Any ray of light moves in the “stationary” system of co-ordinates with the determined velocity  $c$ , whether the ray be emitted by a stationary or by a moving body. Hence

$$\text{velocity} = \frac{\text{light path}}{\text{time interval}}$$

where time interval is to be taken in the sense of the definition in § 1.

Let there be given a stationary rigid rod; and let its length be  $l$  as measured by a measuring-rod which is also stationary. We now imagine the axis of the rod lying along the axis of  $x$  of the stationary system of co-ordinates, and that a uniform motion of parallel translation with velocity  $v$  along the axis of  $x$  in the direction of increasing  $x$  is then imparted to the rod. We now inquire as to the length of the moving rod, and imagine its length to be ascertained by the following two operations:—

(a) The observer moves together with the given measuring-rod and the rod to be measured, and measures the length of the rod directly by superposing the measuring-rod, in just the same way as if all three were at rest.

(b) By means of stationary clocks set up in the stationary system and synchronizing in accordance with § 1, the observer ascertains at what points of the stationary system the two ends of the rod to be measured are located at a definite time. The distance between these two points, measured by the measuring-rod already employed, which in this case is at rest, is also a length which may be designated “the length of the rod.”

In accordance with the principle of relativity the length

to be discovered by the operation (a)—we will call it “the length of the rod in the moving system”—must be equal to the length  $l$  of the stationary rod.

The length to be discovered by the operation (b) we will call “the length of the (moving) rod in the stationary system.” This we shall determine on the basis of our two principles, and we shall find that it differs from  $l$ .

Current kinematics tacitly assumes that the lengths determined by these two operations are precisely equal, or in other words, that a moving rigid body at the epoch  $t$  may in geometrical respects be perfectly represented by *the same* body *at rest* in a definite position.

We imagine further that at the two ends A and B of the rod, clocks are placed which synchronize with the clocks of the stationary system, that is to say that their indications correspond at any instant to the “time of the stationary system” at the places where they happen to be. These clocks are therefore “synchronous in the stationary system.”

We imagine further that with each clock there is a moving observer, and that these observers apply to both clocks the criterion established in § 1 for the synchronization of two clocks. Let a ray of light depart from A at the time \*  $t_A$ , let it be reflected at B at the time  $t_B$ , and reach A again at the time  $t'_A$ . Taking into consideration the principle of the constancy of the velocity of light we find that

$$t_B - t_A = \frac{r_{AB}}{c - v} \text{ and } t'_A - t_B = \frac{r_{AB}}{c + v}$$

where  $r_{AB}$  denotes the length of the moving rod—measured in the stationary system. Observers moving with the moving rod would thus find that the two clocks were not synchronous, while observers in the stationary system would declare the clocks to be synchronous.

So we see that we cannot attach any *absolute* signification to the concept of simultaneity, but that two events which, viewed from a system of co-ordinates, are simultaneous, can no longer be looked upon as simultaneous events when en-

\* “Time” here denotes “time of the stationary system” and also “position of hands of the moving clock situated at the place under discussion.”

visaged from a system which is in motion relatively to that system.

### § 3. Theory of the Transformation of Co-ordinates and Times from a Stationary System to another System in Uniform Motion of Translation Relatively to the Former

Let us in "stationary" space take two systems of co-ordinates, i.e. two systems, each of three rigid material lines, perpendicular to one another, and issuing from a point. Let the axes of  $X$  of the two systems coincide, and their axes of  $Y$  and  $Z$  respectively be parallel. Let each system be provided with a rigid measuring-rod and a number of clocks, and let the two measuring-rods, and likewise all the clocks of the two systems, be in all respects alike.

Now to the origin of one of the two systems ( $k$ ) let a constant velocity  $v$  be imparted in the direction of the increasing  $x$  of the other stationary system ( $K$ ), and let this velocity be communicated to the axes of the co-ordinates, the relevant measuring-rod, and the clocks. To any time of the stationary system  $K$  there then will correspond a definite position of the axes of the moving system, and from reasons of symmetry we are entitled to assume that the motion of  $k$  may be such that the axes of the moving system are at the time  $t$  (this " $t$ " always denotes a time of the stationary system) parallel to the axes of the stationary system.

We now imagine space to be measured from the stationary system  $K$  by means of the stationary measuring-rod, and also from the moving system  $k$  by means of the measuring-rod moving with it; and that we thus obtain the co-ordinates  $x, y, z$ , and  $\xi, \eta, \zeta$  respectively. Further, let the time  $t$  of the stationary system be determined for all points thereof at which there are clocks by means of light signals in the manner indicated in § 1; similarly let the time  $\tau$  of the moving system be determined for all points of the moving system at which there are clocks at rest relatively to that system by applying the method, given in § 1, of light signals between the points at which the latter clocks are located.

To any system of values  $x, y, z, t$ , which completely defines the place and time of an event in the stationary system, there

belongs a system of values  $\xi, \eta, \zeta, \tau$ , determining that event relatively to the system  $k$ , and our task is now to find the system of equations connecting these quantities.

In the first place it is clear that the equations must be *linear* on account of the properties of homogeneity which we attribute to space and time.

If we place  $x' = x - vt$ , it is clear that a point at rest in the system  $k$  must have a system of values  $x', y, z$ , independent of time. We first define  $\tau$  as a function of  $x', y, z$ , and  $t$ . To do this we have to express in equations that  $\tau$  is nothing else than the summary of the data of clocks at rest in system  $k$ , which have been synchronized according to the rule given in § 1.

From the origin of system  $k$  let a ray be emitted at the time  $\tau_0$  along the X-axis to  $x'$ , and at the time  $\tau_1$  be reflected thence to the origin of the co-ordinates, arriving there at the time  $\tau_2$ ; we then must have  $\frac{1}{2}(\tau_0 + \tau_2) = \tau_1$ , or, by inserting the arguments of the function  $\tau$  and applying the principle of the constancy of the velocity of light in the stationary system :—

$$\frac{1}{2} \left[ \tau(0, 0, 0, t) + \tau\left(0, 0, 0, t + \frac{x'}{c-v} + \frac{x'}{c+v}\right) \right] = \tau\left(x', 0, 0, t + \frac{x'}{c-v}\right).$$

Hence, if  $x'$  be chosen infinitesimally small,

$$\frac{1}{2} \left( \frac{1}{c-v} + \frac{1}{c+v} \right) \frac{\partial \tau}{\partial t} = \frac{\partial \tau}{\partial x'} + \frac{1}{c-v} \frac{\partial \tau}{\partial t},$$

or

$$\frac{\partial \tau}{\partial x'} + \frac{v}{c^2 - v^2} \frac{\partial \tau}{\partial t} = 0.$$

It is to be noted that instead of the origin of the co-ordinates we might have chosen any other point for the point of origin of the ray, and the equation just obtained is therefore valid for all values of  $x', y, z$ .

An analogous consideration—applied to the axes of Y and Z—it being borne in mind that light is always propagated along these axes, when viewed from the stationary system, with the velocity  $\sqrt{(c^2 - v^2)}$ , gives us

$$\frac{\partial \tau}{\partial y} = 0, \quad \frac{\partial \tau}{\partial z} = 0.$$

Since  $\tau$  is a *linear* function, it follows from these equations that

$$\tau = a \left( t - \frac{v}{c^2 - v^2} x' \right)$$

where  $a$  is a function  $\phi(v)$  at present unknown, and where for brevity it is assumed that at the origin of  $k$ ,  $\tau = 0$ , when  $t = 0$ .

With the help of this result we easily determine the quantities  $\xi$ ,  $\eta$ ,  $\zeta$  by expressing in equations that light (as required by the principle of the constancy of the velocity of light, in combination with the principle of relativity) is also propagated with velocity  $c$  when measured in the moving system. For a ray of light emitted at the time  $\tau = 0$  in the direction of the increasing  $\xi$

$$\xi = c\tau \text{ or } \xi = ac \left( t - \frac{v}{c^2 - v^2} x' \right).$$

But the ray moves relatively to the initial point of  $k$ , when measured in the stationary system, with the velocity  $c - v$ , so that

$$\frac{x'}{c - v} = t.$$

If we insert this value of  $t$  in the equation for  $\xi$ , we obtain

$$\xi = a \frac{c^2}{c^2 - v^2} x'.$$

In an analogous manner we find, by considering rays moving along the two other axes, that

$$\eta = c\tau = ac \left( t - \frac{v}{c^2 - v^2} x' \right)$$

when

$$\frac{y}{\sqrt{(c^2 - v^2)}} = t, \quad x' = 0.$$

Thus

$$\eta = a \frac{c}{\sqrt{(c^2 - v^2)}} y \text{ and } \zeta = a \frac{c}{\sqrt{(c^2 - v^2)}} z.$$

Substituting for  $x'$  its value, we obtain

$$\tau = \phi(v)\beta(t - vx/c^2),$$

$$\xi = \phi(v)\beta(x - vt),$$

$$\eta = \phi(v)y,$$

$$\zeta = \phi(v)z,$$

where

$$\beta = \frac{1}{\sqrt{1 - v^2/c^2}},$$

and  $\phi$  is an as yet unknown function of  $v$ . If no assumption whatever be made as to the initial position of the moving system and as to the zero point of  $\tau$ , an additive constant is to be placed on the right side of each of these equations.

We now have to prove that any ray of light, measured in the moving system, is propagated with the velocity  $c$ , if, as we have assumed, this is the case in the stationary system; for we have not as yet furnished the proof that the principle of the constancy of the velocity of light is compatible with the principle of relativity.

At the time  $t = \tau = 0$ , when the origin of the co-ordinates is common to the two systems, let a spherical wave be emitted therefrom, and be propagated with the velocity  $c$  in system  $K$ . If  $(x, y, z)$  be a point just attained by this wave, then

$$x^2 + y^2 + z^2 = c^2 t^2.$$

Transforming this equation with the aid of our equations of transformation we obtain after a simple calculation

$$\xi^2 + \eta^2 + \zeta^2 = c^2 \tau^2.$$

The wave under consideration is therefore no less a spherical wave with velocity of propagation  $c$  when viewed in the moving system. This shows that our two fundamental principles are compatible.\*

In the equations of transformation which have been developed there enters an unknown function  $\phi$  of  $v$ , which we will now determine.

For this purpose we introduce a third system of co-ordin-

\* The equations of the Lorentz transformation may be more simply deduced directly from the condition that in virtue of those equations the relation  $x^2 + y^2 + z^2 = c^2 t^2$  shall have as its consequence the second relation  $\xi^2 + \eta^2 + \zeta^2 = c^2 \tau^2$ .

ates  $K'$ , which relatively to the system  $k$  is in a state of parallel translatory motion parallel to the axis of  $X$ , such that the origin of co-ordinates of system  $k$  moves with velocity  $-v$  on the axis of  $X$ . At the time  $t = 0$  let all three origins coincide, and when  $t = x = y = z = 0$  let the time  $t'$  of the system  $K'$  be zero. We call the co-ordinates, measured in the system  $K$ ,  $x, y, z$ , and by a twofold application of our equations of transformation we obtain

$$\begin{aligned} t' &= \phi(-v)\beta(-v)(\tau + v\xi/c^2) = \phi(v)\phi(-v)t, \\ x' &= \phi(-v)\beta(-v)(\xi + v\tau) = \phi(v)\phi(-v)x, \\ y' &= \phi(-v)\eta = \phi(v)\phi(-v)y, \\ z' &= \phi(-v)\zeta = \phi(v)\phi(-v)z. \end{aligned}$$

Since the relations between  $x', y', z'$  and  $x, y, z$  do not contain the time  $t$ , the systems  $K$  and  $K'$  are at rest with respect to one another, and it is clear that the transformation from  $K$  to  $K'$  must be the identical transformation. Thus

$$\phi(v)\phi(-v) = 1.$$

We now inquire into the signification of  $\phi(v)$ . We give our attention to that part of the axis of  $Y$  of system  $k$  which lies between  $\xi = 0, \eta = 0, \zeta = 0$  and  $\xi = 0, \eta = l, \zeta = 0$ . This part of the axis of  $Y$  is a rod moving perpendicularly to its axis with velocity  $v$  relatively to system  $K$ . Its ends possess in  $K$  the co-ordinates

$$x_1 = vt, y_1 = \frac{l}{\phi(v)}, z_1 = 0$$

and

$$x_2 = vt, y_2 = 0, z_2 = 0.$$

The length of the rod measured in  $K$  is therefore  $l/\phi(v)$ ; and this gives us the meaning of the function  $\phi(v)$ . From reasons of symmetry it is now evident that the length of a given rod moving perpendicularly to its axis, measured in the stationary system, must depend only on the velocity and not on the direction and the sense of the motion. The length of the moving rod measured in the stationary system does not change, therefore, if  $v$  and  $-v$  are interchanged. Hence follows that  $l/\phi(v) = l/\phi(-v)$ , or

$$\phi(v) = \phi(-v).$$

It follows from this relation and the one previously found



that  $\phi(v) = 1$ , so that the transformation equations which have been found become

$$\begin{aligned}\tau &= \beta(t - vx/c^2), \\ \xi &= \beta(x - vt), \\ \eta &= y, \\ \zeta &= z,\end{aligned}$$

where

$$\beta = 1/\sqrt{1 - v^2/c^2}.$$

#### § 4. Physical Meaning of the Equations Obtained in Respect to Moving Rigid Bodies and Moving Clocks

We envisage a rigid sphere\* of radius  $R$ , at rest relatively to the moving system  $k$ , and with its centre at the origin of co-ordinates of  $k$ . The equation of the surface of this sphere moving relatively to the system  $K$  with velocity  $v$  is

$$\xi^2 + \eta^2 + \zeta^2 = R^2.$$

The equation of this surface expressed in  $x, y, z$  at the time  $t = 0$  is

$$\frac{x^2}{(\sqrt{1 - v^2/c^2})^2} + y^2 + z^2 = R^2.$$

A rigid body which, measured in a state of rest, has the form of a sphere, therefore has in a state of motion—viewed from the stationary system—the form of an ellipsoid of revolution with the axes

$$R\sqrt{1 - v^2/c^2}, R, R.$$

Thus, whereas the  $Y$  and  $Z$  dimensions of the sphere (and therefore of every rigid body of no matter what form) do not appear modified by the motion, the  $X$  dimension appears shortened in the ratio  $1:\sqrt{1 - v^2/c^2}$ , i.e. the greater the value of  $v$ , the greater the shortening. For  $v = c$  all moving objects—viewed from the “stationary” system—shrivel up into plain figures. For velocities greater than that of light our deliberations become meaningless; we shall, however, find in what follows, that the velocity of light in our theory plays the part, physically, of an infinitely great velocity.

\* That is, a body possessing spherical form when examined at rest.

It is clear that the same results hold good of bodies at rest in the "stationary" system, viewed from a system in uniform motion.

Further, we imagine one of the clocks which are qualified to mark the time  $t$  when at rest relatively to the stationary system, and the time  $\tau$  when at rest relatively to the moving system, to be located at the origin of the co-ordinates of  $k$ , and so adjusted that it marks the time  $\tau$ . What is the rate of this clock, when viewed from the stationary system?

Between the quantities  $x$ ,  $t$ , and  $\tau$ , which refer to the position of the clock, we have, evidently,  $x = vt$  and

$$\tau = \frac{1}{\sqrt{(1 - v^2/c^2)}}(t - vx/c^2).$$

Therefore,

$$\tau = t\sqrt{(1 - v^2/c^2)} = t - (1 - \sqrt{(1 - v^2/c^2)})t$$

whence it follows that the time marked by the clock (viewed in the stationary system) is slow by  $1 - \sqrt{(1 - v^2/c^2)}$  seconds per second, or—neglecting magnitudes of fourth and higher order—by  $\frac{1}{2}v^2/c^2$ .

From this there ensues the following peculiar consequence. If at the points A and B of K there are stationary clocks which, viewed in the stationary system, are synchronous; and if the clock at A is moved with the velocity  $v$  along the line AB to B, then on its arrival at B the two clocks no longer synchronize, but the clock moved from A to B lags behind the other which has remained at B by  $\frac{1}{2}tv^2/c^2$  (up to magnitudes of fourth and higher order),  $t$  being the time occupied in the journey from A to B.

It is at once apparent that this result still holds good if the clock moves from A to B in any polygonal line, and also when the points A and B coincide.

If we assume that the result proved for a polygonal line is also valid for a continuously curved line, we arrive at this result: If one of two synchronous clocks at A is moved in a closed curve with constant velocity until it returns to A, the journey lasting  $t$  seconds, then by the clock which has remained at rest the travelled clock on its arrival at A will be  $\frac{1}{2}tv^2/c^2$  second slow. Thence we conclude that a

balance-clock \* at the equator must go more slowly, by a very small amount, than a precisely similar clock situated at one of the poles under otherwise identical conditions.

### § 5. The Composition of Velocities

In the system  $k$  moving along the axis of  $X$  of the system  $K$  with velocity  $v$ , let a point move in accordance with the equations

$$\xi = w_{\xi}\tau, \eta = w_{\eta}\tau, \zeta = 0,$$

where  $w_{\xi}$  and  $w_{\eta}$  denote constants.

Required: the motion of the point relatively to the system  $K$ . If with the help of the equations of transformation developed in § 3 we introduce the quantities  $x, y, z, t$  into the equations of motion of the point, we obtain

$$\begin{aligned} x &= \frac{w_{\xi} + v}{1 + vw_{\xi}/c^2}t, \\ y &= \frac{\sqrt{(1 - v^2/c^2)}}{1 + vw_{\xi}/c^2}w_{\eta}t, \\ z &= 0. \end{aligned}$$

Thus the law of the parallelogram of velocities is valid according to our theory only to a first approximation. We set

$$\begin{aligned} V^2 &= \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2, \\ w^2 &= w_{\xi}^2 + w_{\eta}^2, \\ \alpha &= \tan^{-1} w_y/w_x, \end{aligned}$$

$\alpha$  is then to be looked upon as the angle between the velocities  $v$  and  $w$ . After a simple calculation we obtain

$$V = \frac{\sqrt{[(v^2 + w^2 + 2vw \cos \alpha) - (vw \sin \alpha/c^2)^2]}}{1 + vw \cos \alpha/c^2}.$$

It is worthy of remark that  $v$  and  $w$  enter into the expression for the resultant velocity in a symmetrical manner. If  $w$  also has the direction of the axis of  $X$ , we get

$$V = \frac{v + w}{1 + vw/c^2}.$$

\* Not a pendulum-clock, which is physically a system to which the Earth belongs. This case had to be excluded.

It follows from this equation that from a composition of two velocities which are less than  $c$ , there always results a velocity less than  $c$ . For if we set  $v = c - \kappa$ ,  $w = c - \lambda$ ,  $\kappa$  and  $\lambda$  being positive and less than  $c$ , then

$$V = c \frac{2c - \kappa - \lambda}{2c - \kappa - \lambda + \kappa\lambda/c} < c.$$

It follows, further, that the velocity of light  $c$  cannot be altered by composition with a velocity less than that of light. For this case we obtain

$$V = \frac{c + w}{1 + w/c} = c.$$

We might also have obtained the formula for  $V$ , for the case when  $v$  and  $w$  have the same direction, by compounding two transformations in accordance with § 3. If in addition to the systems  $K$  and  $k$  figuring in § 3 we introduce still another system of co-ordinates  $k'$  moving parallel to  $k$ , its initial point moving on the axis of  $X$  with the velocity  $w$ , we obtain equations between the quantities  $x, y, z, t$  and the corresponding quantities of  $k'$ , which differ from the equations found in § 3 only in that the place of " $v$ " is taken by the quantity

$$\frac{v + w}{1 + vw/c^2};$$

from which we see that such parallel transformations—necessarily—form a group.

We have now deduced the requisite laws of the theory of kinematics corresponding to our two principles, and we proceed to show their application to electrodynamics.

## II. ELECTRODYNAMICAL PART

### § 6. Transformation of the Maxwell-Hertz Equations for Empty Space. On the Nature of the Electromotive Forces Occurring in a Magnetic Field During Motion

Let the Maxwell-Hertz equations for empty space hold good for the stationary system  $K$ , so that we have

$$\begin{aligned}\frac{1}{c} \frac{\partial X}{\partial t} &= \frac{\partial N}{\partial y} - \frac{\partial M}{\partial z}, & \frac{1}{c} \frac{\partial L}{\partial t} &= \frac{\partial Y}{\partial z} - \frac{\partial Z}{\partial y}, \\ \frac{1}{c} \frac{\partial Y}{\partial t} &= \frac{\partial L}{\partial z} - \frac{\partial N}{\partial x}, & \frac{1}{c} \frac{\partial M}{\partial t} &= \frac{\partial Z}{\partial x} - \frac{\partial X}{\partial z}, \\ \frac{1}{c} \frac{\partial Z}{\partial t} &= \frac{\partial M}{\partial x} - \frac{\partial L}{\partial y}, & \frac{1}{c} \frac{\partial N}{\partial t} &= \frac{\partial X}{\partial y} - \frac{\partial Y}{\partial x},\end{aligned}$$

where (X, Y, Z) denotes the vector of the electric force, and (L, M, N) that of the magnetic force.

If we apply to these equations the transformation developed in § 3, by referring the electromagnetic processes to the system of co-ordinates there introduced, moving with the velocity  $v$ , we obtain the equations

$$\begin{aligned}\frac{1}{c} \frac{\partial X}{\partial \tau} &= \frac{\partial}{\partial \eta} \left\{ \beta \left( N - \frac{v}{c} Y \right) \right\} - \frac{\partial}{\partial \xi} \left\{ \beta \left( M + \frac{v}{c} Z \right) \right\}, \\ \frac{1}{c} \frac{\partial}{\partial \tau} \left\{ \beta \left( Y - \frac{v}{c} N \right) \right\} &= \frac{\partial L}{\partial \xi} - \frac{\partial}{\partial \xi} \left\{ \beta \left( N - \frac{v}{c} Y \right) \right\}, \\ \frac{1}{c} \frac{\partial}{\partial \tau} \left\{ \beta \left( Z + \frac{v}{c} M \right) \right\} &= \frac{\partial}{\partial \xi} \left\{ \beta \left( M + \frac{v}{c} Z \right) \right\} - \frac{\partial L}{\partial \eta}, \\ \frac{1}{c} \frac{\partial L}{\partial \tau} &= \frac{\partial}{\partial \xi} \left\{ \beta \left( Y - \frac{v}{c} N \right) \right\} - \frac{\partial}{\partial \eta} \left\{ \beta \left( Z + \frac{v}{c} M \right) \right\}, \\ \frac{1}{c} \frac{\partial}{\partial \tau} \left\{ \beta \left( M + \frac{v}{c} Z \right) \right\} &= \frac{\partial}{\partial \xi} \left\{ \beta \left( Z + \frac{v}{c} M \right) \right\} - \frac{\partial X}{\partial \xi}, \\ \frac{1}{c} \frac{\partial}{\partial \tau} \left\{ \beta \left( N - \frac{v}{c} Y \right) \right\} &= \frac{\partial X}{\partial \eta} - \frac{\partial}{\partial \xi} \left\{ \beta \left( Y - \frac{v}{c} N \right) \right\},\end{aligned}$$

where

$$\beta = 1/\sqrt{(1 - v^2/c^2)}.$$

Now the principle of relativity requires that if the Maxwell-Hertz equations for empty space hold good in system K, they also hold good in system  $k$ ; that is to say that the vectors of the electric and the magnetic force—(X', Y', Z') and (L', M', N')—of the moving system  $k$ , which are defined by their ponderomotive effects on electric or magnetic masses respectively, satisfy the following equations:—

$$\begin{aligned}\frac{1}{c} \frac{\partial X'}{\partial \tau} &= \frac{\partial N'}{\partial \eta} - \frac{\partial M'}{\partial \xi}, & \frac{1}{c} \frac{\partial L'}{\partial \tau} &= \frac{\partial Y'}{\partial \xi} - \frac{\partial Z'}{\partial \eta}, \\ \frac{1}{c} \frac{\partial Y'}{\partial \tau} &= \frac{\partial L'}{\partial \xi} - \frac{\partial N'}{\partial \xi}, & \frac{1}{c} \frac{\partial M'}{\partial \tau} &= \frac{\partial Z'}{\partial \xi} - \frac{\partial X'}{\partial \xi}, \\ \frac{1}{c} \frac{\partial Z'}{\partial \tau} &= \frac{\partial M'}{\partial \xi} - \frac{\partial L'}{\partial \eta}, & \frac{1}{c} \frac{\partial N'}{\partial \tau} &= \frac{\partial X'}{\partial \eta} - \frac{\partial Y'}{\partial \xi}.\end{aligned}$$

Evidently the two systems of equations found for system  $k$  must express exactly the same thing, since both systems of equations are equivalent to the Maxwell-Hertz equations for system  $K$ . Since, further, the equations of the two systems agree, with the exception of the symbols for the vectors, it follows that the functions occurring in the systems of equations at corresponding places must agree, with the exception of a factor  $\psi(v)$ , which is common for all functions of the one system of equations, and is independent of  $\xi, \eta, \zeta$  and  $\tau$  but depends upon  $v$ . Thus we have the relations

$$\begin{aligned}X' &= \psi(v)X, & L' &= \psi(v)L, \\ Y' &= \psi(v)\beta\left(Y - \frac{v}{c}N\right), & M' &= \psi(v)\beta\left(M + \frac{v}{c}Z\right), \\ Z' &= \psi(v)\beta\left(Z + \frac{v}{c}M\right), & N' &= \psi(v)\beta\left(N - \frac{v}{c}Y\right).\end{aligned}$$

If we now form the reciprocal of this system of equations, firstly by solving the equations just obtained, and secondly by applying the equations to the inverse transformation (from  $k$  to  $K$ ), which is characterized by the velocity  $-v$ , it follows, when we consider that the two systems of equations thus obtained must be identical, that  $\psi(v)\psi(-v) = 1$ . Further, from reasons of symmetry \*  $\psi(v) = \psi(-v)$ , and therefore

$$\psi(v) = 1,$$

and our equations assume the form

\* If, for example,  $X = Y = Z = L = M = 0$ , and  $N \neq 0$ , then from reasons of symmetry it is clear that when  $v$  changes sign without changing its numerical value,  $Y'$  must also change sign without changing its numerical value.

$$X' = X,$$

$$L' = L,$$

$$Y' = \beta \left( Y - \frac{v}{c} N \right), \quad M' = \beta \left( M + \frac{v}{c} Z \right),$$

$$Z' = \beta \left( Z + \frac{v}{c} M \right), \quad N' = \beta \left( N - \frac{v}{c} Y \right).$$

As to the interpretation of these equations we make the following remarks: Let a point charge of electricity have the magnitude "one" when measured in the stationary system K, i.e. let it when at rest in the stationary system exert a force of one dyne upon an equal quantity of electricity at a distance of one cm. By the principle of relativity this electric charge is also of the magnitude "one" when measured in the moving system. If this quantity of electricity is at rest relatively to the stationary system, then by definition the vector  $(X, Y, Z)$  is equal to the force acting upon it. If the quantity of electricity is at rest relatively to the moving system (at least at the relevant instant), then the force acting upon it, measured in the moving system, is equal to the vector  $(X', Y', Z')$ . Consequently the first three equations above allow themselves to be clothed in words in the two following ways:—

1. If a unit electric point charge is in motion in an electromagnetic field, there acts upon it, in addition to the electric force, an "electromotive force" which, if we neglect the terms multiplied by the second and higher powers of  $v/c$ , is equal to the vector-product of the velocity of the charge and the magnetic force, divided by the velocity of light. (Old manner of expression.)

2. If a unit electric point charge is in motion in an electromagnetic field, the force acting upon it is equal to the electric force which is present at the locality of the charge, and which we ascertain by transformation of the field to a system of co-ordinates at rest relatively to the electrical charge. (New manner of expression.)

The analogy holds with "magnetomotive forces." We see that electromotive force plays in the developed theory merely the part of an auxiliary concept, which owes its introduction to the circumstance that electric and magnetic forces

do not exist independently of the state of motion of the system of co-ordinates.

Furthermore it is clear that the asymmetry mentioned in the introduction as arising when we consider the currents produced by the relative motion of a magnet and a conductor, now disappears. Moreover, questions as to the "seat" of electrodynamic electromotive forces (unipolar machines) now have no point.

### § 7. Theory of Doppler's Principle and of Aberration

In the system  $K$ , very far from the origin of co-ordinates, let there be a source of electrodynamic waves, which in a part of space containing the origin of co-ordinates may be represented to a sufficient degree of approximation by the equations

$$X = X_0 \sin \Phi, \quad L = L_0 \sin \Phi,$$

$$Y = Y_0 \sin \Phi, \quad M = M_0 \sin \Phi,$$

$$Z = Z_0 \sin \Phi, \quad N = N_0 \sin \Phi,$$

where

$$\Phi = \omega \left\{ t - \frac{1}{c}(lx + my + nz) \right\}.$$

Here  $(X_0, Y_0, Z_0)$  and  $(L_0, M_0, N_0)$  are the vectors defining the amplitude of the wave-train, and  $l, m, n$  the direction-cosines of the wave-normals. We wish to know the constitution of these waves, when they are examined by an observer at rest in the moving system  $k$ .

Applying the equations of transformation found in § 6 for electric and magnetic forces, and those found in § 3 for the co-ordinates and the time, we obtain directly

$$X' = X_0 \sin \Phi', \quad L' = L_0 \sin \Phi',$$

$$Y' = \beta(Y_0 - vN_0/c) \sin \Phi', \quad M' = \beta(M_0 + vZ_0/c) \sin \Phi',$$

$$Z' = \beta(Z_0 + vM_0/c) \sin \Phi', \quad N' = \beta(N_0 - vY_0/c) \sin \Phi',$$

$$\Phi' = \omega' \left\{ \tau - \frac{1}{c}(l'\xi + m'\eta + n'\zeta) \right\}$$



where

$$\omega' = \omega\beta(1 - lv/c),$$

$$l' = \frac{l - v/c}{1 - lv/c},$$

$$m' = \frac{m}{\beta(1 - lv/c)},$$

$$n' = \frac{n}{\beta(1 - lv/c)}.$$

From the equation for  $\omega'$  it follows that if an observer is moving with velocity  $v$  relatively to an infinitely distant source of light of frequency  $\nu$ , in such a way that the connecting line "source—observer" makes the angle  $\phi$  with the velocity of the observer referred to a system of co-ordinates which is at rest relatively to the source of light, the frequency  $\nu'$  of the light perceived by the observer is given by the equation

$$\nu' = \nu \frac{1 - \cos \phi \cdot v/c}{\sqrt{(1 - v^2/c^2)}}.$$

This is Doppler's principle for any velocities whatever. When  $\phi = 0$  the equation assumes the perspicuous form

$$\nu' = \nu \sqrt{\frac{1 - v/c}{1 + v/c}}.$$

We see that, in contrast with the customary view, when  $v = -c$ ,  $\nu' = \infty$ .

If we call the angle between the wave-normal (direction of the ray) in the moving system and the connecting line "source—observer"  $\phi'$ , the equation for  $l'$  assumes the form

$$\cos \phi' = \frac{\cos \phi - v/c}{1 - \cos \phi \cdot v/c}.$$

This equation expresses the law of aberration in its most general form. If  $\phi = \frac{1}{2}\pi$ , the equation becomes simply

$$\cos \phi' = -v/c.$$

We still have to find the amplitude of the waves, as it appears in the moving system. If we call the amplitude of the electric or magnetic force  $A$  or  $A'$  respectively, accordingly

as it is measured in the stationary system or in the moving system, we obtain

$$A'^2 = A^2 \frac{(1 - \cos \phi \cdot v/c)^2}{1 - v^2/c^2}$$

which equation, if  $\phi = 0$ , simplifies into

$$A'^2 = A^2 \frac{1 - v/c}{1 + v/c}.$$

It follows from these results that to an observer approaching a source of light with the velocity  $c$ , this source of light must appear of infinite intensity.

### § 8. Transformation of the Energy of Light Rays. Theory of the Pressure of Radiation Exerted on Perfect Reflectors

Since  $A^2/8\pi$  equals the energy of light per unit of volume, we have to regard  $A'^2/8\pi$ , by the principle of relativity, as the energy of light in the moving system. Thus  $A'^2/A^2$  would be the ratio of the "measured in motion" to the "measured at rest" energy of a given light complex, if the volume of a light complex were the same, whether measured in  $K$  or in  $k$ . But this is not the case. If  $l, m, n$  are the direction-cosines of the wave-normals of the light in the stationary system, no energy passes through the surface elements of a spherical surface moving with the velocity of light:—

$$(x - lct)^2 + (y - mct)^2 + (z - nct)^2 = R^2.$$

We may therefore say that this surface permanently encloses the same light complex. We inquire as to the quantity of energy enclosed by this surface, viewed in system  $k$ , that is, as to the energy of the light complex relatively to the system  $k$ .

The spherical surface—viewed in the moving system—is an ellipsoidal surface, the equation for which, at the time  $\tau = 0$ , is

$$(\beta\xi - l\beta\xi v/c)^2 + (\eta - m\beta\xi v/c)^2 + (\zeta - n\beta\xi v/c)^2 = R^2.$$

If  $S$  is the volume of the sphere, and  $S'$  that of this ellipsoid,

then by a simple calculation

$$\frac{S'}{S} = \frac{\sqrt{1 - v^2/c^2}}{1 - \cos \phi \cdot v/c}.$$

Thus, if we call the light energy enclosed by this surface  $E$  when it is measured in the stationary system, and  $E'$  when measured in the moving system, we obtain

$$\frac{E'}{E} = \frac{A^2 S'}{A^2 S} = \frac{1 - \cos \phi \cdot v/c}{\sqrt{(1 - v^2/c^2)}},$$

and this formula, when  $\phi = 0$ , simplifies into

$$\frac{E'}{E} = \sqrt{\frac{1 - v/c}{1 + v/c}}.$$

It is remarkable that the energy and the frequency of a light complex vary with the state of motion of the observer in accordance with the same law.

Now let the co-ordinate plane  $\xi = 0$  be a perfectly reflecting surface, at which the plane waves considered in § 7 are reflected. We seek for the pressure of light exerted on the reflecting surface, and for the direction, frequency, and intensity of the light after reflexion.

Let the incidental light be defined by the quantities  $A$ ,  $\cos \phi$ ,  $\nu$  (referred to system  $K$ ). Viewed from  $k$  the corresponding quantities are

$$\begin{aligned} A' &= A \frac{1 - \cos \phi \cdot v/c}{\sqrt{(1 - v^2/c^2)}}, \\ \cos \phi' &= \frac{\cos \phi - v/c}{1 - \cos \phi \cdot v/c}, \\ \nu' &= \nu \frac{1 - \cos \phi \cdot v/c}{\sqrt{(1 - v^2/c^2)}}. \end{aligned}$$

For the reflected light, referring the process to system  $k$ , we obtain

$$\begin{aligned} A'' &= A' \\ \cos \phi'' &= -\cos \phi' \\ \nu'' &= \nu' \end{aligned}$$

Finally, by transforming back to the stationary system  $K$ , we obtain for the reflected light

$$A''' = A'' \frac{1 + \cos \phi'' \cdot v/c}{\sqrt{(1 - v^2/c^2)}} = A \frac{1 - 2 \cos \phi \cdot v/c + v^2/c^2}{1 - v^2/c^2},$$

$$\cos \phi''' = \frac{\cos \phi'' + v/c}{1 + \cos \phi'' \cdot v/c} = - \frac{(1 + v^2/c^2) \cos \phi - 2v/c}{1 - 2 \cos \phi \cdot v/c + v^2/c^2}$$

$$v''' = v'' \frac{1 + \cos \phi'' v/c}{\sqrt{(1 - v^2/c^2)}} = v \frac{1 - 2 \cos \phi \cdot v/c + v^2/c^2}{1 - v^2/c^2}.$$

The energy (measured in the stationary system) which is incident upon unit area of the mirror in unit time is evidently  $A^2(c \cos \phi - v)/8\pi$ . The energy leaving the unit of surface of the mirror in the unit of time is  $A'''^2(-c \cos \phi''' + v)/8\pi$ . The difference of these two expressions is, by the principle of energy, the work done by the pressure of light in the unit of time. If we set down this work as equal to the product  $Pv$ , where  $P$  is the pressure of light, we obtain

$$P = 2 \cdot \frac{A^2}{8\pi} \frac{(\cos \phi - v/c)^2}{1 - v^2/c^2}.$$

In agreement with experiment and with other theories, we obtain to a first approximation

$$P = 2 \cdot \frac{A^2}{8\pi} \cos^2 \phi.$$

All problems in the optics of moving bodies can be solved by the method here employed. What is essential is, that the electric and magnetic force of the light which is influenced by a moving body, be transformed into a system of co-ordinates at rest relatively to the body. By this means all problems in the optics of moving bodies will be reduced to a series of problems in the optics of stationary bodies.

### § 9. Transformation of the Maxwell-Hertz Equations when Convection-Currents are Taken into Account

We start from the equations

$$\begin{aligned} \frac{1}{c} \left\{ \frac{\partial X}{\partial t} + u_x \rho \right\} &= \frac{\partial N}{\partial y} - \frac{\partial M}{\partial z}, & \frac{1}{c} \frac{\partial L}{\partial t} &= \frac{\partial Y}{\partial z} - \frac{\partial Z}{\partial y}, \\ \frac{1}{c} \left\{ \frac{\partial Y}{\partial t} + u_y \rho \right\} &= \frac{\partial L}{\partial z} - \frac{\partial N}{\partial x}, & \frac{1}{c} \frac{\partial M}{\partial t} &= \frac{\partial Z}{\partial x} - \frac{\partial X}{\partial z}, \\ \frac{1}{c} \left\{ \frac{\partial Z}{\partial t} + u_z \rho \right\} &= \frac{\partial M}{\partial x} - \frac{\partial L}{\partial y}, & \frac{1}{c} \frac{\partial N}{\partial t} &= \frac{\partial X}{\partial y} - \frac{\partial Y}{\partial x}, \end{aligned}$$

where

$$\rho = \frac{\partial X}{\partial x} + \frac{\partial Y}{\partial y} + \frac{\partial Z}{\partial z}$$

denotes  $4\pi$  times the density of electricity, and  $(u_x, u_y, u_z)$  the velocity-vector of the charge. If we imagine the electric charges to be invariably coupled to small rigid bodies (ions, electrons), these equations are the electromagnetic basis of the Lorentzian electrodynamics and optics of moving bodies.

Let these equations be valid in the system  $K$ , and transform them, with the assistance of the equations of transformation given in §§ 3 and 6, to the system  $k$ . We then obtain the equations

$$\begin{aligned} \frac{1}{c} \left\{ \frac{\partial X'}{\partial \tau} + u_x \rho' \right\} &= \frac{\partial N'}{\partial \eta} - \frac{\partial M'}{\partial \xi}, \quad \frac{1}{c} \frac{\partial L'}{\partial \tau} = \frac{\partial Y'}{\partial \xi} - \frac{\partial Z'}{\partial \eta}, \\ \frac{1}{c} \left\{ \frac{\partial Y'}{\partial \tau} + u_y \rho' \right\} &= \frac{\partial L'}{\partial \xi} - \frac{\partial N'}{\partial \xi}, \quad \frac{1}{c} \frac{\partial M'}{\partial \tau} = \frac{\partial Z'}{\partial \xi} - \frac{\partial X'}{\partial \xi}, \\ \frac{1}{c} \left\{ \frac{\partial Z'}{\partial \tau} + u_z \rho' \right\} &= \frac{\partial M'}{\partial \xi} - \frac{\partial L'}{\partial \eta}, \quad \frac{1}{c} \frac{\partial N'}{\partial \tau} = \frac{\partial X'}{\partial \eta} - \frac{\partial Y'}{\partial \xi}, \end{aligned}$$

where

$$u_x = \frac{u_x - v}{1 - u_x v / c^2}$$

$$u_y = \frac{u_y}{\beta(1 - u_x v / c^2)}$$

$$u_z = \frac{u_z}{\beta(1 - u_x v / c^2)},$$

and

$$\begin{aligned} \rho' &= \frac{\partial X'}{\partial \xi} + \frac{\partial Y'}{\partial \eta} + \frac{\partial Z'}{\partial \xi} \\ &= \beta(1 - u_x v / c^2) \rho. \end{aligned}$$

Since—as follows from the theorem of addition of velocities (§ 5)—the vector  $(u_x, u_y, u_z)$  is nothing else than the velocity of the electric charge, measured in the system  $k$ , we have the proof that, on the basis of our kinematical principles, the electrodynamic foundation of Lorentz's theory of the electrodynamics of moving bodies is in agreement with the principle of relativity.

In addition I may briefly remark that the following import-

ant law may easily be deduced from the developed equations :  
 If an electrically charged body is in motion anywhere in space without altering its charge when regarded from a system of co-ordinates moving with the body, its charge also remains—when regarded from the “stationary” system K—constant.

### § 10. Dynamics of the Slowly Accelerated Electron

Let there be in motion in an electromagnetic field an electrically charged particle (in the sequel called an “electron”), for the law of motion of which we assume as follows :—

If the electron is at rest at a given epoch, the motion of the electron ensues in the next instant of time according to the equations

$$m \frac{d^2x}{dt^2} = eX$$

$$m \frac{d^2y}{dt^2} = eY$$

$$m \frac{d^2z}{dt^2} = eZ$$

where  $x, y, z$  denote the co-ordinates of the electron, and  $m$  the mass of the electron, as long as its motion is slow.

Now, secondly, let the velocity of the electron at a given epoch be  $v$ . We seek the law of motion of the electron in the immediately ensuing instants of time.

Without affecting the general character of our considerations, we may and will assume that the electron, at the moment when we give it our attention, is at the origin of the co-ordinates, and moves with the velocity  $v$  along the axis of  $X$  of the system  $K$ . It is then clear that at the given moment ( $t = 0$ ) the electron is at rest relatively to a system of co-ordinates which is in parallel motion with velocity  $v$  along the axis of  $X$ .

From the above assumption, in combination with the principle of relativity, it is clear that in the immediately ensuing time (for small values of  $t$ ) the electron, viewed from the system  $k$ , moves in accordance with the equations

$$m \frac{d^2 \xi}{d\tau^2} = \epsilon X',$$

$$m \frac{d^2 \eta}{d\tau^2} = \epsilon Y',$$

$$m \frac{d^2 \zeta}{d\tau^2} = \epsilon Z',$$

in which the symbols  $\xi, \eta, \zeta, \tau, X', Y', Z'$  refer to  $k$ . If, further, we decide that when  $t = x = y = \tau = \xi = \eta = \zeta = 0$ , the transformation equations 6 hold good, so that we have

$$\xi = \beta(x - vt), \eta = y, \zeta = z, \tau = \beta(t - vN/c), \\ X' = X, Y' = \beta(Y - vN/c), Z' = \beta(Z +$$

With the help of these equations we transform equations of motion from system  $k$  to system  $l$

$$\left. \begin{aligned} \frac{d^2 x}{dt^2} &= \frac{\epsilon}{m \beta^3} X \\ \frac{d^2 y}{dt^2} &= \frac{\epsilon}{m \beta} \left( Y - \frac{v}{c} N \right) \\ \frac{d^2 z}{dt^2} &= \frac{\epsilon}{m \beta} \left( Z + \frac{v}{c} M \right) \end{aligned} \right\} .$$

Taking the ordinary point of view we return to the "longitudinal" and the "transverse" moving electron. We write the equations (A)

$$m \beta^3 \frac{d^2 x}{dt^2} = \epsilon X = \epsilon X',$$

$$m \beta^3 \frac{d^2 y}{dt^2} = \epsilon \beta \left( Y - \frac{v}{c} N \right) = \epsilon Y$$

$$m \beta^3 \frac{d^2 z}{dt^2} = \epsilon \beta \left( Z + \frac{v}{c} M \right) = \epsilon Z$$

and remark firstly that  $\epsilon X', \epsilon Y', \epsilon Z'$  are the ponderomotive force acting upon the electron indeed as viewed in a system moving at the same velocity as the electron, with the same velocity as the electron might be measured, for example, by a spring

in the last-mentioned system.) Now if we call this force simply "the force acting upon the electron,"\* and maintain the equation—mass  $\times$  acceleration = force—and if we also decide that the accelerations are to be measured in the stationary system K, we derive from the above equations

$$\text{Longitudinal mass} = \frac{m}{(\sqrt{1 - v^2/c^2})^3}.$$

$$\text{Transverse mass} = \frac{m}{1 - v^2/c^2}.$$

With a different definition of force and acceleration we should naturally obtain other values for the masses. This shows us that in comparing different theories of the motion of the electron we must proceed very cautiously.

We remark that these results as to the mass are also valid for ponderable material points, because a ponderable material point can be made into an electron (in our sense of the word) by the addition of an electric charge, *no matter how small*.

We will now determine the kinetic energy of the electron. If an electron moves from rest at the origin of co-ordinates of the system K along the axis of X under the action of an electrostatic force X, it is clear that the energy withdrawn from the electrostatic field has the value  $\int \epsilon X dx$ . As the electron is to be slowly accelerated, and consequently may not give off any energy in the form of radiation, the energy withdrawn from the electrostatic field must be put down as equal to the energy of motion W of the electron. Bearing in mind that during the whole process of motion which we are considering, the first of the equations (A) applies, we therefore obtain

$$\begin{aligned} W &= \int \epsilon X dx = m \int_0^v \beta^3 v dv \\ &= mc^2 \left\{ \frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right\}. \end{aligned}$$

Thus, when  $v = c$ , W becomes infinite. Velocities

\* The definition of force here given is not advantageous, as was first shown by M. Planck. It is more to the point to define force in such a way that the laws of momentum and energy assume the simplest form.



greater than that of light have—as in our previous results—no possibility of existence.

This expression for the kinetic energy must also, by virtue of the argument stated above, apply to ponderable masses as well.

We will now enumerate the properties of the motion of the electron which result from the system of equations (A), and are accessible to experiment.

1. From the second equation of the system (A) it follows that an electric force  $Y$  and a magnetic force  $N$  have an equally strong deflective action on an electron moving with the velocity  $v$ , when  $Y = Nv/c$ . Thus we see that it is possible by our theory to determine the velocity of the electron from the ratio of the magnetic power of deflexion  $A_m$  to the electric power of deflexion  $A_e$ , for any velocity, by applying the law

$$\frac{A_m}{A_e} = \frac{v}{c}.$$

This relationship may be tested experimentally, since the velocity of the electron can be directly measured, e.g. by means of rapidly oscillating electric and magnetic fields.

2. From the deduction for the kinetic energy of the electron it follows that between the potential difference,  $P$ , traversed and the acquired velocity  $v$  of the electron there must be the relationship

$$P = \int X dx = \frac{m}{\epsilon} c^2 \left\{ \frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right\}$$

3. We calculate the radius of curvature of the path of the electron when a magnetic force  $N$  is present (as the only deflective force), acting perpendicularly to the velocity of the electron. From the second of the equations (A) we obtain

$$-\frac{d^2y}{dt^2} = \frac{v^2}{R} = \frac{\epsilon}{m} \frac{v}{c} N \sqrt{1 - \frac{v^2}{c^2}}$$

or

$$R = \frac{mc^2}{\epsilon} \cdot \frac{v/c}{\sqrt{(1 - v^2/c^2)}} \cdot \frac{1}{N}.$$

These three relationships are a complete expression for

## ELECTRODYNAMICS

65

the laws according to which, by the theory here advanced, the electron must move.

In conclusion I wish to say that in working at the problem here dealt with I have had the loyal assistance of my friend and colleague M. Besso, and that I am indebted to him for several valuable suggestions.



# DOES THE INERTIA OF A BODY DEPEND UPON ITS ENERGY-CONTENT ?

BY

A. EINSTEIN

*Translated from "Ist die Trägheit eines Körpers von seinem  
Energiegehalt abhängig?" Annalen der Physik, 17,  
1905.*



# DOES THE INERTIA OF A BODY DEPEND UPON ITS ENERGY-CONTENT?

By A. EINSTEIN

THE results of the previous investigation lead to a very interesting conclusion, which is here to be deduced.

I based that investigation on the Maxwell-Hertz equations for empty space, together with the Maxwellian expression for the electromagnetic energy of space, and in addition the principle that:—

*The laws by which the states of physical systems alter are independent of the alternative, to which of two systems of co-ordinates, in uniform motion of parallel translation relatively to each other, these alterations of state are referred (principle of relativity).*

With these principles \* as my basis I deduced *inter alia* the following result (§ 8):—

Let a system of plane waves of light, referred to the system of co-ordinates  $(x, y, z)$ , possess the energy  $l$ ; let the direction of the ray (the wave-normal) make an angle  $\phi$  with the axis of  $x$  of the system. If we introduce a new system of co-ordinates  $(\xi, \eta, \zeta)$  moving in uniform parallel translation with respect to the system  $(x, y, z)$ , and having its origin of co-ordinates in motion along the axis of  $x$  with the velocity  $v$ , then this quantity of light—measured in the system  $(\xi, \eta, \zeta)$ —possesses the energy

$$l^* = l \frac{1 - \frac{v}{c} \cos \phi}{\sqrt{1 - v^2/c^2}}$$

\*The principle of the constancy of the velocity of light is of course contained in Maxwell's equations.

where  $c$  denotes the velocity of light. We shall make use of this result in what follows.

Let there be a stationary body in the system  $(x, y, z)$ , and let its energy—referred to the system  $(x, y, z)$ —be  $E_0$ . Let the energy of the body relative to the system  $(\xi, \eta, \zeta)$ , moving as above with the velocity  $v$ , be  $H_0$ .

Let this body send out, in a direction making an angle  $\phi$  with the axis of  $x$ , plane waves of light, of energy  $\frac{1}{2}L$  measured relatively to  $(x, y, z)$ , and simultaneously an equal quantity of light in the opposite direction. Meanwhile the body remains at rest with respect to the system  $(x, y, z)$ . The principle of energy must apply to this process, and in fact (by the principle of relativity) with respect to both systems of co-ordinates. If we call the energy of the body after the emission of light  $E_1$  or  $H_1$  respectively, measured relatively to the system  $(x, y, z)$  or  $(\xi, \eta, \zeta)$  respectively, then by employing the relation given above we obtain

$$E_0 = E_1 + \frac{1}{2}L + \frac{1}{2}L,$$

$$\begin{aligned} H_0 &= H_1 + \frac{1}{2}L \frac{1 - \frac{v}{c} \cos \phi}{\sqrt{1 - v^2/c^2}} + \frac{1}{2}L \frac{1 + \frac{v}{c} \cos \phi}{\sqrt{1 - v^2/c^2}} \\ &= H_1 + \frac{L}{\sqrt{1 - v^2/c^2}}. \end{aligned}$$

By subtraction we obtain from these equations

$$H_0 - E_0 - (H_1 - E_1) = L \left\{ \frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right\}.$$

The two differences of the form  $H - E$  occurring in this expression have simple physical significations.  $H$  and  $E$  are energy values of the same body referred to two systems of co-ordinates which are in motion relatively to each other, the body being at rest in one of the two systems (system  $(x, y, z)$ ). Thus it is clear that the difference  $H - E$  can differ from the kinetic energy  $K$  of the body, with respect to the other system  $(\xi, \eta, \zeta)$ , only by an additive constant  $C$ , which depends on the choice of the arbitrary additive constants of the energies  $H$  and  $E$ . Thus we may place

$$\begin{aligned} H_0 - E_0 &= K_0 + C, \\ H_1 - E_1 &= K_1 + C, \end{aligned}$$

since  $C$  does not change during the emission of light. So we have

$$K_0 - K_1 = L \left\{ \frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right\}.$$

The kinetic energy of the body with respect to  $(\xi, \eta, \zeta)$  diminishes as a result of the emission of light, and the amount of diminution is independent of the properties of the body. Moreover, the difference  $K_0 - K_1$ , like the kinetic energy of the electron (§ 10), depends on the velocity.

Neglecting magnitudes of fourth and higher orders we may place

$$K_0 - K_1 = \frac{1}{2} \frac{L}{c^2} v^2.$$

From this equation it directly follows that:—

*If a body gives off the energy  $L$  in the form of radiation, its mass diminishes by  $L/c^2$ .* The fact that the energy withdrawn from the body becomes energy of radiation evidently makes no difference, so that we are led to the more general conclusion that

The mass of a body is a measure of its energy-content; if the energy changes by  $L$ , the mass changes in the same sense by  $L/9 \times 10^{20}$ , the energy being measured in ergs, and the mass in grammes.

It is not impossible that with bodies whose energy-content is variable to a high degree (e.g. with radium salts) the theory may be successfully put to the test.

If the theory corresponds to the facts, radiation conveys inertia between the emitting and absorbing bodies.





# SPACE AND TIME

BY

H. MINKOWSKI

*A Translation of an Address delivered at the 80th Assembly  
of German Natural Scientists and Physicians, at Cologne,  
21 September, 1908.*



## SPACE AND TIME

By H. MINKOWSKI

THE views of space and time which I wish to lay before you have sprung from the soil of experimental physics, and therein lies their strength. They are radical. Henceforth space by itself, and time by itself, are doomed to fade away into mere shadows, and only a kind of union of the two will preserve an independent reality.

### I

First of all I should like to show how it might be possible, setting out from the accepted mechanics of the present day, along a purely mathematical line of thought, to arrive at changed ideas of space and time. The equations of Newton's mechanics exhibit a two-fold invariance. Their form remains unaltered, firstly, if we subject the underlying system of spatial co-ordinates to any arbitrary *change of position*; secondly, if we change its state of motion, namely, by imparting to it any *uniform translatory motion*; furthermore, the zero point of time is given no part to play. We are accustomed to look upon the axioms of geometry as finished with, when we feel ripe for the axioms of mechanics, and for that reason the two invariances are probably rarely mentioned in the same breath. Each of them by itself signifies, for the differential equations of mechanics, a certain group of transformations. The existence of the first group is looked upon as a fundamental characteristic of space. The second group is preferably treated with disdain, so that we with untroubled minds may overcome the difficulty of never being able to decide, from physical phenomena, whether space, which is supposed to be stationary, may not be after all in a

state of uniform translation. Thus the two groups, side by side, lead their lives entirely apart. Their utterly heterogeneous character may have discouraged any attempt to compound them. But it is precisely when they are compounded that the complete group, as a whole, gives us to think.

We will try to visualize the state of things by the graphic method. Let  $x, y, z$  be rectangular co-ordinates for space, and let  $t$  denote time. The objects of our perception invariably include places and times in combination. Nobody has ever noticed a place except at a time, or a time except at a place. But I still respect the dogma that both space and time have independent significance. A point of space at a point of time, that is, a system of values  $x, y, z, t$ , I will call a *world-point*. The multiplicity of all thinkable  $x, y, z, t$  systems of values we will christen the *world*. With this most valiant piece of chalk I might project upon the blackboard four world-axes. Since merely one chalky axis, as it is, consists of molecules all a-thrill, and moreover is taking part in the earth's travels in the universe, it already affords us ample scope for abstraction; the somewhat greater abstraction associated with the number four is for the mathematician no infliction. Not to leave a yawning void anywhere, we will imagine that everywhere and everywhen there is something perceptible. To avoid saying "matter" or "electricity" I will use for this something the word "substance." We fix our attention on the substantial point which is at the world-point  $x, y, z, t$ , and imagine that we are able to recognize this substantial point at any other time. Let the variations  $dx, dy, dz$  of the space co-ordinates of this substantial point correspond to a time element  $dt$ . Then we obtain, as an image, so to speak, of the everlasting career of the substantial point, a curve in the world, a *world-line*, the points of which can be referred unequivocally to the parameter  $t$  from  $-\infty$  to  $+\infty$ . The whole universe is seen to resolve itself into similar world-lines, and I would fain anticipate myself by saying that in my opinion physical laws might find their most perfect expression as reciprocal relations between these world-lines.

The concepts, space and time, cause the  $x, y, z$ -manifold  $t = 0$  and its two sides  $t > 0$  and  $t < 0$  to fall asunder. If,

for simplicity, we retain the same zero point of space and time, the first-mentioned group signifies in mechanics that we may subject the axes of  $x, y, z$  at  $t = 0$  to any rotation we choose about the origin, corresponding to the homogeneous linear transformations of the expression

$$x^2 + y^2 + z^2.$$

But the second group means that we may—also without changing the expression of the laws of mechanics—replace  $x, y, z, t$  by  $x - at, y - \beta t, z - \gamma t, t$  with any constant values of  $a, \beta, \gamma$ . Hence we may give to the time axis whatever direction we choose towards the upper half of the world,  $t > 0$ . Now what has the requirement of orthogonality in space to do with this perfect freedom of the time axis in an upward direction?

To establish the connexion, let us take a positive parameter  $c$ , and consider the graphical representation of

$$c^2 t^2 - x^2 - y^2 - z^2 = 1.$$

It consists of two surfaces separated by  $t = 0$ , on the analogy of a hyperboloid of two sheets. We consider the sheet in the region  $t > 0$ , and now take those homogeneous linear transformations of  $x, y, z, t$  into four new variables  $x', y', z', t'$ , for which the expression for this sheet in the new variables is of the same form. It is evident that the rotations of space about the origin pertain to these transformations. Thus we gain full comprehension of the rest of the transformations simply by taking into consideration one among them, such that  $y$  and  $z$  remain unchanged. We draw (Fig. 1) the section of this sheet by the plane of the axes of  $x$  and  $t$ —the upper branch of the hyperbola  $c^2 t^2 - x^2 = 1$ , with its asymptotes. From the origin  $O$  we draw any radius vector  $OA'$  of this branch of the hyperbola; draw the tangent to the hyperbola at  $A'$  to cut the asymptote on the right at  $B'$ ; complete the parallelogram  $OA'B'C'$ ; and finally, for subsequent use, produce  $B'C'$  to cut the axis of  $x$  at  $D'$ . Now if we take  $OC'$  and  $OA'$  as axes of oblique co-ordinates  $x', t'$ , with the measures  $OC' = 1, OA' = 1/c$ , then that branch of the hyperbola again acquires the expression  $c^2 t'^2 - x'^2 = 1, t' > 0$ , and the transition from  $x, y, z, t$  to  $x', y', z', t'$  is one of

the transformations in question. With these transformations we now associate the arbitrary displacements of the zero point of space and time, and thereby constitute a group of transformations, which is also, evidently, dependent on the parameter  $c$ . This group I denote by  $G_c$ .

If we now allow  $c$  to increase to infinity, and  $1/c$  therefore to converge towards zero, we see from the figure that the




FIG. 1.

branch of the hyperbola bends more and more towards the axis of  $x$ , the angle of the asymptotes becomes more and more obtuse, and that in the limit this special transformation changes into one in which the axis of  $t'$  may have any upward direction whatever, while  $x'$  approaches more and more exactly to  $x$ . In view of this it is clear that group  $G_c$  in the limit when  $c = \infty$ , that is the group  $G_\infty$ , becomes no other than that complete group which is appropriate to Newtonian

mechanics. This being so, and since  $G_c$  is mathematically more intelligible than  $G_\infty$ , it looks as though the thought might have struck some mathematician, fancy-free, that after all, as a matter of fact, natural phenomena do not possess an invariance with the group  $G_\infty$ , but rather with a group  $G_c$ ,  $c$  being finite and determinate, but in ordinary units of measure, *extremely great*. Such a premonition would have been an extraordinary triumph for pure mathematics. Well, mathematics, though it now can display only staircase-wit, has the satisfaction of being wise after the event, and is able, thanks to its happy antecedents, with its senses sharpened by an unhampered outlook to far horizons, to grasp forthwith the far-reaching consequences of such a metamorphosis of our concept of nature.

I will state at once what is the value of  $c$  with which we shall finally be dealing. It is the velocity of the propagation of light in empty space. To avoid speaking either of space or of emptiness, we may define this magnitude in another way, as the ratio of the electromagnetic to the electrostatic unit of electricity.

The existence of the invariance of natural laws for the relevant group  $G_c$  would have to be taken, then, in this way:—

From the totality of natural phenomena it is possible, by successively enhanced approximations, to derive more and more exactly a system of reference  $x, y, z, t$ , space and time, by means of which these phenomena then present themselves in agreement with definite laws. But when this is done, this system of reference is by no means unequivocally determined by the phenomena. *It is still possible to make any change in the system of reference that is in conformity with the transformations of the group  $G_c$ , and leave the expression of the laws of nature unaltered.*

For example, in correspondence with the figure described above, we may also designate time  $t'$ , but then must of necessity, in connexion therewith, define space by the manifold of the three parameters  $x', y, z$ , in which case physical laws would be expressed in exactly the same way by means of  $x', y, z, t'$  as by means of  $x, y, z, t$ . We should then have in the world no longer *space*, but an infinite number of spaces,



analogously as there are in three-dimensional space an infinite number of planes. Three-dimensional geometry becomes a chapter in four-dimensional physics. Now you know why I said at the outset that space and time are to fade away into shadows, and only a world in itself will subsist.

## II

The question now is, what are the circumstances which force this changed conception of space and time upon us? Does it actually never contradict experience? And finally, is it advantageous for describing phenomena?

Before going into these questions, I must make an important remark. If we have in any way individualized space and time, we have, as a world-line corresponding to a stationary substantial point, a straight line parallel to the axis of  $t$ ; corresponding to a substantial point in uniform motion, a straight line at an angle to the axis of  $t$ ; to a substantial point in varying motion, a world-line in some form of a curve. If at any world-point  $x, y, z, t$  we take the world-line passing through that point, and find it parallel to any radius vector  $OA'$  of the above-mentioned hyperboloidal sheet, we can introduce  $OA'$  as a new axis of time, and with the new concepts of space and time thus given, the substance at the world-point concerned appears as at rest. We will now introduce this fundamental axiom:—

*The substance at any world-point may always, with the appropriate determination of space and time, be looked upon as at rest.*

The axiom signifies that at any world-point the expression

$$c^2 dt^2 - dx^2 - dy^2 - dz^2$$

always has a positive value, or, what comes to the same thing, that any velocity  $v$  always proves less than  $c$ . Accordingly  $c$  would stand as the upper limit for all substantial velocities, and that is precisely what would reveal the deeper significance of the magnitude  $c$ . In this second form the first impression made by the axiom is not altogether pleasing. But we must bear in mind that a modified form of mechanics, in which the square root of this quadratic differential expression appears,

will now make its way, so that cases with a velocity greater than that of light will henceforward play only some such part as that of figures with imaginary co-ordinates in geometry.

Now the impulse and true motive for assuming the group  $G_c$  came from the fact that the differential equation for the propagation of light in empty space possesses that group  $G_c$ .\* On the other hand, the concept of rigid bodies has meaning only in mechanics satisfying the group  $G_\infty$ . If we have a theory of optics with  $G_c$ , and if on the other hand there were rigid bodies, it is easy to see that one and the same direction of  $t$  would be distinguished by the two hyperboloidal sheets appropriate to  $G_c$  and  $G_\infty$ , and this would have the further consequence, that we should be able, by employing suitable rigid optical instruments in the laboratory, to perceive some alteration in the phenomena when the orientation with respect to the direction of the earth's motion is changed. But all efforts directed towards this goal, in particular the famous interference experiment of Michelson, have had a negative result. To explain this failure, H. A. Lorentz set up an hypothesis, the success of which lies in this very invariance in optics for the group  $G_c$ . According to Lorentz any moving body must have undergone a contraction in the direction of its motion, and in fact with a velocity  $v$ , a contraction in the ratio

$$1 : \sqrt{1 - v^2/c^2}.$$

This hypothesis sounds extremely fantastical, for the contraction is not to be looked upon as a consequence of resistances in the ether, or anything of that kind, but simply as a gift from above,—as an accompanying circumstance of the circumstance of motion.

I will now show by our figure that the Lorentzian hypothesis is completely equivalent to the new conception of space and time, which, indeed, makes the hypothesis much more intelligible. If for simplicity we disregard  $y$  and  $z$ , and imagine a world of one spatial dimension, then a parallel band, upright like the axis of  $t$ , and another inclining to the axis of  $t$  (see Fig. 1)

\* An application of this fact in its essentials has already been given by W. Voigt, *Göttinger Nachrichten*, 1887, p. 41.

represent, respectively, the career of a body at rest or in uniform motion, preserving in each case a constant spatial extent. If  $OA'$  is parallel to the second band, we can introduce  $t'$  as the time, and  $x'$  as the space co-ordinate, and then the second body appears at rest, the first in uniform motion. We now assume that the first body, envisaged as at rest, has the length  $l$ , that is, the cross section  $PP$  of the first band on the axis of  $x$  is equal to  $l \cdot OC$ , where  $OC$  denotes the unit of measure on the axis of  $x$ ; and on the other hand, that the second body, envisaged as at rest, has the same length  $l$ , which then means that the cross section  $Q'Q'$  of the second band, measured parallel to the axis of  $x'$ , is equal to  $l \cdot OC'$ . We now have in these two bodies images of two equal Lorentzian electrons, one at rest and one in uniform motion. But if we retain the original co-ordinates  $x, t$ , we must give as the extent of the second electron the cross section of its appropriate band parallel to the axis of  $x$ . Now since  $Q'Q' = l \cdot OC'$ , it is evident that  $QQ = l \cdot OD'$ . If  $dx/dt$  for the second band is equal to  $v$ , an easy calculation gives

$$OD' = OC \sqrt{1 - v^2/c^2},$$

therefore also  $PP : QQ = 1 : \sqrt{1 - v^2/c^2}$ . But this is the meaning of Lorentz's hypothesis of the contraction of electrons in motion. If on the other hand we envisage the second electron as at rest, and therefore adopt the system of reference  $x' t'$ , the length of the first must be denoted by the cross section  $P'P'$  of its band parallel to  $OC'$ , and we should find the first electron in comparison with the second to be contracted in exactly the same proportion; for in the figure

$$P'P' : Q'Q' = OD : OC' = OD' : OC = QQ : PP.$$

Lorentz called the  $t'$  combination of  $x$  and  $t$  the local time of the electron in uniform motion, and applied a physical construction of this concept, for the better understanding of the hypothesis of contraction. But the credit of first recognizing clearly that the time of the one electron is just as good as that of the other, that is to say, that  $t$  and  $t'$  are to be treated identically, belongs to A. Einstein.\* Thus time, as a

\* A. Einstein, Ann. d. Phys., 17, 1905, p. 891; Jahrb. d. Radioaktivität und Elektronik, 4, 1907, p. 411.

concept unequivocally determined by phenomena, was first deposed from its high seat. Neither Einstein nor Lorentz made any attack on the concept of space, perhaps because in the above-mentioned special transformation, where the plane of  $x', t'$  coincides with the plane of  $x, t$ , an interpretation is possible by saying that the  $x$ -axis of space maintains its position. One may expect to find a corresponding violation of the concept of space appraised as another act of audacity on the part of the higher mathematics. Nevertheless, this further step is indispensable for the true understanding of the group  $G_c$ , and when it has been taken, the word *relativity-postulate* for the requirement of an invariance with the group  $G_c$  seems to me very feeble. Since the postulate comes to mean that only the four-dimensional world in space and time is given by phenomena, but that the projection in space and in time may still be undertaken with a certain degree of freedom, I prefer to call it the *postulate of the absolute world* (or briefly, the world-postulate).

### III

The world-postulate permits identical treatment of the four co-ordinates  $x, y, z, t$ . By this means, as I shall now show, the forms in which the laws of physics are displayed gain in intelligibility. In particular the idea of acceleration acquires a clear-cut character.

I will use a geometrical manner of expression, which suggests itself at once if we tacitly disregard  $z$  in the triplex  $x, y, z$ . I take any world-point  $O$  as the zero-point of space-time. The cone  $c^2t^2 - x^2 - y^2 - z^2 = 0$  with apex  $O$  (Fig. 2) consists of two parts, one with values  $t < 0$ , the other with values  $t > 0$ . The former, the front cone of  $O$ , consists, let us say, of all the world-points which "send light to  $O$ ," the latter, the back cone of  $O$ , of all the world-points which "receive light from  $O$ ." The territory bounded by the front cone alone, we may call "before"  $O$ , that which is bounded by the back cone alone, "after"  $O$ . The hyperboloidal sheet already discussed

$$F = c^2t^2 - x^2 - y^2 - z^2 = 1, \quad t > 0$$

lies after  $O$ . The territory between the cones is filled by the

one-sheeted hyperboloidal figures

$$-F = x^2 + y^2 + z^2 - c^2t^2 = k^2$$

for all constant positive values of  $k$ . We are specially interested in the hyperbolas with  $O$  as centre, lying on the latter figures. The single branches of these hyperbolas may be called briefly the internal hyperbolas with centre  $O$ . One of these branches, regarded as a world-line, would represent a motion which, for  $t = -\infty$  and  $t = +\infty$ , rises asymptotically to the velocity of light,  $c$ .

If we now, on the analogy of vectors in space, call a directed length in the manifold of  $x, y, z, t$  a vector, we have to distinguish between the time-like vectors with directions from  $O$  to the sheet  $+F = 1$ ,  $t > 0$ , and the space-like vectors




FIG. 2.

with directions from  $O$  to  $-F = 1$ . The time axis may run parallel to any vector of the former kind. Any world-point between the front and back cones of  $O$  can be arranged by means of the system of reference so as to be simultaneous with  $O$ , but also just as well so as to be earlier than  $O$  or later than  $O$ . Any world-point within the front cone of  $O$  is necessarily always before  $O$ ; any world-point within the back cone of  $O$  necessarily always after  $O$ . Corresponding to passing to the limit,  $c = \infty$ , there would be a complete flattening out of the wedge-shaped segment between the cones into the plane manifold  $t = 0$ . In the figures this segment is intentionally drawn with different widths.

We divide up any vector we choose, e.g. that from  $O$  to  $x, y, z, t$ , into the four components  $x, y, z, t$ . If the directions

of two vectors are, respectively, that of a radius vector OR from O to one of the surfaces  $\mp F = 1$ , and that of a tangent RS at the point R of the same surface, the vectors are said to be normal to one another. Thus the condition that the vectors with components  $x, y, z, t$  and  $x_1, y_1, z_1, t_1$  may be normal to each other is

$$c^2 tt_1 - xx_1 - yy_1 - zz_1 = 0.$$

For the measurement of vectors in different directions the units of measure are to be fixed by assigning to a space-like vector from O to  $-F = 1$  always the magnitude 1, and to a time-like vector from O to  $+F = 1, t > 0$  always the magnitude  $1/c$ .

If we imagine at a world-point P ( $x, y, z, t$ ) the world-line of a substantial point running through that point, the magnitude corresponding to the time-like vector  $dx, dy, dz, dt$  laid off along the line is therefore

$$d\tau = \frac{1}{c} \sqrt{c^2 dt^2 - dx^2 - dy^2 - dz^2}.$$

The integral  $\int d\tau = \tau$  of this amount, taken along the world-line from any fixed starting-point  $P_0$  to the variable end-point P, we call the proper time of the substantial point at P. On the world-line we regard  $x, y, z, t$ —the components of the vector OP—as functions of the proper time  $\tau$ ; denote their first differential coefficients with respect to  $\tau$  by  $\dot{x}, \dot{y}, \dot{z}, \dot{t}$ ; their second differential coefficients with respect to  $\tau$  by  $\ddot{x}, \ddot{y}, \ddot{z}, \ddot{t}$ ; and give names to the appropriate vectors, calling the derivative of the vector OP with respect to  $\tau$  the velocity vector at P, and the derivative of this velocity vector with respect to  $\tau$  the acceleration vector at P. Hence, since

$$c^2 \dot{t}^2 - \dot{x}^2 - \dot{y}^2 - \dot{z}^2 = c^2,$$

we have

$$c^2 \ddot{t} \dot{t} - \dot{x} \ddot{x} - \dot{y} \ddot{y} - \dot{z} \ddot{z} = 0,$$

i.e. the velocity vector is the time-like vector of unit magnitude in the direction of the world-line at P, and the acceleration vector at P is normal to the velocity vector at P, and is therefore in any case a space-like vector.

Now, as is readily seen, there is a definite hyperbola

which has three infinitely proximate points in common with the world-line at P, and whose asymptotes are generators of a "front cone" and a "back cone" (Fig. 3).




FIG. 8.

Let this hyperbola be called the hyperbola of curvature at P. If M is the centre of this hyperbola, we here have to do with an internal hyperbola with centre M. Let  $\rho$  be the magnitude of the vector MP; then we recognize the acceleration vector at P as the vector in the direction MP of magnitude  $c^2/\rho$ .

If  $\ddot{x}, \ddot{y}, \ddot{z}, \ddot{t}$  are all zero, the hyperbola of curvature reduces to the straight line touching the world-line in P, and we must put  $\rho = \infty$ .

#### IV

To show that the assumption of group  $G_c$  for the laws of physics never leads to a contradiction, it is unavoidable to undertake a revision of the whole of physics on the basis of this assumption. This revision has to some extent already been successfully carried out for questions of thermodynamics and heat radiation,\* for electromagnetic processes, and finally, with the retention of the concept of mass, for mechanics.†

For this last branch of physics it is of prime importance to raise the question—When a force with the components X, Y, Z parallel to the axes of space acts at a world-point P ( $x, y, z, t$ ), where the velocity vector is  $\dot{x}, \dot{y}, \dot{z}, \dot{t}$ , what must we take this force to be when the system of reference is in any way changed? Now there exist certain approved statements as to the ponderomotive force in the electromagnetic field in the cases where the group  $G_c$  is undoubtedly admissible. These statements lead up to the simple rule:—When the system of reference is changed, the force in question transforms into a force in the new space co-ordinates in such a way that the appropriate vector with the components  $\dot{t}X$ ,

\* M. Planck, "Zur Dynamik bewegter Systeme," Berliner Berichte, 1907, p. 542; also in Ann. d. Phys., 26, 1908, p. 1.

† H. Minkowski, "Die Grundgleichungen für die elektromagnetischen Vorgänge in bewegten Körpern," Göttinger Nachrichten, 1908, p. 58.

$tY, tZ, tT$ , where

$$T = \frac{1}{c^2} \left( \frac{\dot{x}}{t} X + \frac{\dot{y}}{t} Y + \frac{\dot{z}}{t} Z \right)$$

is the rate at which work is done by the force at the world-point divided by  $c$ , remains unchanged. This vector is always normal to the velocity vector at  $P$ . A force vector of this kind, corresponding to a force at  $P$ , is to be called a "motive force vector" at  $P$ .

I shall now describe the world-line of a substantial point with constant mechanical mass  $m$ , passing through  $P$ . Let the velocity vector at  $P$ , multiplied by  $m$ , be called the "momentum vector" at  $P$ , and the acceleration vector at  $P$ , multiplied by  $m$ , be called the "force vector" of the motion at  $P$ . With these definitions, the law of motion of a point of mass with given motive force vector runs thus:—\* *The Force Vector of Motion is Equal to the Motive Force Vector*. This assertion comprises four equations for the components corresponding to the four axes, and since both vectors mentioned are *a priori* normal to the velocity vector, the fourth equation may be looked upon as a consequence of the other three. In accordance with the above signification of  $T$ , the fourth equation undoubtedly represents the law of energy. Therefore the component of the momentum vector along the axis of  $t$ , multiplied by  $c$ , is to be defined as the kinetic energy of the point mass. The expression for this is

$$mc^2 \frac{dt}{d\tau} = mc^2 / \sqrt{1 - v^2/c^2},$$

i.e., after removal of the additive constant  $mc^2$ , the expression  $\frac{1}{2}mv^2$  of Newtonian mechanics down to magnitudes of the order  $1/c^2$ . It comes out very clearly in this way, how the energy depends on the system of reference. But as the axis of  $t$  may be laid in the direction of any time-like vector, the law of energy, framed for all possible systems of reference, already contains, on the other hand, the whole system of the equations of motion. At the limiting transition which we have discussed, to  $c = \infty$ , this fact retains its importance for

\* H. Minkowski, loc. cit., p. 107. Cf. also M. Planck, Verhandlungen der physikalischen Gesellschaft, 4, 1906, p. 186.





sents by its components along the axes of  $x, y, z$ , the vector potential multiplied by  $c$ , and by the component along the axis of  $t$ , the scalar potential of the field excited by  $e$  at the world-point  $P$ . Herein lie the elementary laws formulated by A. Liénard and E. Wiechert.\*

Then in the description of the field produced by the electron we see that the separation of the field into electric and magnetic force is a relative one with regard to the underlying time axis; the most perspicuous way of describing the two forces together is on a certain analogy with the wrench in mechanics, though the analogy is not complete.

I will now describe the ponderomotive action of a moving point charge on another moving point charge. Let us imagine the world-line of a second point electron of the charge  $e_1$ , passing through the world-point  $P_1$ . We define  $P, Q, r$  as before, then construct (Fig. 4) the centre  $M$  of the hyperbola of curvature at  $P$ , and finally the normal  $MN$  from  $M$  to a straight line imagined through  $P$  parallel to  $QP_1$ . With  $P$  as starting-point we now determine a system of reference as follows:—The axis of  $t$  in the direction  $PQ$ , the axis of  $x$  in direction  $QP_1$ , the axis of  $y$  in direction  $MN$ , whereby finally the direction of the axis of  $z$  is also defined as normal to the axes of  $t, x, y$ . Let the acceleration vector at  $P$  be  $\ddot{x}, \ddot{y}, \ddot{z}, \dot{t}$ , the velocity vector at  $P_1$  be  $\dot{x}_1, \dot{y}_1, \dot{z}_1, \dot{t}_1$ . The motive force vector exerted at  $P_1$  by the first moving electron  $e$  on the second moving electron  $e_1$  now takes the form

$$- ee_1 \left( \dot{t}_1 - \frac{\dot{x}_1}{c} \right) \mathfrak{F},$$

where the components  $\mathfrak{F}_x, \mathfrak{F}_y, \mathfrak{F}_z, \mathfrak{F}_t$  of the vector  $\mathfrak{F}$  satisfy the three relations

$$c\mathfrak{F}_t - \mathfrak{F}_x = \frac{1}{r^2}, \quad \mathfrak{F}_y = \frac{\ddot{y}}{c^2 r}, \quad \mathfrak{F}_z = 0,$$

and where, fourthly, this vector  $\mathfrak{F}$  is normal to the velocity vector at  $P_1$ , and through this circumstance alone stands in dependence on the latter velocity vector.

\* A. Liénard, "Champ électrique et magnétique produit par une charge concentrée en un point et animée d'un mouvement quelconque," *L'Eclairage Electrique*, 16, 1898, pp. 5, 53, 106; E. Wiechert, "Elektrodynamische Elementargesetze," *Arch. Néerl.* (2), 5, 1900, p. 549.

When we compare this statement with previous formulations\* of the same elementary law of the ponderomotive action of moving point charges on one another, we are compelled to admit that it is only in four dimensions that the relations here taken under consideration reveal their inner being in full simplicity, and that on a three dimensional space forced upon us *a priori* they cast only a very complicated projection.

In mechanics as reformed in accordance with the world-postulate, the disturbing lack of harmony between Newtonian mechanics and modern electrodynamics disappears of its own accord. Before concluding I will just touch upon the attitude of Newton's law of attraction toward this postulate. I shall assume that when two points of mass  $m, m_1$  describe their world-lines, a motive force vector is exerted by  $m$  on  $m_1$ , of exactly the same form as that just given in the case of electrons, except that  $+mm_1$  must now take the place of  $-ee_1$ . We now specially consider the case where the acceleration vector of  $m$  is constantly zero. Let us then introduce  $t$  in such a way that  $m$  is to be taken as at rest, and let only  $m_1$  move under the motive force vector which proceeds from  $m$ . If we now modify this given vector in the first place by adding the factor  $t^{-1} = \sqrt{1 - v^2/c^2}$ , which, to the order of  $1/c^2$ , is equal to 1, it will be seen † that for the positions  $x_1, y_1, z_1$ , of  $m_1$  and their variations in time, we should arrive exactly at Kepler's laws again, except that the proper times  $\tau_1$  of  $m_1$  would take the place of the times  $t_1$ . From this simple remark it may then be seen that the proposed law of attraction combined with the new mechanics is no less well adapted to explain astronomical observations than the Newtonian law of attraction combined with Newtonian mechanics.

The fundamental equations for electromagnetic processes in ponderable bodies also fit in completely with the world-postulate. As I shall show elsewhere, it is not even by any means necessary to abandon the derivation of these funda-

\*K. Schwarzwald, Göttinger Nachr., 1908, p. 182; H. A. Lorentz, Enzykl. d. math. Wissensch., V, Art. 14, p. 199.

†H. Minkowski, loc. cit., p. 110.

mental equations from ideas of the electronic theory, as taught by Lorentz, in order to adapt them to the world-postulate.

The validity without exception of the world-postulate, I like to think, is the true nucleus of an electromagnetic image of the world, which, discovered by Lorentz, and further revealed by Einstein, now lies open in the full light of day. In the development of its mathematical consequences there will be ample suggestions for experimental verifications of the postulate, which will suffice to conciliate even those to whom the abandonment of old-established views is unsympathetic or painful, by the idea of a pre-established harmony between pure mathematics and physics.

## NOTES

by

A. SOMMERFELD

The following notes are given in an appendix so as to interfere in no way with Minkowski's text. They are by no means essential, having no other purpose than that of removing certain small formal mathematical difficulties which might hinder the comprehension of Minkowski's great thoughts. The bibliographical references are confined to the literature dealing expressly with the subject of his address. From the physical point of view there is nothing in what Minkowski says that must now be withdrawn, with the exception of the final remark on Newton's law of attraction. What will be the epistemological attitude towards Minkowski's conception of the time-space problem is another question, but, as it seems to me, a question which does not essentially touch his physics.

(1) Page 81, line 8. "On the other hand, the concept of rigid bodies has meaning only in mechanics satisfying the group  $G_{\infty}$ ." This sentence was confirmed in the widest sense in a discussion on a paper by his disciple M. Born, a year after Minkowski's death. Born (Ann. d. Physik, 80, 1909, p. 1) had defined a relatively rigid body as one in which every element of volume, even in accelerated motions, undergoes the Lorentzian contraction appropriate to its velocity. Ehrenfest (Phys. Zeitschr., 10, 1909, p. 918) showed that such a body cannot be set in rotation; Herglotz (Ann. d. Phys., 31, 1910, p. 393) and F. Nöther (Ann. d. Phys., 31, 1910, p. 919) that it has only three degrees of freedom of movement. The attempt was also made to define a relatively rigid body with six or nine degrees of freedom. But Planck (Phys. Zeitschr., 11, 1910, p. 294) expressed the view that the theory of relativity can operate only with more or less elastic bodies, and Laue (Phys. Zeitschr., 12, 1911, p. 48), employing Minkowski's methods, and his Fig. 2 in the text above, proved that in the theory of relativity every solid body must have an infinite number of degrees of freedom. Finally Herglotz (Ann. d. Physik, 36, 1911, p. 453) developed a relativistic theory of elasticity, according to which elastic tensions always occur if the motion of the body is not relatively rigid in Born's sense. Thus the relatively rigid body plays the same part in this theory of elasticity as the ordinary rigid body plays in the ordinary theory of elasticity.

(2) Page 82, line 18. "If  $da/dt$  for the second band is equal to  $v$ , an easy calculation gives  $OD' = OC\sqrt{1 - v^2/c^2}$ ." In Fig. 1, let  $\alpha = \angle A'OA$ ,  $\beta = \angle B'OA' = \angle C'OB'$ , in which the equality of the last two angles follows from the symmetrical position of the asymptotes with respect to the new axes of co-ordinates (conjugate diameters of the hyperbola).\* Since  $\alpha + \beta = \frac{1}{2}\pi$ ,

$$\sin 2\beta = \cos 2\alpha.$$

\* Sommerfeld seems to take  $ct$  as a co-ordinate in the graph in place of  $t$  as used by Minkowski.—TRANS.

In the triangle  $OD'O'$  the law of sines gives

$$\frac{OD'}{OC'} = \frac{\sin 2\beta}{\cos \alpha} = \frac{\cos 2\alpha}{\cos \alpha}$$

or, as  $OC' = OA'$ ,

$$OD' = OA' \frac{\cos 2\alpha}{\cos \alpha} = OA' \cos \alpha (1 - \tan^2 \alpha) \quad . \quad . \quad (1)$$

If  $x, t$  are the co-ordinates of the point  $A'$  in the  $x, t$  system, and therefore  $x \cdot OA$  and  $ct \cdot OC = ct \cdot OA$  respectively are the corresponding distances from the axes of co-ordinates, we have

$$x \cdot OA = \sin \alpha \cdot OA', \quad ct \cdot OA = \cos \alpha \cdot OA', \quad \frac{x}{ct} = \tan \alpha = \frac{v}{c} \quad . \quad (2)$$

Inserting these values of  $x$  and  $ct$  in the equation of the hyperbola, we find

$$OA'^2 (\cos^2 \alpha - \sin^2 \alpha) = OA^2, \quad OA' = \frac{OA}{\cos \alpha \sqrt{1 - \tan^2 \alpha}} \quad . \quad (3)$$

therefore, on account of (1) and (2),

$$OD' = OA \sqrt{1 - \tan^2 \alpha} = OA \sqrt{1 - v^2/c^2}.$$

This, because  $OA = OC$ , is the formula to be proved.

Further, in the right-angled triangle  $ODC$ ,

$$OD = \frac{OC}{\cos \alpha} = \frac{OA}{\cos \alpha}.$$

Equation (3) may therefore be also written in this way,

$$OA' = \frac{OD}{\sqrt{1 - \tan^2 \alpha}} \text{ or } \frac{OD}{OA'} = \sqrt{1 - \frac{v^2}{c^2}}.$$

This, together with (4), gives the proportion,

$$OD : OA' = OD' : OA,$$

which, as  $OA' = OC'$  and  $OA = OC$ , is identical with

$$OD : OC' = OD' : OC$$

employed on page 82, line 29.

(3) Page 84, line 15. "Any world-point between the front and back cones of  $O$  can be arranged, by means of the system of reference, so as to be simultaneous with  $O$ , but also just as well so as to be earlier than  $O$ , or later than  $O$ ." M. Lane (Phys. Zeitschr., 12, 1911, p. 48) traces to this observation the proof of Einstein's theorem: In the theory of relativity no process of causality can be propagated with a velocity greater than that of light ("Signal velocity  $\leq c$ "). Assume that an event  $O$  causes another event  $P$ , and that the world-point  $P$  lies in the region between the cones of  $O$ . In this case the effect would have been conveyed from  $O$  to  $P$  with a velocity greater than that of light, relatively to the system of reference  $x, t$  in question, in which, of course, the effect  $P$  is assumed to be later than the cause  $O$ ,  $t_P > 0$ . But now, in accordance with the words quoted above, the system of reference may be changed, so that  $P$  comes to be earlier than  $O$ , that is to say, a system  $x', t'$  may be chosen in

infinitely many ways so that  $t'_P$  becomes  $< 0$ . This is irreconcilable with the idea of causality.  $P$  must therefore lie either "after"  $O$  or on the back cone of  $O$ , i.e. the velocity of propagation of a signal to be sent from  $O$ , which is to cause a second event at the world-point  $P$ , must of necessity be  $\leq c$ . (Of course it is possible, even in the theory of relativity, to define processes propagated with velocity greater than light. This can be done geometrically, for example, in a very simple way. But such processes can never serve as signals, i.e. it is impossible to introduce them arbitrarily and by them, for example, to set a relay in motion at a distant place. There may be e.g. optical media, in which the "velocity of light" is greater than  $c$ . But in that case what is understood by the velocity of light is the propagation of phases in an infinite periodic wave-train. These can never be used for signalling. On the other hand a wave-front is propagated, in all circumstances and with any constitution of the optical medium, with the velocity  $c$ ; cf. e.g. A. Sommerfeld, "Festschrift Heinrich Weber," Leipzig, Teubner, 1912, p. 338, or Ann. d. Physik, 44, 1914, p. 177.

(4) Page 85, line 18. As Minkowski once remarked to me, the element of proper time  $d\tau$  is not a complete differential. Thus if we connect two world-points  $O$  and  $P$  by two different world-lines 1 and 2, then

$$\int_1 d\tau \neq \int_2 d\tau.$$

If 1 runs parallel to the  $t$ -axis, so that the first transition in the chosen system of reference signifies rest, it is evident that

$$\int_1 d\tau = t, \quad \int_2 d\tau < t.$$

On this depends the retardation of the moving clock compared with the clock at rest. The assertion is based, as Einstein has pointed out, on the unprovable assumption that the clock in motion actually indicates its own proper time, i.e. that it always gives the time corresponding to the state of velocity, regarded as constant, at any instant. The moving clock must naturally have been moved with acceleration (with changes of speed or direction) in order to be compared with the stationary clock at the world-point  $P$ . The retardation of the moving clock does not therefore actually indicate "motion," but "accelerated motion." Hence this does not contradict the principle of relativity.

(5) Page 86, line 4. The term "hyperbola of curvature" is formed exactly on the model of the elementary concept of the circle of curvature. The analogy becomes analytical identity if instead of the real co-ordinate of time  $t$  the imaginary  $u = ict$  is employed, that is,  $c$  times the co-ordinate employed by Minkowski, page 88, line 6.

By page 84 an internal hyperbola in the  $x, t$ -plane has the equation, with  $k = \rho$ ,

$$x^2 - c^2 t^2 = \rho^2,$$

therefore in the  $x, u$  plane

$$x^2 + u^2 = \rho^2.$$

Hence it may be written in parametric form, when  $\phi$  denotes a purely imaginary angle,

$$x = \rho \cos \phi, \quad u = \rho \sin \phi.$$

So, as I suggested in the *Ann. d. Phys.*, 33, p. 649, § 8, hyperbolic motion may also be denoted as "cyclic motion," whereby its chief properties (convection of the field, occurrence of a kind of centrifugal force) are characterized with particular clearness. For the hyperbolic motion we have

$$\dot{d}\tau = \frac{1}{c} \sqrt{(-du^2 - dx^2)} = \frac{c}{c} |d\phi|$$

and thus

$$\dot{x} = \frac{dx}{d\tau} = -ic \sin \phi, \quad \dot{u} = \frac{du}{d\tau} = +ic \cos \phi$$

$$\ddot{x} = \frac{d\dot{x}}{d\tau} = \frac{c^2}{\rho} \cos \phi, \quad \ddot{u} = \frac{d\dot{u}}{d\tau} = \frac{c^2}{\rho} \sin \phi.$$

The magnitude of the acceleration vector in hyperbolic motion is therefore  $c^2/\rho$ . Since any given world-line is touched by the hyperbola of curvature at three points, it has the same acceleration vector as the hyperbolic motion, and its magnitude is  $c^2/\rho$ , as indicated on page 86, line 11.

The centre M of the cyclic motion  $x^2 + u^2 = \rho^2$  is evidently the point  $x = 0, u = 0$ , and from this centre all points of the hyperbola have the constant "distance," i.e. a constant magnitude of the radius vector. Therefore  $\rho$  denotes the interval marked MP in Fig. 8.

(6) Page 87, line 1. A force X, Y, Z, to be made into a "force vector," must be multiplied by  $\dot{t} = dt/d\tau$ . This may be explained as follows.

According to Minkowski, page 87, line 10, the momentum vector is defined by  $m\dot{x}, m\dot{y}, m\dot{z}, m\dot{t}$ , where  $m$  denotes the "constant mechanical mass," or, as Minkowski says more plainly elsewhere, the "rest mass." If we retain Newton's law of motion (time rate of change of momentum equal to force), we have to set

$$\frac{d}{dt}(m\dot{x}) = X, \quad \frac{d}{dt}(m\dot{y}) = Y, \quad \frac{d}{dt}(m\dot{z}) = Z.$$

Multiplication by  $\dot{t}$  makes the left-hand sides into vector components in Minkowski's sense. Therefore  $\dot{t}X, \dot{t}Y, \dot{t}Z$  are also the first three components of the "force vector." The fourth component T follows without ambiguity from the requirement that the force vector is to be normal to the motion vector. Minkowski's equations for the mechanics of the mass point are therefore, with constant rest mass,

$$m\ddot{x} = \dot{t}X, \quad m\ddot{y} = \dot{t}Y, \quad m\ddot{z} = \dot{t}Z, \quad m\ddot{t} = \dot{t}T.$$

The assumption of constancy of rest mass can only be maintained, however, when the energy-content of the body is not changed in its motion, or in the words of Planck, when the motion ensues "adiabatically and isochorically."

(7) Pages 88 and 89. What is characteristic of the constructions here given, is their complete independence of any special system of reference. They give, as Minkowski postulates on page 88, "reciprocal relations between world-lines" (or world-points) as "the most perfect expression of physical laws." On page 89, for example, the electrodynamic potential (four-potential) is not referred to the axes of co-ordinates  $x, y, z, t$  until it is to be conventionally divided into a scalar and a vector portion, which have no independent invariant meaning from the relativistic standpoint.

By way of commentary to Minkowski I have deduced, from Maxwell's



equations, by Minkowski's methods, an invariant analytical form for the four-potential and the ponderomotive action between two electrons, and so given another view of these constructions of Minkowski. Instead of going into details here, I may refer to my article in *Ann. d. Phys.*, 33, 1910, p. 649, § 7, or to M. Laue, "Das Relativitätsprinzip," Braunschweig, Vieweg, 1913, § 19. Compare also Minkowski's address on the principle of relativity, edited by myself, in *Ann. d. Phys.*, 47, 1915, p. 927, where the four-potential is placed at the head of electrodynamics, and this theory thus reduced to its simplest form.

(8) Page 89, line 6. The invariant representation of the electromagnetic field by a "vector of the second kind" (or, as I proposed to call it, a "six-vector," a term which seems to be winning acceptance) is a particularly important part of Minkowski's view of electrodynamics. Whereas Minkowski's ideas on the vector of the first kind, or four-vector, were in part anticipated by Poincaré (*Rend. Circ. Mat. Palermo*, 21, 1906), the introduction of the six-vector is new. Like the six-vector, the wrench of mechanics (standing for a single force and a couple) depends on six independent parameters. And as in the electromagnetic field "the separation into electric and magnetic force is a relative one," so with the wrench, as is well known, the division into single force and couple can be made in very many ways.

(9) Page 90, line 9. Minkowski's relativistic form of Newton's law for the special case of zero acceleration mentioned in the text is included in the more general form proposed by Poincaré (*loc. cit.*). On the other hand, in taking acceleration into consideration, it goes further than the latter. Minkowski's or Poincaré's formulation of the law of gravitation shows that it is possible in many ways to reconcile Newton's law with the theory of relativity. That law is viewed as a point law, and gravitation therefore in a certain sense as action at a distance. The general theory of relativity, which Einstein has been developing from 1907 on, gets a deeper grip of the problem of gravitation. Gravitation is not only regarded as a field action and described by space-time differential equations—which seems from the present standpoint irrefutable—but it is also united organically with the principle of relativity extended to any transformations, whereas Minkowski and Poincaré had adapted it to the postulate of relativity in a more external manner. In the general theory of relativity the space-time structure is determined, from or together with, gravitation. Thus the principle of relativity, by an extension of Minkowski's ideas, is so formulated that it postulates the co-variance of physical quantities with reference to all point transformations, so that the coefficients of the invariant linear element enter into the laws of physics.

(10) Page 90, line 83. The "fundamental equations for electromagnetic processes in ponderable bodies" are developed by Minkowski in *Göttinger Nachrichten*, 1907. It was not granted him to complete the "deduction of this equation on the basis of the theory of electrons." His essays in this direction have been worked out by M. Born, and together with the "Fundamental Equations" make up the first volume of the series of monographs edited by Otto Blumenthal (Leipzig, 1910).

# ON THE INFLUENCE OF GRAVITATION ON THE PROPAGATION OF LIGHT

BY

A. EINSTEIN

*Translated from "Über den Einfluss der Schwerkraft auf  
die Ausbreitung des Lichtes," Annalen der Physik, 35,  
1911.*



# ON THE INFLUENCE OF GRAVITATION ON THE PROPAGATION OF LIGHT

By A. EINSTEIN

**I**N a memoir published four years ago \* I tried to answer the question whether the propagation of light is influenced by gravitation. I return to this theme, because my previous presentation of the subject does not satisfy me, and for a stronger reason, because I now see that one of the most important consequences of my former treatment is capable of being tested experimentally. For it follows from the theory here to be brought forward, that rays of light, passing close to the sun, are deflected by its gravitational field, so that the angular distance between the sun and a fixed star appearing near to it is apparently increased by nearly a second of arc.

In the course of these reflexions further results are yielded which relate to gravitation. But as the exposition of the entire group of considerations would be rather difficult to follow, only a few quite elementary reflexions will be given in the following pages, from which the reader will readily be able to inform himself as to the suppositions of the theory and its line of thought. The relations here deduced, even if the theoretical foundation is sound, are valid only to a first approximation.

## § 1. A Hypothesis as to the Physical Nature of the Gravitational Field

In a homogeneous gravitational field (acceleration of gravity  $\gamma$ ) let there be a stationary system of co-ordinates  $K$ , orientated so that the lines of force of the gravitational field run in the negative direction of the axis of  $z$ . In a space free

\* A. Einstein, Jahrbuch für Radioakt. und Elektronik, 4, 1907.

of gravitational fields let there be a second system of co-ordinates  $K'$ , moving with uniform acceleration ( $\gamma$ ) in the positive direction of its axis of  $z$ . To avoid unnecessary complications, let us for the present disregard the theory of relativity, and regard both systems from the customary point of view of kinematics, and the movements occurring in them from that of ordinary mechanics.

Relatively to  $K$ , as well as relatively to  $K'$ , material points which are not subjected to the action of other material points, move in keeping with the equations

$$\frac{d^2x}{dt^2} = 0, \quad \frac{d^2y}{dt^2} = 0, \quad \frac{d^2z}{dt^2} = -\gamma.$$

For the accelerated system  $K'$  this follows directly from Galileo's principle, but for the system  $K$ , at rest in a homogeneous gravitational field, from the experience that all bodies in such a field are equally and uniformly accelerated. This experience, of the equal falling of all bodies in the gravitational field, is one of the most universal which the observation of nature has yielded; but in spite of that the law has not found any place in the foundations of our edifice of the physical universe.

But we arrive at a very satisfactory interpretation of this law of experience, if we assume that the systems  $K$  and  $K'$  are physically exactly equivalent, that is, if we assume that we may just as well regard the system  $K$  as being in a space free from gravitational fields, if we then regard  $K$  as uniformly accelerated. This assumption of exact physical equivalence makes it impossible for us to speak of the absolute acceleration of the system of reference, just as the usual theory of relativity forbids us to talk of the absolute velocity of a system;\* and it makes the equal falling of all bodies in a gravitational field seem a matter of course.

As long as we restrict ourselves to purely mechanical processes in the realm where Newton's mechanics holds sway, we are certain of the equivalence of the systems  $K$  and  $K'$ .

\* Of course we cannot replace any arbitrary gravitational field by a state of motion of the system without a gravitational field, any more than, by a transformation of relativity, we can transform all points of a medium in any kind of motion to rest.

But this view of ours will not have any deeper significance unless the systems  $K$  and  $K'$  are equivalent with respect to all physical processes, that is, unless the laws of nature with respect to  $K$  are in entire agreement with those with respect to  $K'$ . By assuming this to be so, we arrive at a principle which, if it is really true, has great heuristic importance. For by theoretical consideration of processes which take place relatively to a system of reference with uniform acceleration, we obtain information as to the career of processes in a homogeneous gravitational field. We shall now show, first of all, from the standpoint of the ordinary theory of relativity, what degree of probability is inherent in our hypothesis.

## § 2. On the Gravitation of Energy

One result yielded by the theory of relativity is that the inertia mass of a body increases with the energy it contains; if the increase of energy amounts to  $E$ , the increase in inertia mass is equal to  $E/c^2$ , when  $c$  denotes the velocity of light. Now is there an increase of gravitating mass corresponding to this increase of inertia mass? If not, then a body would fall in the same gravitational field with varying acceleration according to the energy it contained. That highly satisfactory result of the theory of relativity by which the law of the conservation of mass is merged in the law of conservation of energy could not be maintained, because it would compel us to abandon the law of the conservation of mass in its old form for inertia mass, and maintain it for gravitating mass.

But this must be regarded as very improbable. On the other hand, the usual theory of relativity does not provide us with any argument from which to infer that the weight of a body depends on the energy contained in it. But we shall show that our hypothesis of the equivalence of the systems  $K$  and  $K'$  gives us gravitation of energy as a necessary consequence.

Let the two material systems  $S_1$  and  $S_2$ , provided with instruments of measurement, be situated on the  $z$ -axis of  $K$  at the distance  $h$  from each other,\* so that the gravitation potential in  $S_2$  is greater than that in  $S_1$  by  $\gamma h$ . Let a definite quantity

\* The dimensions of  $S_1$  and  $S_2$  are regarded as infinitely small in comparison with  $h$ .

of energy  $E$  be emitted from  $S_2$  towards  $S_1$ . Let the quantities of energy in  $S_1$  and  $S_2$  be measured by contrivances which—brought to one place in the system  $z$  and there compared—shall be perfectly alike. As to the process of this conveyance of energy by radiation we can make no *a priori* assertion, because we do not know the influence of the gravitational field on the radiation and the measuring instruments in  $S_1$  and  $S_2$ .

But by our postulate of the equivalence of  $K$  and  $K'$  we are able, in place of the system  $K$  in a homogeneous gravitational field, to set the gravitation-free system  $K'$ , which moves with uniform acceleration in the direction of positive  $z$ , and with the  $z$ -axis of which the material systems  $S_1$  and  $S_2$  are rigidly connected.

We judge of the process of the transference of energy by radiation from  $S_2$  to  $S_1$  from a system  $K_0$ , which is to be free from acceleration. At the moment when the radiation energy  $E_2$  is emitted from  $S_2$  toward  $S_1$ , let the velocity of  $K'$  relatively to  $K_0$  be zero. The radiation will arrive at  $S_1$  when the time  $h/c$  has elapsed (to a first approximation). But at this moment the velocity of  $S_1$  relatively to  $K_0$  is  $\gamma h/c = v$ . Therefore by the ordinary theory of relativity the radiation arriving at  $S_1$  does not possess the energy  $E_2$ , but a greater energy  $E_1$ , which is related to  $E_2$  to a first approximation by the equation\*




FIG. 5.

$$E_1 = E_2 \left( 1 + \frac{v}{c} \right) = E_2 \left( 1 + \gamma \frac{h}{c^2} \right) \quad . \quad . \quad (1)$$

By our assumption exactly the same relation holds if the same process takes place in the system  $K$ , which is not accelerated, but is provided with a gravitational field. In this case we may replace  $\gamma h$  by the potential  $\Phi$  of the gravitation vector in  $S_2$ , if the arbitrary constant of  $\Phi$  in  $S_1$  is equated to zero. We then have the equation

$$E_1 = E_2 + \frac{E_2 \Phi}{c^2} \quad . \quad . \quad (1a)$$

\* See above, pp. 69-71.

This equation expresses the law of energy for the process under observation. The energy  $E_1$  arriving at  $S_1$  is greater than the energy  $E_2$ , measured by the same means, which was emitted in  $S_2$ , the excess being the potential energy of the mass  $E_2/c^2$  in the gravitational field. It thus proves that for the fulfilment of the principle of energy we have to ascribe to the energy  $E$ , before its emission in  $S_2$ , a potential energy due to gravity, which corresponds to the gravitational mass  $E/c^2$ . Our assumption of the equivalence of  $K$  and  $K'$  thus removes the difficulty mentioned at the beginning of this paragraph which is left unsolved by the ordinary theory of relativity.

The meaning of this result is shown particularly clearly if we consider the following cycle of operations:—

1. The energy  $E$ , as measured in  $S_2$ , is emitted in the form of radiation in  $S_2$  towards  $S_1$ , where, by the result just obtained, the energy  $E(1 + \gamma h/c^2)$ , as measured in  $S_1$ , is absorbed.

2. A body  $W$  of mass  $M$  is lowered from  $S_2$  to  $S_1$ , work  $M\gamma h$  being done in the process.

3. The energy  $E$  is transferred from  $S_1$  to the body  $W$  while  $W$  is in  $S_1$ . Let the gravitational mass  $M$  be thereby changed so that it acquires the value  $M'$ .

4. Let  $W$  be again raised to  $S_2$ , work  $M'\gamma h$  being done in the process.

5. Let  $E$  be transferred from  $W$  back to  $S_2$ .

The effect of this cycle is simply that  $S_1$  has undergone the increase of energy  $E\gamma h/c^2$ , and that the quantity of energy  $M'\gamma h - M\gamma h$  has been conveyed to the system in the form of mechanical work. By the principle of energy, we must therefore have

$$E\gamma \frac{h}{c^2} = M'\gamma h - M\gamma h,$$

or

$$M' - M = E/c^2. \quad . \quad . \quad . \quad (1b)$$

The increase in gravitational mass is thus equal to  $E/c^2$ , and therefore equal to the increase in inertia mass as given by the theory of relativity.

The result emerges still more directly from the equivalence



## 104 GRAVITATION AND LIGHT

of the systems  $K$  and  $K'$ , according to which the gravitational mass in respect of  $K$  is exactly equal to the inertia mass in respect of  $K'$ ; energy must therefore possess a gravitational mass which is equal to its inertia mass. If a mass  $M_0$  be suspended on a spring balance in the system  $K'$ , the balance will indicate the apparent weight  $M_0\gamma$  on account of the inertia of  $M_0$ . If the quantity of energy  $E$  be transferred to  $M_0$ , the spring balance, by the law of the inertia of energy, will indicate  $(M_0 + E/c^2)\gamma$ . By reason of our fundamental assumption exactly the same thing must occur when the experiment is repeated in the system  $K$ , that is, in the gravitational field.

### § 3. Time and the Velocity of Light in the Gravitational Field

If the radiation emitted in the uniformly accelerated system  $K'$  in  $S_2$  toward  $S_1$  had the frequency  $\nu_2$  relatively to the clock in  $S_2$ , then, relatively to  $S_1$ , at its arrival in  $S_1$  it no longer has the frequency  $\nu_2$  relatively to an identical clock in  $S_1$ , but a greater frequency  $\nu_1$ , such that to a first approximation

$$\nu_1 = \nu_2 \left( 1 + \gamma \frac{h}{c^2} \right). \quad (2)$$

For if we again introduce the unaccelerated system of reference  $K_0$ , relatively to which, at the time of the emission of light,  $K'$  has no velocity, then  $S_1$ , at the time of arrival of the radiation at  $S_1$ , has, relatively to  $K_0$ , the velocity  $\gamma h/c$ , from which, by Doppler's principle, the relation as given results immediately.

In agreement with our assumption of the equivalence of the systems  $K'$  and  $K$ , this equation also holds for the stationary system of co-ordinates  $K$ , provided with a uniform gravitational field, if in it the transference by radiation takes place as described. It follows, then, that a ray of light emitted in  $S_2$  with a definite gravitational potential, and possessing at its emission the frequency  $\nu_2$ —compared with a clock in  $S_2$ —will, at its arrival in  $S_1$ , possess a different frequency  $\nu_1$ —measured by an identical clock in  $S_1$ . For  $\gamma h$  we substitute the gravitational potential  $\Phi$  of  $S_2$ —that of  $S_1$

being taken as zero—and assume that the relation which we have deduced for the homogeneous gravitational field also holds for other forms of field. Then

$$\nu_1 = \nu_2 \left( 1 + \frac{\Phi}{c^2} \right) . . . . . (2a)$$

This result (which by our deduction is valid to a first approximation) permits, in the first place, of the following application. Let  $\nu_0$  be the vibration-number of an elementary light-generator, measured by a delicate clock at the same place. Let us imagine them both at a place on the surface of the Sun (where our  $S_2$  is located). Of the light there emitted, a portion reaches the Earth ( $S_1$ ), where we measure the frequency of the arriving light with a clock U in all respects resembling the one just mentioned. Then by (2a),

$$\nu = \nu_0 \left( 1 + \frac{\Phi}{c^2} \right),$$

where  $\Phi$  is the (negative) difference of gravitational potential between the surface of the Sun and the Earth. Thus according to our view the spectral lines of sunlight, as compared with the corresponding spectral lines of terrestrial sources of light, must be somewhat displaced toward the red, in fact by the relative amount

$$\frac{\nu_0 - \nu}{\nu_0} = - \frac{\Phi}{c^2} = 2 \cdot 10^{-6}$$

If the conditions under which the solar bands arise were exactly known, this shifting would be susceptible of measurement. But as other influences (pressure, temperature) affect the position of the centres of the spectral lines, it is difficult to discover whether the inferred influence of the gravitational potential really exists.\*

On a superficial consideration equation (2), or (2a), respectively, seems to assert an absurdity. If there is constant transmission of light from  $S_2$  to  $S_1$ , how can any other number of periods per second arrive in  $S_1$  than is emitted

\* L. F. Jewell (Journ. de Phys., 6, 1897, p. 84) and particularly Ch. Fabry and H. Boisson (Comptes rendus, 148, 1909, pp. 688-690) have actually found such displacements of fine spectral lines toward the red end of the spectrum, of the order of magnitude here calculated, but have ascribed them to an effect of pressure in the absorbing layer.

in  $S_2$ ? But the answer is simple. We cannot regard  $\nu_2$  or respectively  $\nu_1$  simply as frequencies (as the number of periods per second) since we have not yet determined the time in system K. What  $\nu_2$  denotes is the number of periods with reference to the time-unit of the clock U in  $S_2$ , while  $\nu_1$  denotes the number of periods per second with reference to the identical clock in  $S_1$ . Nothing compels us to assume that the clocks U in different gravitation potentials must be regarded as going at the same rate. On the contrary, we must certainly define the time in K in such a way that the number of wave crests and troughs between  $S_2$  and  $S_1$  is independent of the absolute value of time; for the process under observation is by nature a stationary one. If we did not satisfy this condition, we should arrive at a definition of time by the application of which time would merge explicitly into the laws of nature, and this would certainly be unnatural and unpractical. Therefore the two clocks in  $S_1$  and  $S_2$  do not both give the "time" correctly. If we measure time in  $S_1$  with the clock U, then we must measure time in  $S_2$  with a clock which goes  $1 + \Phi/c^2$  times more slowly than the clock U when compared with U at one and the same place. For when measured by such a clock the frequency of the ray of light which is considered above is at its emission in  $S_2$

$$\nu_2 \left( 1 + \frac{\Phi}{c^2} \right)$$

and is therefore, by (2a), equal to the frequency  $\nu_1$  of the same ray of light on its arrival in  $S_1$ .

This has a consequence which is of fundamental importance for our theory. For if we measure the velocity of light at different places in the accelerated, gravitation-free-system K', employing clocks U of identical constitution, we obtain the same magnitude at all these places. The same holds good, by our fundamental assumption, for the system K as well. But from what has just been said we must use clocks of unlike constitution, for measuring time at places with differing gravitation potential. For measuring time at a place which, relatively to the origin of the co-ordinates, has the gravitation potential  $\Phi$ , we must employ a clock which—when removed to the origin of co-ordinates—goes  $(1 + \Phi/c^2)$  times more slowly than the clock used for measuring time at

the origin of co-ordinates. If we call the velocity of light at the origin of co-ordinates  $c_0$ , then the velocity of light  $c$  at a place with the gravitation potential  $\Phi$  will be given by the relation

$$c = c_0 \left(1 + \frac{\Phi}{c^2}\right) \quad . \quad . \quad . \quad . \quad (3)$$

The principle of the constancy of the velocity of light holds good according to this theory in a different form from that which usually underlies the ordinary theory of relativity.

#### § 4. Bending of Light-Rays in the Gravitational Field

From the proposition which has just been proved, that the velocity of light in the gravitational field is a function of the place, we may easily infer, by means of Huyghens's principle, that light-rays propagated across a gravitational field undergo deflexion. For let  $E$  be a wave front of a plane light-wave at the time  $t$ , and let  $P_1$  and  $P_2$  be two points in that plane at




FIG. 6.

unit distance from each other.  $P_1$  and  $P_2$  lie in the plane of the paper, which is chosen so that the differential coefficient of  $\Phi$ , taken in the direction of the normal to the plane, vanishes, and therefore also that of  $c$ . We obtain the corresponding wave front at time  $t + dt$ , or, rather, its line of section with the plane of the paper, by describing circles round the points  $P_1$  and  $P_2$  with radii  $c_1 dt$  and  $c_2 dt$  respectively, where  $c_1$  and  $c_2$  denote the velocity of light at the points  $P_1$  and  $P_2$  respectively, and by drawing the tangent to these circles. The angle through which the light-ray is deflected in the path  $c dt$  is therefore

$$(c_1 - c_2)dt = - \frac{\partial c}{\partial n'} dt,$$

if we calculate the angle positively when the ray is bent toward the side of increasing  $n'$ . The angle of deflexion per unit of path of the light-ray is thus

$$- \frac{1}{c} \frac{\partial c}{\partial n'}, \text{ or by (3) } - \frac{1}{c^2} \frac{\partial \Phi}{\partial n'}.$$

Finally, we obtain for the deflexion which a light-ray experiences toward the side  $n'$  on any path ( $s$ ) the expression

$$\alpha = - \frac{1}{c^2} \int \frac{\partial \Phi}{\partial n'} ds \quad . \quad . \quad . \quad (4)$$

We might have obtained the same result by directly considering the propagation of a ray of light in the uniformly accelerated system  $K'$ , and transferring the result to the system  $K$ , and thence to the case of a gravitational field of any form.

By equation (4) a ray of light passing along by a heavenly body suffers a deflexion to the side of the diminishing gravitational potential, that is, on the side directed toward the heavenly body, of the magnitude

$$\alpha = \frac{1}{c^2} \int_{\theta = -\frac{1}{2}\pi}^{\theta = \frac{1}{2}\pi} \frac{kM}{r^2} \cos \theta ds = 2 \frac{kM}{c^2 \Delta}$$

where  $k$  denotes the constant of gravitation,  $M$  the mass of the heavenly body,  $\Delta$  the distance of the ray from the centre of the body. A ray of light going past the Sun would accordingly undergo deflexion to the amount of  $4 \cdot 10^{-6} = \cdot 83$  seconds of arc. The angular distance of the star from the centre of the Sun appears to be increased by this amount. As the fixed stars in the parts of the sky near the Sun are visible during total eclipses of the Sun, this consequence of the theory may be compared with experience. With the planet Jupiter the displacement to be expected reaches to about  $\frac{1}{100}$  of the amount




FIG. 7.

given. It would be a most desirable thing if astronomers would take up the question here raised. For apart from any theory there is the question whether it is possible with the equipment at present available to detect an influence of gravitational fields on the propagation of light.

# THE FOUNDATION OF THE GENERAL THEORY OF RELATIVITY

BY

A. EINSTEIN

*Translated from "Die Grundlage der allgemeinen Relativitätstheorie," Annalen der Physik, 49, 1916.*

---



# THE FOUNDATION OF THE GENERAL THEORY OF RELATIVITY

By A. EINSTEIN

## A. FUNDAMENTAL CONSIDERATIONS ON THE POSTULATE OF RELATIVITY

### § 1. Observations on the Special Theory of Relativity

THE special theory of relativity is based on the following postulate, which is also satisfied by the mechanics of Galileo and Newton.

If a system of co-ordinates  $K$  is chosen so that, in relation to it, physical laws hold good in their simplest form, the *same* laws also hold good in relation to any other system of co-ordinates  $K'$  moving in uniform translation relatively to  $K$ . This postulate we call the "special principle of relativity." The word "special" is meant to intimate that the principle is restricted to the case when  $K'$  has a motion of uniform translation relatively to  $K$ , but that the equivalence of  $K'$  and  $K$  does not extend to the case of non-uniform motion of  $K'$  relatively to  $K$ .

Thus the special theory of relativity does not depart from classical mechanics through the postulate of relativity, but through the postulate of the constancy of the velocity of light *in vacuo*, from which, in combination with the special principle of relativity, there follow, in the well-known way, the relativity of simultaneity, the Lorentzian transformation, and the related laws for the behaviour of moving bodies and clocks.

The modification to which the special theory of relativity has subjected the theory of space and time is indeed far-reaching, but one important point has remained unaffected.



For the laws of geometry, even according to the special theory of relativity, are to be interpreted directly as laws relating to the possible relative positions of solid bodies at rest; and, in a more general way, the laws of kinematics are to be interpreted as laws which describe the relations of measuring bodies and clocks. To two selected material points of a stationary rigid body there always corresponds a distance of quite definite length, which is independent of the locality and orientation of the body, and is also independent of the time. To two selected positions of the hands of a clock at rest relatively to the privileged system of reference there always corresponds an interval of time of a definite length, which is independent of place and time. We shall soon see that the general theory of relativity cannot adhere to this simple physical interpretation of space and time.

## § 2. The Need for an Extension of the Postulate of Relativity

In classical mechanics, and no less in the special theory of relativity, there is an inherent epistemological defect which was, perhaps for the first time, clearly pointed out by Ernst Mach. We will elucidate it by the following example:—Two fluid bodies of the same size and nature hover freely in space at so great a distance from each other and from all other masses that only those gravitational forces need be taken into account which arise from the interaction of different parts of the same body. Let the distance between the two bodies be invariable, and in neither of the bodies let there be any relative movements of the parts with respect to one another. But let either mass, as judged by an observer at rest relatively to the other mass, rotate with constant angular velocity about the line joining the masses. This is a verifiable relative motion of the two bodies. Now let us imagine that each of the bodies has been surveyed by means of measuring instruments at rest relatively to itself, and let the surface of  $S_1$  prove to be a sphere, and that of  $S_2$  an ellipsoid of revolution. Thereupon we put the question—What is the reason for this difference in the two bodies? No answer can

be admitted as epistemologically satisfactory,\* unless the reason given is an *observable fact of experience*. The law of causality has not the significance of a statement as to the world of experience, except when *observable facts* ultimately appear as causes and effects.

Newtonian mechanics does not give a satisfactory answer to this question. It pronounces as follows:—The laws of mechanics apply to the space  $R_1$ , in respect to which the body  $S_1$  is at rest, but not to the space  $R_2$ , in respect to which the body  $S_2$  is at rest. But the privileged space  $R_1$  of Galileo, thus introduced, is a merely *factitious* cause, and not a thing that can be observed. It is therefore clear that Newton's mechanics does not really satisfy the requirement of causality in the case under consideration, but only apparently does so, since it makes the factitious cause  $R_1$  responsible for the observable difference in the bodies  $S_1$  and  $S_2$ .

The only satisfactory answer must be that the physical system consisting of  $S_1$  and  $S_2$  reveals within itself no imaginable cause to which the differing behaviour of  $S_1$  and  $S_2$  can be referred. The cause must therefore lie *outside* this system. We have to take it that the general laws of motion, which in particular determine the shapes of  $S_1$  and  $S_2$ , must be such that the mechanical behaviour of  $S_1$  and  $S_2$  is partly conditioned, in quite essential respects, by distant masses which we have not included in the system under consideration. These distant masses and their motions relative to  $S_1$  and  $S_2$  must then be regarded as the seat of the causes (which must be susceptible to observation) of the different behaviour of our two bodies  $S_1$  and  $S_2$ . They take over the rôle of the factitious cause  $R_1$ . Of all imaginable spaces  $R_1$ ,  $R_2$ , etc., in any kind of motion relatively to one another, there is none which we may look upon as privileged *a priori* without reviving the above-mentioned epistemological objection. *The laws of physics must be of such a nature that they apply to systems of reference in any kind of motion.* Along this road we arrive at an extension of the postulate of relativity.

In addition to this weighty argument from the theory of

\* Of course an answer may be satisfactory from the point of view of epistemology, and yet be unsound physically, if it is in conflict with other experiences.

knowledge, there is a well-known physical fact which favours an extension of the theory of relativity. Let  $K$  be a Galilean system of reference, i.e. a system relatively to which (at least in the four-dimensional region under consideration) a mass, sufficiently distant from other masses, is moving with uniform motion in a straight line. Let  $K'$  be a second system of reference which is moving relatively to  $K$  in *uniformly accelerated* translation. Then, relatively to  $K'$ , a mass sufficiently distant from other masses would have an accelerated motion such that its acceleration and direction of acceleration are independent of the material composition and physical state of the mass.

Does this permit an observer at rest relatively to  $K'$  to infer that he is on a "really" accelerated system of reference? The answer is in the negative; for the above-mentioned relation of freely movable masses to  $K'$  may be interpreted equally well in the following way. The system of reference  $K'$  is unaccelerated, but the space-time territory in question is under the sway of a gravitational field, which generates the accelerated motion of the bodies relatively to  $K'$ .

This view is made possible for us by the teaching of experience as to the existence of a field of force, namely, the gravitational field, which possesses the remarkable property of imparting the same acceleration to all bodies.\* The mechanical behaviour of bodies relatively to  $K'$  is the same as presents itself to experience in the case of systems which we are wont to regard as "stationary" or as "privileged." Therefore, from the physical standpoint, the assumption readily suggests itself that the systems  $K$  and  $K'$  may both with equal right be looked upon as "stationary," that is to say, they have an equal title as systems of reference for the physical description of phenomena.

It will be seen from these reflexions that in pursuing the general theory of relativity we shall be led to a theory of gravitation, since we are able to "produce" a gravitational field merely by changing the system of co-ordinates. It will also be obvious that the principle of the constancy of the velocity of light *in vacuo* must be modified, since we easily

\* Eötvös has proved experimentally that the gravitational field has this property in great accuracy.

recognize that the path of a ray of light with respect to  $K'$  must in general be curvilinear, if with respect to  $K$  light is propagated in a straight line with a definite constant velocity.

### § 3. The Space-Time Continuum. Requirement of General Co-Variance for the Equations Expressing General Laws of Nature

In classical mechanics, as well as in the special theory of relativity, the co-ordinates of space and time have a direct physical meaning. To say that a point-event has the  $X_1$  co-ordinate  $x_1$  means that the projection of the point-event on the axis of  $X_1$ , determined by rigid rods and in accordance with the rules of Euclidean geometry, is obtained by measuring off a given rod (the unit of length)  $x_1$  times from the origin of co-ordinates along the axis of  $X_1$ . To say that a point-event has the  $X_4$  co-ordinate  $x_4 = t$ , means that a standard clock, made to measure time in a definite unit period, and which is stationary relatively to the system of co-ordinates and practically coincident in space with the point-event,\* will have measured off  $x_4 = t$  periods at the occurrence of the event.

This view of space and time has always been in the minds of physicists, even if, as a rule, they have been unconscious of it. This is clear from the part which these concepts play in physical measurements; it must also have underlain the reader's reflexions on the preceding paragraph (§ 2) for him to connect any meaning with what he there read. But we shall now show that we must put it aside and replace it by a more general view, in order to be able to carry through the postulate of general relativity, if the special theory of relativity applies to the special case of the absence of a gravitational field.

In a space which is free of gravitational fields we introduce a Galilean system of reference  $K(x, y, z, t)$ , and also a system of co-ordinates  $K'(x', y', z', t')$  in uniform rotation relatively to  $K$ . Let the origins of both systems, as well as their axes

\* We assume the possibility of verifying "simultaneity" for events immediately proximate in space, or—to speak more precisely—for immediate proximity or coincidence in space-time, without giving a definition of this fundamental concept.

of  $Z$ , permanently coincide. We shall show that for a space-time measurement in the system  $K'$  the above definition of the physical meaning of lengths and times cannot be maintained. For reasons of symmetry it is clear that a circle around the origin in the  $X, Y$  plane of  $K$  may at the same time be regarded as a circle in the  $X', Y'$  plane of  $K'$ . We suppose that the circumference and diameter of this circle have been measured with a unit measure infinitely small compared with the radius, and that we have the quotient of the two results. If this experiment were performed with a measuring-rod at rest relatively to the Galilean system  $K$ , the quotient would be  $\pi$ . With a measuring-rod at rest relatively to  $K'$ , the quotient would be greater than  $\pi$ . This is readily understood if we envisage the whole process of measuring from the "stationary" system  $K$ , and take into consideration that the measuring-rod applied to the periphery undergoes a Lorentzian contraction, while the one applied along the radius does not. Hence Euclidean geometry does not apply to  $K'$ . The notion of co-ordinates defined above, which presupposes the validity of Euclidean geometry, therefore breaks down in relation to the system  $K'$ . So, too, we are unable to introduce a time corresponding to physical requirements in  $K'$ , indicated by clocks at rest relatively to  $K'$ . To convince ourselves of this impossibility, let us imagine two clocks of identical constitution placed, one at the origin of co-ordinates, and the other at the circumference of the circle, and both envisaged from the "stationary" system  $K$ . By a familiar result of the special theory of relativity, the clock at the circumference—judged from  $K$ —goes more slowly than the other, because the former is in motion and the latter at rest. An observer at the common origin of co-ordinates, capable of observing the clock at the circumference by means of light, would therefore see it lagging behind the clock beside him. As he will not make up his mind to let the velocity of light along the path in question depend explicitly on the time, he will interpret his observations as showing that the clock at the circumference "really" goes more slowly than the clock at the origin. So he will be obliged to define time in such a way that the rate of a clock depends upon where the clock may be.

We therefore reach this result :—In the general theory of relativity, space and time cannot be defined in such a way that differences of the spatial co-ordinates can be directly measured by the unit measuring-rod, or differences in the time co-ordinate by a standard clock.

The method hitherto employed for laying co-ordinates into the space-time continuum in a definite manner thus breaks down, and there seems to be no other way which would allow us to adapt systems of co-ordinates to the four-dimensional universe so that we might expect from their application a particularly simple formulation of the laws of nature. So there is nothing for it but to regard all imaginable systems of co-ordinates, on principle, as equally suitable for the description of nature. This comes to requiring that:—

*The general laws of nature are to be expressed by equations which hold good for all systems of co-ordinates, that is, are co-variant with respect to any substitutions whatever (generally co-variant).*

It is clear that a physical theory which satisfies this postulate will also be suitable for the general postulate of relativity. For the sum of *all* substitutions in any case includes those which correspond to all relative motions of three-dimensional systems of co-ordinates. That this requirement of general co-variance, which takes away from space and time the last remnant of physical objectivity, is a natural one, will be seen from the following reflexion. All our space-time verifications invariably amount to a determination of space-time coincidences. If, for example, events consisted merely in the motion of material points, then ultimately nothing would be observable but the meetings of two or more of these points. Moreover, the results of our measurements are nothing but verifications of such meetings of the material points of our measuring instruments with other material points, coincidences between the hands of a clock and points on the clock dial, and observed point-events happening at the same place at the same time.

The introduction of a system of reference serves no other purpose than to facilitate the description of the totality of such coincidences. We allot to the universe four space-time variables  $x_1, x_2, x_3, x_4$  in such a way that for every point-event

there is a corresponding system of values of the variables  $x_1 \dots x_4$ . To two coincident point-events there corresponds one system of values of the variables  $x_1 \dots x_4$ , i.e. coincidence is characterized by the identity of the co-ordinates. If, in place of the variables  $x_1 \dots x_4$ , we introduce functions of them,  $x'_1, x'_2, x'_3, x'_4$ , as a new system of co-ordinates, so that the systems of values are made to correspond to one another without ambiguity, the equality of all four co-ordinates in the new system will also serve as an expression for the space-time coincidence of the two point-events. As all our physical experience can be ultimately reduced to such coincidences, there is no immediate reason for preferring certain systems of co-ordinates to others, that is to say, we arrive at the requirement of general co-variance.

#### § 4. The Relation of the Four Co-ordinates to Measurement in Space and Time

It is not my purpose in this discussion to represent the general theory of relativity as a system that is as simple and logical as possible, and with the minimum number of axioms; but my main object is to develop this theory in such a way that the reader will feel that the path we have entered upon is psychologically the natural one, and that the underlying assumptions will seem to have the highest possible degree of security. With this aim in view let it now be granted that:—

For infinitely small four-dimensional regions the theory of relativity in the restricted sense is appropriate, if the co-ordinates are suitably chosen.

For this purpose we must choose the acceleration of the infinitely small ("local") system of co-ordinates so that no gravitational field occurs; this is possible for an infinitely small region. Let  $X_1, X_2, X_3$ , be the co-ordinates of space, and  $X_4$  the appertaining co-ordinate of time measured in the appropriate unit.\* If a rigid rod is imagined to be given as the unit measure, the co-ordinates, with a given orientation of the system of co-ordinates, have a direct physical meaning

\* The unit of time is to be chosen so that the velocity of light *in vacuo* as measured in the "local" system of co-ordinates is to be equal to unity.

in the sense of the special theory of relativity. By the special theory of relativity the expression

$$ds^2 = -dX_1^2 - dX_2^2 - dX_3^2 + dX_4^2 \quad . \quad . \quad (1)$$

then has a value which is independent of the orientation of the local system of co-ordinates, and is ascertainable by measurements of space and time. The magnitude of the linear element pertaining to points of the four-dimensional continuum in infinite proximity, we call  $ds$ . If the  $ds$  belonging to the element  $dX_1 \dots dX_4$  is positive, we follow Minkowski in calling it time-like; if it is negative, we call it space-like.

To the "linear element" in question, or to the two infinitely proximate point-events, there will also correspond definite differentials  $dx_1 \dots dx_4$  of the four-dimensional co-ordinates of any chosen system of reference. If this system, as well as the "local" system, is given for the region under consideration, the  $dX_\nu$  will allow themselves to be represented here by definite linear homogeneous expressions of the  $dx_\sigma$ :—

$$dX_\nu = \sum_{\sigma} a_{\nu\sigma} dx_{\sigma} \quad . \quad . \quad . \quad (2)$$

Inserting these expressions in (1), we obtain

$$ds^2 = \sum_{\sigma\tau} g_{\sigma\tau} dx_{\sigma} dx_{\tau}, \quad . \quad . \quad . \quad (3)$$

where the  $g_{\sigma\tau}$  will be functions of the  $x_{\sigma}$ . These can no longer be dependent on the orientation and the state of motion of the "local" system of co-ordinates, for  $ds^2$  is a quantity ascertainable by rod-clock measurement of point-events infinitely proximate in space-time, and defined independently of any particular choice of co-ordinates. The  $g_{\sigma\tau}$  are to be chosen here so that  $g_{\sigma\tau} = g_{\tau\sigma}$ ; the summation is to extend over all values of  $\sigma$  and  $\tau$ , so that the sum consists of  $4 \times 4$  terms, of which twelve are equal in pairs.

The case of the ordinary theory of relativity arises out of the case here considered, if it is possible, by reason of the particular relations of the  $g_{\sigma\tau}$  in a finite region, to choose the system of reference in the finite region in such a way that the  $g_{\sigma\tau}$  assume the constant values



$$\left. \begin{array}{cccc} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & +1 \end{array} \right\} \cdot \cdot \cdot \quad (4)$$

We shall find hereafter that the choice of such co-ordinates is, in general, not possible for a finite region.

From the considerations of § 2 and § 3 it follows that the quantities  $g_{\sigma\tau}$  are to be regarded from the physical standpoint as the quantities which describe the gravitational field in relation to the chosen system of reference. For, if we now assume the special theory of relativity to apply to a certain four-dimensional region with the co-ordinates properly chosen, then the  $g_{\sigma\tau}$  have the values given in (4). A free material point then moves, relatively to this system, with uniform motion in a straight line. Then if we introduce new space-time co-ordinates  $x_1, x_2, x_3, x_4$ , by means of any substitution we choose, the  $g^{\sigma\tau}$  in this new system will no longer be constants, but functions of space and time. At the same time the motion of the free material point will present itself in the new co-ordinates as a curvilinear non-uniform motion, and the law of this motion will be independent of the nature of the moving particle. We shall therefore interpret this motion as a motion under the influence of a gravitational field. We thus find the occurrence of a gravitational field connected with a space-time variability of the  $g_{\sigma}$ . So, too, in the general case, when we are no longer able by a suitable choice of co-ordinates to apply the special theory of relativity to a finite region, we shall hold fast to the view that the  $g_{\sigma\tau}$  describe the gravitational field.

Thus, according to the general theory of relativity, gravitation occupies an exceptional position with regard to other forces, particularly the electromagnetic forces, since the ten functions representing the gravitational field at the same time define the metrical properties of the space measured.

## B. MATHEMATICAL AIDS TO THE FORMULATION OF GENERALLY COVARIANT EQUATIONS

Having seen in the foregoing that the general postulate of relativity leads to the requirement that the equations of

physics shall be covariant in the face of any substitution of the co-ordinates  $x_1 \dots x_4$ , we have to consider how such generally covariant equations can be found. We now turn to this purely mathematical task, and we shall find that in its solution a fundamental rôle is played by the invariant  $ds$  given in equation (3), which, borrowing from Gauss's theory of surfaces, we have called the "linear element."

The fundamental idea of this general theory of covariants is the following:—Let certain things ("tensors") be defined with respect to any system of co-ordinates by a number of functions of the co-ordinates, called the "components" of the tensor. There are then certain rules by which these components can be calculated for a new system of co-ordinates, if they are known for the original system of co-ordinates, and if the transformation connecting the two systems is known. The things hereafter called tensors are further characterized by the fact that the equations of transformation for their components are linear and homogeneous. Accordingly, all the components in the new system vanish, if they all vanish in the original system. If, therefore, a law of nature is expressed by equating all the components of a tensor to zero, it is generally covariant. By examining the laws of the formation of tensors, we acquire the means of formulating generally covariant laws.

### § 5. Contravariant and Covariant Four-vectors

*Contravariant Four-vectors.*—The linear element is defined by the four "components"  $dx_\nu$ , for which the law of transformation is expressed by the equation

$$dx'_\sigma = \sum_\nu \frac{\partial x'_\sigma}{\partial x_\nu} dx_\nu, \quad . \quad . \quad . \quad (5)$$

The  $dx'_\sigma$  are expressed as linear and homogeneous functions of the  $dx_\nu$ . Hence we may look upon these co-ordinate differentials as the components of a "tensor" of the particular kind which we call a contravariant four-vector. Any thing which is defined relatively to the system of co-ordinates by four quantities  $A^\nu$ , and which is transformed by the same law

$$A'^\sigma = \sum_\nu \frac{\partial x'_\sigma}{\partial x_\nu} A^\nu, \quad . \quad . \quad . \quad (5a)$$

we also call a contravariant four-vector. From (5a) it follows at once that the sums  $A^\sigma \pm B^\sigma$  are also components of a four-vector, if  $A^\sigma$  and  $B^\sigma$  are such. Corresponding relations hold for all "tensors" subsequently to be introduced. (Rule for the addition and subtraction of tensors.)

*Covariant Four-vectors.*—We call four quantities  $A_\nu$  the components of a covariant four-vector, if for any arbitrary choice of the contravariant four-vector  $B^\nu$

$$\sum_\nu A_\nu B^\nu = \text{Invariant} \quad . \quad . \quad . \quad (6)$$

The law of transformation of a covariant four-vector follows from this definition. For if we replace  $B^\nu$  on the right-hand side of the equation

$$\sum_\sigma A'_\sigma B'^\sigma = \sum_\nu A_\nu B^\nu$$

by the expression resulting from the inversion of (5a),

$$\sum_\sigma \frac{\partial x_\nu}{\partial x'_\sigma} B'^\sigma,$$

we obtain

$$\sum_\sigma B'^\sigma \sum_\nu \frac{\partial x_\nu}{\partial x'_\sigma} A_\nu = \sum_\sigma B'^\sigma A'_\sigma.$$

Since this equation is true for arbitrary values of the  $B'^\sigma$ , it follows that the law of transformation is

$$A'_\sigma = \sum_\nu \frac{\partial x_\nu}{\partial x'_\sigma} A_\nu \quad . \quad . \quad . \quad (7)$$

*Note on a Simplified Way of Writing the Expressions.*—A glance at the equations of this paragraph shows that there is always a summation with respect to the indices which occur twice under a sign of summation (e.g. the index  $\nu$  in (5)), and only with respect to indices which occur twice. It is therefore possible, without loss of clearness, to omit the sign of summation. In its place we introduce the convention:—If an index occurs twice in one term of an expression, it is always to be summed unless the contrary is expressly stated.

The difference between covariant and contravariant four-vectors lies in the law of transformation ((7) or (5) respectively). Both forms are tensors in the sense of the general remark above. Therein lies their importance. Following Ricci and

Levi-Civita, we denote the contravariant character by placing the index above, the covariant by placing it below.

### § 6. Tensors of the Second and Higher Ranks

*Contravariant Tensors.*—If we form all the sixteen products  $A^{\mu\nu}$  of the components  $A^\mu$  and  $B^\nu$  of two contravariant four-vectors

$$A^{\mu\nu} = A^\mu B^\nu \quad . \quad . \quad . \quad (8)$$

then by (8) and (5a)  $A^{\mu\nu}$  satisfies the law of transformation

$$A'^{\sigma\tau} = \frac{\partial x'_\sigma}{\partial x_\mu} \frac{\partial x'_\tau}{\partial x_\nu} A^{\mu\nu} \quad . \quad . \quad . \quad (9)$$

We call a thing which is described relatively to any system of reference by sixteen quantities, satisfying the law of transformation (9), a contravariant tensor of the second rank. Not every such tensor allows itself to be formed in accordance with (8) from two four-vectors, but it is easily shown that any given sixteen  $A^{\mu\nu}$  can be represented as the sums of the  $A^\mu B^\nu$  of four appropriately selected pairs of four-vectors. Hence we can prove nearly all the laws which apply to the tensor of the second rank defined by (9) in the simplest manner by demonstrating them for the special tensors of the type (8).

*Contravariant Tensors of Any Rank.*—It is clear that, on the lines of (8) and (9), contravariant tensors of the third and higher ranks may also be defined with  $4^3$  components, and so on. In the same way it follows from (8) and (9) that the contravariant four-vector may be taken in this sense as a contravariant tensor of the first rank.

*Covariant Tensors.*—On the other hand, if we take the sixteen products  $A_{\mu\nu}$  of two covariant four-vectors  $A_\mu$  and  $B_\nu$ ,

$$A_{\mu\nu} = A_\mu B_\nu, \quad . \quad . \quad . \quad (10)$$

the law of transformation for these is

$$A'_{\sigma\tau} = \frac{\partial x_\mu}{\partial x'_\sigma} \frac{\partial x_\nu}{\partial x'_\tau} A_{\mu\nu} \quad . \quad . \quad . \quad (11)$$

This law of transformation defines the covariant tensor of the second rank. All our previous remarks on contravariant tensors apply equally to covariant tensors.

NOTE.—It is convenient to treat the scalar (or invariant) both as a contravariant and a covariant tensor of zero rank.

*Mixed Tensors.*—We may also define a tensor of the second rank of the type

$$A^{\nu}_{\mu} = A_{\mu} B^{\nu} \quad . \quad . \quad . \quad . \quad . \quad (12)$$

which is covariant with respect to the index  $\mu$ , and contravariant with respect to the index  $\nu$ . Its law of transformation is

$$A'^{\tau}_{\sigma} = \frac{\partial x'_{\tau}}{\partial x_{\nu}} \frac{\partial x_{\mu}}{\partial x'_{\sigma}} A^{\nu}_{\mu} \quad . \quad . \quad . \quad . \quad . \quad (13)$$

Naturally there are mixed tensors with any number of indices of covariant character, and any number of indices of contravariant character. Covariant and contravariant tensors may be looked upon as special cases of mixed tensors.

*Symmetrical Tensors.*—A contravariant, or a covariant tensor, of the second or higher rank is said to be symmetrical if two components, which are obtained the one from the other by the interchange of two indices, are equal. The tensor  $A^{\mu\nu}$ , or the tensor  $A_{\mu\nu}$ , is thus symmetrical if for any combination of the indices  $\mu, \nu$ ,

$$A^{\mu\nu} = A^{\nu\mu}, \quad . \quad . \quad . \quad . \quad . \quad (14)$$

or respectively,

$$A_{\mu\nu} = A_{\nu\mu} \quad . \quad . \quad . \quad . \quad . \quad (14a)$$

It has to be proved that the symmetry thus defined is a property which is independent of the system of reference. It follows in fact from (9), when (14) is taken into consideration, that

$$A'^{\sigma\tau} = \frac{\partial x'_{\sigma}}{\partial x_{\mu}} \frac{\partial x'_{\tau}}{\partial x_{\nu}} A^{\mu\nu} = \frac{\partial x'_{\sigma}}{\partial x_{\mu}} \frac{\partial x'_{\tau}}{\partial x_{\nu}} A^{\nu\mu} = \frac{\partial x'_{\sigma}}{\partial x_{\nu}} \frac{\partial x'_{\tau}}{\partial x_{\mu}} A^{\mu\nu} = A'^{\tau\sigma}.$$

The last equation but one depends upon the interchange of the summation indices  $\mu$  and  $\nu$ , i.e. merely on a change of notation.

*Antisymmetrical Tensors.*—A contravariant or a covariant tensor of the second, third, or fourth rank is said to be antisymmetrical if two components, which are obtained the one from the other by the interchange of two indices, are equal and of opposite sign. The tensor  $A^{\mu\nu}$ , or the tensor  $A_{\mu\nu}$ , is therefore antisymmetrical, if always

$$A^{\mu\nu} = -A^{\nu\mu}, \quad (15)$$

or respectively,

$$A_{\mu\nu} = -A_{\nu\mu} \quad . \quad . \quad . \quad . \quad (15a)$$

Of the sixteen components  $A^{\mu\nu}$ , the four components  $A^{\mu\mu}$  vanish; the rest are equal and of opposite sign in pairs, so that there are only six components numerically different (a six-vector). Similarly we see that the antisymmetrical tensor of the third rank  $A^{\mu\nu\sigma}$  has only four numerically different components, while the antisymmetrical tensor  $A^{\mu\nu\sigma\tau}$  has only one. There are no antisymmetrical tensors of higher rank than the fourth in a continuum of four dimensions.

## § 7. Multiplication of Tensors

*Outer Multiplication of Tensors.*—We obtain from the components of a tensor of rank  $n$  and of a tensor of rank  $m$  the components of a tensor of rank  $n + m$  by multiplying each component of the one tensor by each component of the other. Thus, for example, the tensors  $T$  arise out of the tensors  $A$  and  $B$  of different kinds.

$$\begin{aligned} T_{\mu\nu\sigma} &= A_{\mu\nu}B_{\sigma}, \\ T^{\mu\nu\sigma} &= A^{\mu\nu}B^{\sigma}, \\ T^{\sigma}_{\mu\nu} &= A_{\mu\nu}B^{\sigma}. \end{aligned}$$

The proof of the tensor character of  $T$  is given directly by the representations (8), (10), (12), or by the laws of transformation (9), (11), (13). The equations (8), (10), (12) are themselves examples of outer multiplication of tensors of the first rank.

“*Contraction*” of a *Mixed Tensor*.—From any mixed tensor we may form a tensor whose rank is less by two, by equating an index of covariant with one of contravariant character, and summing with respect to this index (“contraction”). Thus, for example, from the mixed tensor of the fourth rank  $A^{\sigma\tau}_{\mu\nu}$ , we obtain the mixed tensor of the second rank,

$$A_{\nu}^{\tau} = A_{\mu\nu}^{\mu\tau} \quad (= \sum_{\mu} A_{\mu\nu}^{\mu\tau}),$$

and from this, by a second contraction, the tensor of zero rank.

$$A = A'_y = A^{\mu\nu}_{\mu\nu}$$

The proof that the result of contraction really possesses the tensor character is given either by the representation of a tensor according to the generalization of (12) in combination with (6), or by the generalization of (13).

*Inner and Mixed Multiplication of Tensors.*—These consist in a combination of outer multiplication with contraction.

*Examples.*—From the covariant tensor of the second rank  $A_{\mu\nu}$  and the contravariant tensor of the first rank  $B^\sigma$  we form by outer multiplication the mixed tensor

$$D_{\mu\nu}^\sigma = A_{\mu\nu}B^\sigma.$$

On contraction with respect to the indices  $\nu$  and  $\sigma$ , we obtain the covariant four-vector

$$D_\mu = D_{\mu\nu}^\nu = A_{\mu\nu}B^\nu.$$

This we call the inner product of the tensors  $A_{\mu\nu}$  and  $B^\sigma$ . Analogously we form from the tensors  $A_{\mu\nu}$  and  $B^{\sigma\tau}$ , by outer multiplication and double contraction, the inner product  $A_{\mu\nu}B^{\mu\nu}$ . By outer multiplication and one contraction, we obtain from  $A_{\mu\nu}$  and  $B^{\sigma\tau}$  the mixed tensor of the second rank  $D_\mu^\tau = A_{\mu\nu}B^{\nu\tau}$ . This operation may be aptly characterized as a mixed one, being "outer" with respect to the indices  $\mu$  and  $\tau$ , and "inner" with respect to the indices  $\nu$  and  $\sigma$ .

We now prove a proposition which is often useful as evidence of tensor character. From what has just been explained,  $A_{\mu\nu}B^{\mu\nu}$  is a scalar if  $A_{\mu\nu}$  and  $B^{\sigma\tau}$  are tensors. But we may also make the following assertion: If  $A_{\mu\nu}B^{\mu\nu}$  is a scalar for any choice of the tensor  $B^{\mu\nu}$ , then  $A_{\mu\nu}$  has tensor character. For, by hypothesis, for any substitution,

$$A'_{\sigma\tau}B'^{\sigma\tau} = A_{\mu\nu}B^{\mu\nu}.$$

But by an inversion of (9)

$$B^{\mu\nu} = \frac{\partial x_\mu}{\partial x'_\sigma} \frac{\partial x_\nu}{\partial x'_\tau} B'^{\sigma\tau}.$$

This, inserted in the above equation, gives

$$\left( A'_{\sigma\tau} - \frac{\partial x_\mu}{\partial x'_\sigma} \frac{\partial x_\nu}{\partial x'_\tau} A_{\mu\nu} \right) B'^{\sigma\tau} = 0.$$

This can only be satisfied for arbitrary values of  $B'^{\sigma\tau}$  if the

bracket vanishes. The result then follows by equation (11). This rule applies correspondingly to tensors of any rank and character, and the proof is analogous in all cases.

The rule may also be demonstrated in this form: If  $B^\mu$  and  $C^\nu$  are any vectors, and if, for all values of these, the inner product  $A_{\mu\nu}B^\mu C^\nu$  is a scalar, then  $A_{\mu\nu}$  is a covariant tensor. This latter proposition also holds good even if only the more special assertion is correct, that with any choice of the four-vector  $B^\mu$  the inner product  $A_{\mu\nu}B^\mu B^\nu$  is a scalar, if in addition it is known that  $A_{\mu\nu}$  satisfies the condition of symmetry  $A_{\mu\nu} = A_{\nu\mu}$ . For by the method given above we prove the tensor character of  $(A_{\mu\nu} + A_{\nu\mu})$ , and from this the tensor character of  $A_{\mu\nu}$  follows on account of symmetry. This also can be easily generalized to the case of covariant and contravariant tensors of any rank.

Finally, there follows from what has been proved, this law, which may also be generalized for any tensors: If for any choice of the four-vector  $B^\nu$  the quantities  $A_{\mu\nu}B^\nu$  form a tensor of the first rank, then  $A_{\mu\nu}$  is a tensor of the second rank. For, if  $C^\mu$  is any four-vector, then on account of the tensor character of  $A_{\mu\nu}B^\nu$ , the inner product  $A_{\mu\nu}B^\nu C^\mu$  is a scalar for any choice of the two four-vectors  $B^\nu$  and  $C^\mu$ . From which the proposition follows.

### § 8. Some Aspects of the Fundamental Tensor $g_{\mu\nu}$

*The Covariant Fundamental Tensor.*—In the invariant expression for the square of the linear element,

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu,$$

the part played by the  $dx_\mu$  is that of a contravariant vector which may be chosen at will. Since further,  $g_{\mu\nu} = g_{\nu\mu}$ , it follows from the considerations of the preceding paragraph that  $g_{\mu\nu}$  is a covariant tensor of the second rank. We call it the "fundamental tensor." In what follows we deduce some properties of this tensor which, it is true, apply to any tensor of the second rank. But as the fundamental tensor plays a special part in our theory, which has its physical basis in the peculiar effects of gravitation, it so happens that the relations to be developed are of importance to us only in the case of the fundamental tensor.



*The Contravariant Fundamental Tensor.*—If in the determinant formed by the elements  $g_{\mu\nu}$ , we take the co-factor of each of the  $g_{\mu\nu}$  and divide it by the determinant  $g = |g_{\mu\nu}|$ , we obtain certain quantities  $g^{\mu\nu} (= g^{\nu\mu})$  which, as we shall demonstrate, form a contravariant tensor.

By a known property of determinants

$$g_{\mu\sigma}g^{\nu\sigma} = \delta_{\mu}^{\nu} \quad . \quad . \quad . \quad (16)$$

where the symbol  $\delta_{\mu}^{\nu}$  denotes 1 or 0, according as  $\mu = \nu$  or  $\mu \neq \nu$ .

Instead of the above expression for  $ds^2$  we may thus write

$$g_{\mu\sigma}\delta_{\nu}^{\sigma}dx_{\mu}dx_{\nu}$$

or, by (16)

$$g_{\mu\sigma}g_{\nu\tau}g^{\sigma\tau}dx_{\mu}dx_{\nu}.$$

But, by the multiplication rules of the preceding paragraphs, the quantities

$$d\xi_{\sigma} = g_{\mu\sigma}dx_{\mu}$$

form a covariant four-vector, and in fact an arbitrary vector, since the  $dx_{\mu}$  are arbitrary. By introducing this into our expression we obtain

$$ds^2 = g^{\sigma\tau}d\xi_{\sigma}d\xi_{\tau}.$$

Since this, with the arbitrary choice of the vector  $d\xi_{\sigma}$ , is a scalar, and  $g^{\sigma\tau}$  by its definition is symmetrical in the indices  $\sigma$  and  $\tau$ , it follows from the results of the preceding paragraph that  $g^{\sigma\tau}$  is a contravariant tensor.

It further follows from (16) that  $\delta_{\mu}^{\nu}$  is also a tensor, which we may call the mixed fundamental tensor.

*The Determinant of the Fundamental Tensor.*—By the rule for the multiplication of determinants

$$|g_{\mu\alpha}g^{\alpha\nu}| = |g_{\mu\alpha}| \times |g^{\alpha\nu}|.$$

On the other hand

$$|g_{\mu\alpha}g^{\alpha\nu}| = |\delta_{\mu}^{\nu}| = 1.$$

It therefore follows that

$$|g_{\mu\nu}| \times |g^{\mu\nu}| = 1 \quad . \quad . \quad . \quad (17)$$

*The Volume Scalar.*—We seek first the law of transfor-

mation of the determinant  $g = |g_{\mu\nu}|$ . In accordance with (11)

$$g' = \left| \frac{\partial x_\mu}{\partial x'_\sigma} \frac{\partial x}{\partial x'_\tau} g_{\mu\nu} \right|.$$

Hence, by a double application of the rule for the multiplication of determinants, it follows that

$$g' = \left| \frac{\partial x_\mu}{\partial x'_\sigma} \right| \cdot \left| \frac{\partial x_\nu}{\partial x'_\tau} \right| \cdot |g_{\mu\nu}| = \left| \frac{\partial x_\mu}{\partial x'_\sigma} \right|^2 g,$$

or

$$\sqrt{g'} = \left| \frac{\partial x_\mu}{\partial x'_\sigma} \right| \sqrt{g}.$$

On the other hand, the law of transformation of the element of volume

$$d\tau = \int dx_1 dx_2 dx_3 dx_4$$

is, in accordance with the theorem of Jacobi,

$$d\tau' = \left| \frac{\partial x'_\sigma}{\partial x_\mu} \right| d\tau.$$

By multiplication of the last two equations, we obtain

$$\sqrt{g'} d\tau' = \sqrt{g} d\tau \quad . \quad . \quad . \quad (18).$$

Instead of  $\sqrt{g}$ , we introduce in what follows the quantity  $\sqrt{-g}$ , which is always real on account of the hyperbolic character of the space-time continuum. The invariant  $\sqrt{-g} d\tau$  is equal to the magnitude of the four-dimensional element of volume in the "local" system of reference, as measured with rigid rods and clocks in the sense of the special theory of relativity.

*Note on the Character of the Space-time Continuum.*—Our assumption that the special theory of relativity can always be applied to an infinitely small region, implies that  $ds^2$  can always be expressed in accordance with (1) by means of real quantities  $dX_1 \dots dX_4$ . If we denote by  $d\tau_0$  the "natural" element of volume  $dX_1, dX_2, dX_3, dX_4$ , then

$$d\tau_0 = \sqrt{-g} d\tau \quad . \quad . \quad . \quad (18a)$$

If  $\sqrt{-g}$  were to vanish at a point of the four-dimensional continuum, it would mean that at this point an infinitely small "natural" volume would correspond to a finite volume in the co-ordinates. Let us assume that this is never the case. Then  $g$  cannot change sign. We will assume that, in the sense of the special theory of relativity,  $g$  always has a finite negative value. This is a hypothesis as to the physical nature of the continuum under consideration, and at the same time a convention as to the choice of co-ordinates.

But if  $-g$  is always finite and positive, it is natural to settle the choice of co-ordinates *a posteriori* in such a way that this quantity is always equal to unity. We shall see later that by such a restriction of the choice of co-ordinates it is possible to achieve an important simplification of the laws of nature.

In place of (18), we then have simply  $d\tau' = d\tau$ , from which, in view of Jacobi's theorem, it follows that

$$\left| \frac{\partial x'_\sigma}{\partial x_\mu} \right| = 1 \quad . \quad . \quad . \quad (19)$$

Thus, with this choice of co-ordinates, only substitutions for which the determinant is unity are permissible.

But it would be erroneous to believe that this step indicates a partial abandonment of the general postulate of relativity. We do not ask "What are the laws of nature which are co-variant in face of all substitutions for which the determinant is unity?" but our question is "What are the generally co-variant laws of nature?" It is not until we have formulated these that we simplify their expression by a particular choice of the system of reference.

*The Formation of New Tensors by Means of the Fundamental Tensor.*—Inner, outer, and mixed multiplication of a tensor by the fundamental tensor give tensors of different character and rank. For example,

$$\begin{aligned} A^\mu &= g^{\mu\sigma} A_\sigma, \\ A &= g_{\mu\nu} A^{\mu\nu}. \end{aligned}$$

The following forms may be specially noted:—

$$\begin{aligned} A^{\mu\nu} &= g^{\mu\alpha} g^{\nu\beta} A_{\alpha\beta}, \\ A_{\mu\nu} &= g_{\mu\alpha} g_{\nu\beta} A^{\alpha\beta} \end{aligned}$$

the "complements" of covariant and contravariant tensors respectively), and

$$B_{\mu\nu} = g_{\mu\nu} g^{\alpha\beta} A_{\alpha\beta}.$$

We call  $B_{\mu\nu}$  the reduced tensor associated with  $A_{\mu\nu}$ . Similarly,

$$B^{\mu\nu} = g^{\mu\nu} g_{\alpha\beta} A^{\alpha\beta}.$$

It may be noted that  $g^{\mu\nu}$  is nothing more than the complement of  $g_{\mu\nu}$ , since

$$g^{\mu\alpha} g^{\nu\beta} g_{\alpha\beta} = g^{\mu\alpha} \delta_{\alpha}^{\nu} = g^{\mu\nu}.$$

### § 9. The Equation of the Geodetic Line. The Motion of a Particle

As the linear element  $ds$  is defined independently of the system of co-ordinates, the line drawn between two points P and P' of the four-dimensional continuum in such a way that  $\int ds$  is stationary—a geodetic line—has a meaning which also is independent of the choice of co-ordinates. Its equation is

$$\delta \int_P^{P'} ds = 0 \quad . \quad . \quad . \quad (20)$$

Carrying out the variation in the usual way, we obtain from this equation four differential equations which define the geodetic line; this operation will be inserted here for the sake of completeness. Let  $\lambda$  be a function of the co-ordinates  $x_\nu$ , and let this define a family of surfaces which intersect the required geodetic line as well as all the lines in immediate proximity to it which are drawn through the points P and P'. Any such line may then be supposed to be given by expressing its co-ordinates  $x_\nu$  as functions of  $\lambda$ . Let the symbol  $\delta$  indicate the transition from a point of the required geodetic to the point corresponding to the same  $\lambda$  on a neighbouring line. Then for (20) we may substitute

$$\left. \begin{aligned} \int_{\lambda_1}^{\lambda_2} \delta w d\lambda &= 0 \\ w^2 &= g_{\mu\nu} \frac{dx_\mu}{d\lambda} \frac{dx_\nu}{d\lambda} \end{aligned} \right\} \quad . \quad . \quad . \quad (20a)$$

But since

$$\delta w = \frac{1}{w} \left\{ \frac{1}{2} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \frac{dx_\mu}{d\lambda} \frac{dx_\nu}{d\lambda} \delta x_\sigma + g_{\mu\nu} \frac{dx_\mu}{d\lambda} \delta \left( \frac{dx_\nu}{d\lambda} \right) \right\},$$

and

$$\delta \left( \frac{dx_\nu}{d\lambda} \right) = \frac{d}{d\lambda} (\delta x_\nu),$$

we obtain from (20a), after a partial integration,

$$\int_{\lambda_1}^{\lambda_2} \kappa_\sigma \delta x_\sigma d\lambda = 0,$$

where

$$\kappa_\sigma = \frac{d}{d\lambda} \left\{ \frac{g_{\mu\nu}}{w} \frac{dx_\mu}{d\lambda} \right\} - \frac{1}{2w} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \frac{dx_\mu}{d\lambda} \frac{dx_\nu}{d\lambda} \quad (20b)$$

Since the values of  $\delta x_\sigma$  are arbitrary, it follows from this that

$$\kappa_\sigma = 0 \quad (20c)$$

are the equations of the geodetic line.

If  $ds$  does not vanish along the geodetic line we may choose the "length of the arc"  $s$ , measured along the geodetic line, for the parameter  $\lambda$ . Then  $w = 1$ , and in place of (20c) we obtain

$$g_{\mu\nu} \frac{d^2 x_\mu}{ds^2} + \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \frac{dx_\sigma}{ds} \frac{dx_\mu}{ds} - \frac{1}{2} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0$$

or, by a mere change of notation,

$$g_{\alpha\sigma} \frac{d^2 x_\alpha}{ds^2} + [\mu\nu, \sigma] \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0 \quad (20d)$$

where, following Christoffel, we have written

$$[\mu\nu, \sigma] = \frac{1}{2} \left( \frac{\partial g_{\mu\sigma}}{\partial x_\nu} + \frac{\partial g_{\nu\sigma}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\sigma} \right) \quad (21)$$

Finally, if we multiply (20d) by  $g^{\sigma\tau}$  (outer multiplication with respect to  $\tau$ , inner with respect to  $\sigma$ ), we obtain the equations of the geodetic line in the form

$$\frac{d^2 x_\tau}{ds^2} + \{\mu\nu, \tau\} \frac{dx_\mu}{ds} \frac{dx_\nu}{ds} = 0 \quad (22)$$

where, following Christoffel, we have set

$$\{\mu\nu, \tau\} = g^{\tau\alpha} [\mu\nu, \alpha] \quad (23)$$



and since by (23) and (21)  $\{\mu\nu, \tau\}$  is symmetrical in  $\mu$  and  $\nu$ , it follows that the expression in brackets is symmetrical in  $\mu$  and  $\nu$ . Since a geodetic line can be drawn in any direction from a point of the continuum, and therefore  $dx_\mu/ds$  is a four-vector with the ratio of its components arbitrary, it follows from the results of § 7 that

$$A_{\mu\nu} = \frac{\partial^2 \phi}{\partial x_\mu \partial x_\nu} - \{\mu\nu, \tau\} \frac{\partial \phi}{\partial x_\tau} \quad (25)$$

is a covariant tensor of the second rank. We have therefore come to this result: from the covariant tensor of the first rank

$$A_\mu = \frac{\partial \phi}{\partial x_\mu}$$

we can, by differentiation, form a covariant tensor of the second rank

$$A_{\mu\nu} = \frac{\partial A_\mu}{\partial x_\nu} - \{\mu\nu, \tau\} A_\tau \quad (26)$$

We call the tensor  $A_{\mu\nu}$  the "extension" (covariant derivative) of the tensor  $A_\mu$ . In the first place we can readily show that the operation leads to a tensor, even if the vector  $A_\mu$  cannot be represented as a gradient. To see this, we first observe that

$$\psi \frac{\partial \phi}{\partial x_\mu}$$

is a covariant vector, if  $\psi$  and  $\phi$  are scalars. The sum of four such terms

$$S_\mu = \psi^{(1)} \frac{\partial \phi^{(1)}}{\partial x_\mu} + \dots + \psi^{(4)} \frac{\partial \phi^{(4)}}{\partial x_\mu},$$

is also a covariant vector, if  $\psi^{(1)}, \phi^{(1)}, \dots, \psi^{(4)}, \phi^{(4)}$  are scalars. But it is clear that any covariant vector can be represented in the form  $S_\mu$ . For, if  $A_\mu$  is a vector whose components are any given functions of the  $x_\nu$ , we have only to put (in terms of the selected system of co-ordinates)

$$\begin{aligned} \psi^{(1)} &= A_1, & \phi^{(1)} &= x_1, \\ \psi^{(2)} &= A_2, & \phi^{(2)} &= x_2, \\ \psi^{(3)} &= A_3, & \phi^{(3)} &= x_3, \\ \psi^{(4)} &= A_4, & \phi^{(4)} &= x_4, \end{aligned}$$

in order to ensure that  $S_\mu$  shall be equal to  $A_\mu$ .

Therefore, in order to demonstrate that  $A_{\mu\nu}$  is a tensor if *any* covariant vector is inserted on the right-hand side for  $A_{\mu}$ , we only need show that this is so for the vector  $S_{\mu}$ . But for this latter purpose it is sufficient, as a glance at the right-hand side of (26) teaches us, to furnish the proof for the case

$$A_{\mu} = \psi \frac{\partial \phi}{\partial x_{\mu}}.$$

Now the right-hand side of (25) multiplied by  $\psi$ ,

$$\psi \frac{\partial^2 \phi}{\partial x_{\mu} \partial x_{\nu}} - \{\mu\nu, \tau\} \psi \frac{\partial \phi}{\partial x_{\tau}}$$

is a tensor. Similarly

$$\frac{\partial \psi}{\partial x_{\mu}} \frac{\partial \phi}{\partial x_{\nu}}$$

being the outer product of two vectors, is a tensor. By addition, there follows the tensor character of

$$\frac{\partial}{\partial x_{\nu}} \left( \psi \frac{\partial \phi}{\partial x_{\mu}} \right) - \{\mu\nu, \tau\} \left( \psi \frac{\partial \phi}{\partial x_{\tau}} \right).$$

As a glance at (26) will show, this completes the demonstration for the vector

$$\psi \frac{\partial \phi}{\partial x_{\mu}}$$

and consequently, from what has already been proved, for any vector  $A_{\mu}$ .

By means of the extension of the vector, we may easily define the "extension" of a covariant tensor of any rank. This operation is a generalization of the extension of a vector. We restrict ourselves to the case of a tensor of the second rank, since this suffices to give a clear idea of the law of formation.

As has already been observed, any covariant tensor of the second rank can be represented \* as the sum of tensors of the

\* By outer multiplication of the vector with arbitrary components  $A_{11}$ ,  $A_{12}$ ,  $A_{13}$ ,  $A_{14}$  by the vector with components 1, 0, 0, 0, we produce a tensor with components

$A_{11}$	$A_{12}$	$A_{13}$	$A_{14}$
0	0	0	0
0	0	0	0
0	0	0	0

By the addition of four tensors of this type, we obtain the tensor  $A_{\mu\nu}$  with any assigned components.



type  $A_\mu B_\nu$ . It will therefore be sufficient to deduce the expression for the extension of a tensor of this special type. By (26) the expressions

$$\frac{\partial A_\mu}{\partial x_\sigma} - \{\sigma\mu, \tau\}A_\tau,$$

$$\frac{\partial B_\nu}{\partial x_\sigma} - \{\sigma\nu, \tau\}B_\tau,$$

are tensors. On outer multiplication of the first by  $B_\nu$ , and of the second by  $A_\mu$ , we obtain in each case a tensor of the third rank. By adding these, we have the tensor of the third rank

$$A_{\mu\nu\sigma} = \frac{\partial A_{\mu\nu}}{\partial x_\sigma} - \{\sigma\mu, \tau\}A_{\tau\nu} - \{\sigma\nu, \tau\}A_{\mu\tau}. \quad (27)$$

where we have put  $A_{\mu\nu} = A_\mu B_\nu$ . As the right-hand side of (27) is linear and homogeneous in the  $A_{\mu\nu}$  and their first derivatives, this law of formation leads to a tensor, not only in the case of a tensor of the type  $A_\mu B_\nu$ , but also in the case of a sum of such tensors, i.e. in the case of any covariant tensor of the second rank. We call  $A_{\mu\nu\sigma}$  the extension of the tensor  $A_{\mu\nu}$ .

It is clear that (26) and (24) concern only special cases of extension (the extension of the tensors of rank one and zero respectively).

In general, all special laws of formation of tensors are included in (27) in combination with the multiplication of tensors.

## § 11. Some Cases of Special Importance

*The Fundamental Tensor.*—We will first prove some lemmas which will be useful hereafter. By the rule for the differentiation of determinants

$$dg = g^{\mu\nu}g dg_{\mu\nu} = -g_{\mu\nu}g dg^{\mu\nu} \quad (28)$$

The last member is obtained from the last but one, if we bear in mind that  $g_{\mu\nu}g^{\mu'\nu} = \delta_\mu^{\mu'}$ , so that  $g_{\mu\nu}g^{\mu\nu} = 4$ , and consequently

$$g_{\mu\nu}dg^{\mu\nu} + g^{\mu\nu}dg_{\mu\nu} = 0.$$

From (28), it follows that

$$\frac{1}{\sqrt{-g}} \frac{\partial \sqrt{-g}}{\partial x_\sigma} = \frac{1}{2} \frac{\partial \log(-g)}{\partial x_\sigma} = \frac{1}{2} g^{\mu\nu} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} = \frac{1}{2} g^{\mu\nu} \frac{\partial g^{\mu\nu}}{\partial x_\sigma}. \quad (29)$$

Further, from  $g_{\mu\sigma} g^{\nu\sigma} = \delta_\mu^\nu$ , it follows on differentiation that

$$\left. \begin{aligned} g_{\mu\sigma} dg^{\nu\sigma} &= -g^{\nu\sigma} dg_{\mu\sigma} \\ g_{\mu\sigma} \frac{\partial g^{\nu\sigma}}{\partial x_\lambda} &= -g^{\nu\sigma} \frac{\partial g_{\mu\sigma}}{\partial x_\lambda} \end{aligned} \right\} \quad . \quad . \quad . \quad (30)$$

From these, by mixed multiplication by  $g^{\sigma\tau}$  and  $g_{\nu\lambda}$  respectively, and a change of notation for the indices, we have

$$\left. \begin{aligned} dg^{\mu\nu} &= -g^{\mu\alpha} g^{\nu\beta} dg_{\alpha\beta} \\ \frac{\partial g^{\mu\nu}}{\partial x_\sigma} &= -g^{\mu\alpha} g^{\nu\beta} \frac{\partial g_{\alpha\beta}}{\partial x_\sigma} \end{aligned} \right\} \quad . \quad . \quad . \quad (31)$$

and

$$\left. \begin{aligned} dg_{\mu\nu} &= -g_{\mu\alpha} g_{\nu\beta} dg^{\alpha\beta} \\ \frac{\partial g_{\mu\nu}}{\partial x_\sigma} &= -g_{\mu\alpha} g_{\nu\beta} \frac{\partial g^{\alpha\beta}}{\partial x_\sigma} \end{aligned} \right\} \quad . \quad . \quad . \quad (32)$$

The relation (31) admits of a transformation, of which we also have frequently to make use. From (21)

$$\frac{\partial g_{\alpha\beta}}{\partial x_\sigma} = [a\sigma, \beta] + [\beta\sigma, \alpha] \quad . \quad . \quad . \quad (33)$$

Inserting this in the second formula of (31), we obtain, in view of (23)

$$\frac{\partial g^{\mu\nu}}{\partial x_\sigma} = -g^{\mu\tau} \{\tau\sigma, \nu\} - g^{\nu\tau} \{\tau\sigma, \mu\} \quad . \quad . \quad . \quad (34)$$

Substituting the right-hand side of (34) in (29), we have

$$\frac{1}{\sqrt{-g}} \frac{\partial \sqrt{-g}}{\partial x_\sigma} = \{\mu\sigma, \mu\} \quad . \quad . \quad . \quad (29a)$$

*The "Divergence" of a Contravariant Vector.*—If we take the inner product of (26) by the contravariant fundamental tensor  $g^{\mu\nu}$ , the right-hand side, after a transformation of the first term, assumes the form

$$\frac{\partial}{\partial x_\nu} (g^{\mu\nu} A_\mu) - A_\mu \frac{\partial g^{\mu\nu}}{\partial x_\nu} - \frac{1}{2} g^{\tau\alpha} \left( \frac{\partial g_{\mu\alpha}}{\partial x_\nu} + \frac{\partial g_{\nu\alpha}}{\partial x_\mu} - \frac{\partial g_{\mu\nu}}{\partial x_\alpha} \right) g^{\mu\nu} A_\tau.$$

In accordance with (31) and (29), the last term of this expression may be written

$$\frac{1}{2} \frac{\partial g^{\tau\nu}}{\partial x_\nu} A_\tau + \frac{1}{2} \frac{\partial g^{\tau\mu}}{\partial x_\mu} A_\tau + \frac{1}{\sqrt{-g}} \frac{\partial \sqrt{-g}}{\partial x_\alpha} g^{\mu\nu} A_\tau.$$

As the symbols of the indices of summation are immaterial, the first two terms of this expression cancel the second of the one above. If we then write  $g^{\mu\nu} A_\mu = A^\nu$ , so that  $A^\nu$  like  $A_\mu$  is an arbitrary vector, we finally obtain

$$\Phi = \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x_\nu} (\sqrt{-g} A^\nu). \quad (35)$$

This scalar is the *divergence* of the contravariant vector  $A^\nu$ .

*The "Curl" of a Covariant Vector.*—The second term in (26) is symmetrical in the indices  $\mu$  and  $\nu$ . Therefore  $A_{\mu\nu} - A_{\nu\mu}$  is a particularly simply constructed antisymmetrical tensor. We obtain

$$B_{\mu\nu} = \frac{\partial A_\mu}{\partial x_\nu} - \frac{\partial A_\nu}{\partial x_\mu} \quad (36)$$

*Antisymmetrical Extension of a Six-vector.*—Applying (27) to an antisymmetrical tensor of the second rank  $A_{\mu\nu}$ , forming in addition the two equations which arise through cyclic permutations of the indices, and adding these three equations, we obtain the tensor of the third rank

$$B_{\mu\nu\sigma} = A_{\mu\nu\sigma} + A_{\nu\sigma\mu} + A_{\sigma\mu\nu} = \frac{\partial A_{\mu\nu}}{\partial x_\sigma} + \frac{\partial A_{\nu\sigma}}{\partial x_\mu} + \frac{\partial A_{\sigma\mu}}{\partial x_\nu} \quad (37)$$

which it is easy to prove is antisymmetrical.

*The Divergence of a Six-vector.*—Taking the mixed product of (27) by  $g^{\mu\alpha} g^{\nu\beta}$ , we also obtain a tensor. The first term on the right-hand side of (27) may be written in the form

$$\frac{\partial}{\partial x_\sigma} (g^{\mu\alpha} g^{\nu\beta} A_{\mu\nu}) - g^{\mu\alpha} \frac{\partial g^{\nu\beta}}{\partial x_\sigma} A_{\mu\nu} - g^{\nu\beta} \frac{\partial g^{\mu\alpha}}{\partial x_\sigma} A_{\mu\nu}.$$

If we write  $A_\sigma^{\alpha\beta}$  for  $g^{\mu\alpha} g^{\nu\beta} A_{\mu\nu\sigma}$  and  $A^{\alpha\beta}$  for  $g^{\mu\alpha} g^{\nu\beta} A_{\mu\nu}$ , and in the transformed first term replace

$$\frac{\partial g^{\nu\beta}}{\partial x_\sigma} \text{ and } \frac{\partial g^{\mu\alpha}}{\partial x_\sigma}$$

by their values as given by (34), there results from the right-hand side of (27) an expression consisting of seven terms, of which four cancel, and there remains

$$A^{\alpha\beta}_{\sigma} = \frac{\partial A^{\alpha\beta}}{\partial x_{\sigma}} + \{\sigma\gamma, \alpha\}A^{\gamma\beta} + \{\sigma\gamma, \beta\}A^{\alpha\gamma}. \quad (38)$$

This is the expression for the extension of a contravariant tensor of the second rank, and corresponding expressions for the extension of contravariant tensors of higher and lower rank may also be formed.

We note that in an analogous way we may also form the extension of a mixed tensor :—

$$A^{\alpha}_{\mu\sigma} = \frac{\partial A^{\alpha}_{\mu}}{\partial x_{\sigma}} - \{\sigma\mu, \tau\}A^{\alpha}_{\tau} + \{\sigma\tau, \alpha\}A^{\tau}_{\mu}. \quad (39)$$

On contracting (38) with respect to the indices  $\beta$  and  $\sigma$  (inner multiplication by  $\delta^{\sigma}_{\beta}$ ), we obtain the vector

$$A^{\alpha} = \frac{\partial A^{\alpha\beta}}{\partial x_{\beta}} + \{\beta\gamma, \beta\}A^{\alpha\gamma} + \{\beta\gamma, \alpha\}A^{\gamma\beta}.$$

On account of the symmetry of  $\{\beta\gamma, \alpha\}$  with respect to the indices  $\beta$  and  $\gamma$ , the third term on the right-hand side vanishes, if  $A^{\alpha\beta}$  is, as we will assume, an antisymmetrical tensor. The second term allows itself to be transformed in accordance with (29a). Thus we obtain

$$A^{\alpha} = \frac{1}{\sqrt{-g}} \frac{\partial(\sqrt{-g}A^{\alpha\beta})}{\partial x_{\beta}}. \quad (40)$$

This is the expression for the divergence of a contravariant six-vector.

*The Divergence of a Mixed Tensor of the Second Rank.*—Contracting (39) with respect to the indices  $\alpha$  and  $\sigma$ , and taking (29a) into consideration, we obtain

$$\sqrt{-g}A_{\mu} = \frac{\partial(\sqrt{-g}A^{\sigma}_{\mu})}{\partial x_{\sigma}} - \{\sigma\mu, \tau\}\sqrt{-g}A^{\sigma}_{\tau}. \quad (41)$$

If we introduce the contravariant tensor  $A^{\rho\sigma} = g^{\rho\tau}A^{\sigma}_{\tau}$  in the last term, it assumes the form

$$- [\sigma\mu, \rho]\sqrt{-g}A^{\rho\sigma}.$$

If, further, the tensor  $A^{\rho\sigma}$  is symmetrical, this reduces to

$$-\frac{1}{2}\sqrt{-g}\frac{\partial g^{\rho\sigma}}{\partial x_\mu}A^{\rho\sigma}.$$

Had we introduced, instead of  $A^{\rho\sigma}$ , the covariant tensor  $A_{\rho\sigma} = g_{\rho\alpha}g_{\sigma\beta}A^{\alpha\beta}$ , which is also symmetrical, the last term, by virtue of (31), would assume the form

$$\frac{1}{2}\sqrt{-g}\frac{\partial g^{\rho\sigma}}{\partial x_\mu}A_{\rho\sigma}.$$

In the case of symmetry in question, (41) may therefore be replaced by the two forms

$$\sqrt{-g}A_\mu = \frac{\partial(\sqrt{-g}A_\mu^\sigma)}{\partial x_\sigma} - \frac{1}{2}\frac{\partial g^{\rho\sigma}}{\partial x_\mu}\sqrt{-g}A^{\rho\sigma}. \quad (41a)$$

$$\sqrt{-g}A_\mu = \frac{\partial(\sqrt{-g}A_\mu^\sigma)}{\partial x_\sigma} + \frac{1}{2}\frac{\partial g^{\rho\sigma}}{\partial x_\mu}\sqrt{-g}A_{\rho\sigma}. \quad (41b)$$

which we have to employ later on.

## § 12. The Riemann-Christoffel Tensor

We now seek the tensor which can be obtained from the fundamental tensor *alone*, by differentiation. At first sight the solution seems obvious. We place the fundamental tensor of the  $g_{\mu\nu}$  in (27) instead of any given tensor  $A_{\mu\nu}$ , and thus have a new tensor, namely, the extension of the fundamental tensor. But we easily convince ourselves that this extension vanishes identically. We reach our goal, however, in the following way. In (27) place

$$A_{\mu\nu} = \frac{\partial A_\mu}{\partial x_\nu} - \{\mu\nu, \rho\}A_\rho,$$

i.e. the extension of the four-vector  $A_\mu$ . Then (with a somewhat different naming of the indices) we get the tensor of the third rank

$$\begin{aligned} A_{\mu\sigma\tau} = & \frac{\partial^2 A_\mu}{\partial x_\sigma \partial x_\tau} - \{\mu\sigma, \rho\}\frac{\partial A_\rho}{\partial x_\tau} - \{\mu\tau, \rho\}\frac{\partial A_\rho}{\partial x_\sigma} - \{\sigma\tau, \rho\}\frac{\partial A_\mu}{\partial x_\rho} \\ & + \left[ -\frac{\partial}{\partial x_\tau}\{\mu\sigma, \rho\} + \{\mu\tau, \alpha\}\{\alpha\sigma, \rho\} + \{\sigma\tau, \alpha\}\{\alpha\mu, \rho\} \right] A_\rho. \end{aligned}$$

This expression suggests forming the tensor  $A_{\mu\sigma\tau} - A_{\mu\tau\sigma}$ . For, if we do so, the following terms of the expression for  $A_{\mu\sigma\tau}$  cancel those of  $A_{\mu\tau\sigma}$ , the first, the fourth, and the member corresponding to the last term in square brackets; because all these are symmetrical in  $\sigma$  and  $\tau$ . The same holds good for the sum of the second and third terms. Thus we obtain

$$A_{\mu\sigma\tau} - A_{\mu\tau\sigma} = B_{\mu\sigma\tau}^{\rho} A_{\rho} \quad . \quad . \quad . \quad (42)$$

where

$$B_{\mu\sigma\tau}^{\rho} = - \frac{\partial}{\partial x_{\tau}} \{\mu\sigma, \rho\} + \frac{\partial}{\partial x_{\sigma}} \{\mu\tau, \rho\} - \{\mu\sigma, \alpha\} \{\alpha\tau, \rho\} \\ + \{\mu\tau, \alpha\} \{\alpha\sigma, \rho\} \quad (43)$$

The essential feature of the result is that on the right side of (42) the  $A_{\rho}$  occur alone, without their derivatives. From the tensor character of  $A_{\mu\sigma\tau} - A_{\mu\tau\sigma}$  in conjunction with the fact that  $A_{\rho}$  is an arbitrary vector, it follows, by reason of § 7, that  $B_{\mu\sigma\tau}^{\rho}$  is a tensor (the Riemann-Christoffel tensor).

The mathematical importance of this tensor is as follows: If the continuum is of such a nature that there is a co-ordinate system with reference to which the  $g_{\mu\nu}$  are constants, then all the  $B_{\mu\sigma\tau}^{\rho}$  vanish. If we choose any new system of co-ordinates in place of the original ones, the  $g_{\mu\nu}$  referred thereto will not be constants, but in consequence of its tensor nature, the transformed components of  $B_{\mu\sigma\tau}^{\rho}$  will still vanish in the new system. Thus the vanishing of the Riemann tensor is a necessary condition that, by an appropriate choice of the system of reference, the  $g_{\mu\nu}$  may be constants. In our problem this corresponds to the case in which,\* with a suitable choice of the system of reference, the special theory of relativity holds good for a *finite* region of the continuum.

Contracting (43) with respect to the indices  $\tau$  and  $\rho$  we obtain the covariant tensor of second rank

\* The mathematicians have proved that this is also a *sufficient* condition,

$$\left. \begin{aligned}
 G_{\mu\nu} &= B_{\mu\nu\rho}^{\rho} = R_{\mu\nu} + S_{\mu\nu} \\
 \text{where} \quad R_{\mu\nu} &= -\frac{\partial}{\partial x_a} \{ \mu\nu, \alpha \} + \{ \mu\alpha, \beta \} \{ \nu\beta, \alpha \} \\
 S_{\mu\nu} &= \frac{\partial^2 \log \sqrt{-g}}{\partial x_\mu \partial x_\nu} - \{ \mu\nu, \alpha \} \frac{\partial \log \sqrt{-g}}{\partial x_\alpha}
 \end{aligned} \right\} \quad (44)$$

*Note on the Choice of Co-ordinates.*—It has already been observed in § 8, in connexion with equation (18a), that the choice of co-ordinates may with advantage be made so that  $\sqrt{-g} = 1$ . A glance at the equations obtained in the last two sections shows that by such a choice the laws of formation of tensors undergo an important simplification. This applies particularly to  $G_{\mu\nu}$ , the tensor just developed, which plays a fundamental part in the theory to be set forth. For this specialization of the choice of co-ordinates brings about the vanishing of  $S_{\mu\nu}$ , so that the tensor  $G_{\mu\nu}$  reduces to  $R_{\mu\nu}$ .

On this account I shall hereafter give all relations in the simplified form which this specialization of the choice of co-ordinates brings with it. It will then be an easy matter to revert to the *generally* covariant equations, if this seems desirable in a special case.

## C. THEORY OF THE GRAVITATIONAL FIELD

### § 13. Equations of Motion of a Material Point in the Gravitational Field. Expression for the Field-components of Gravitation

A freely movable body not subjected to external forces moves, according to the special theory of relativity, in a straight line and uniformly. This is also the case, according to the general theory of relativity, for a part of four-dimensional space in which the system of co-ordinates  $K_0$ , may be, and is, so chosen that they have the special constant values given in (4).

If we consider precisely this movement from any chosen system of co-ordinates  $K_1$ , the body, observed from  $K_1$ , moves, according to the considerations in § 2, in a gravitational field. The law of motion with respect to  $K_1$  results without diffi-

culty from the following consideration. With respect to  $K_0$  the law of motion corresponds to a four-dimensional straight line, i.e. to a geodetic line. Now since the geodetic line is defined independently of the system of reference, its equations will also be the equation of motion of the material point with respect to  $K_1$ . If we set

$$\Gamma_{\mu\nu}^{\tau} = -\{\mu\nu, \tau\} \quad . \quad . \quad . \quad (45)$$

the equation of the motion of the point with respect to  $K_1$ , becomes

$$\frac{d^2 x_{\tau}}{ds^2} = \Gamma_{\mu\nu}^{\tau} \frac{dx_{\mu}}{ds} \frac{dx_{\nu}}{ds} \quad . \quad . \quad . \quad (46)$$

We now make the assumption, which readily suggests itself, that this covariant system of equations also defines the motion of the point in the gravitational field in the case when there is no system of reference  $K_0$ , with respect to which the special theory of relativity holds good in a finite region. We have all the more justification for this assumption as (46) contains only *first* derivatives of the  $g_{\mu\nu}$ , between which even in the special case of the existence of  $K_0$ , no relations subsist.\*

If the  $\Gamma_{\mu\nu}^{\tau}$  vanish, then the point moves uniformly in a straight line. These quantities therefore condition the deviation of the motion from uniformity. They are the components of the gravitational field.

#### § 14. The Field Equations of Gravitation in the Absence of Matter

We make a distinction hereafter between "gravitational field" and "matter" in this way, that we denote everything but the gravitational field as "matter." Our use of the word therefore includes not only matter in the ordinary sense, but the electromagnetic field as well.

Our next task is to find the field equations of gravitation in the absence of matter. Here we again apply the method

\* It is only between the second (and first) derivatives that, by § 12, the relations  $B_{\mu\sigma\tau}^{\rho} = 0$  subsist.



employed in the preceding paragraph in formulating the equations of motion of the material point. A special case in which the required equations must in any case be satisfied is that of the special theory of relativity, in which the  $g_{\mu\nu}$  have certain constant values. Let this be the case in a certain finite space in relation to a definite system of co-ordinates  $K_0$ . Relatively to this system all the components of the Riemann tensor  $B_{\mu\sigma\tau}^{\rho}$ , defined in (43), vanish. For the space under consideration they then vanish, also in any other system of co-ordinates.

Thus the required equations of the matter-free gravitational field must in any case be satisfied if all  $B_{\mu\sigma\tau}^{\rho}$  vanish. But this condition goes too far. For it is clear that, e.g., the gravitational field generated by a material point in its environment certainly cannot be "transformed away" by any choice of the system of co-ordinates, i.e. it cannot be transformed to the case of constant  $g_{\mu\nu}$ .

This prompts us to require for the matter-free gravitational field that the symmetrical tensor  $G_{\mu\nu}$ , derived from the tensor  $B_{\mu\nu\tau}^{\rho}$ , shall vanish. Thus we obtain ten equations for the ten quantities  $g_{\mu\nu}$ , which are satisfied in the special case of the vanishing of all  $B_{\mu\nu\tau}^{\rho}$ . With the choice which we have made of a system of co-ordinates, and taking (44) into consideration, the equations for the matter-free field are

$$\left. \begin{aligned} \frac{\partial I_{\mu\nu}^{\alpha}}{\partial x_{\alpha}} + I_{\mu\beta}^{\alpha} I_{\nu\alpha}^{\beta} &= 0 \\ \sqrt{-g} &= 1 \end{aligned} \right\} \quad (47)$$

It must be pointed out that there is only a minimum of arbitrariness in the choice of these equations. For besides  $G_{\mu\nu}$  there is no tensor of second rank which is formed from the  $g_{\mu\nu}$  and its derivatives, contains no derivations higher than second, and is linear in these derivatives.\*

These equations, which proceed, by the method of pure

\* Properly speaking, this can be affirmed only of the tensor

$$G_{\mu\nu} + \lambda g_{\mu\nu} g^{\alpha\beta} G_{\alpha\beta},$$

where  $\lambda$  is a constant. If, however, we set this tensor = 0, we come back again to the equations  $G_{\mu\nu} = 0$ .

mathematics, from the requirement of the general theory of relativity, give us, in combination with the equations of motion (46), to a first approximation Newton's law of attraction, and to a second approximation the explanation of the motion of the perihelion of the planet Mercury discovered by Leverrier (as it remains after corrections for perturbation have been made). These facts must, in my opinion, be taken as a convincing proof of the correctness of the theory.

### § 15. The Hamiltonian Function for the Gravitational Field. Laws of Momentum and Energy

To show that the field equations correspond to the laws of momentum and energy, it is most convenient to write them in the following Hamiltonian form:—

$$\left. \begin{aligned} \delta \int H d\tau &= 0 \\ H &= g^{\mu\nu} \Gamma_{\mu\beta}^{\alpha} \Gamma_{\nu\alpha}^{\beta} \\ \sqrt{-g} &= 1 \end{aligned} \right\} \quad . \quad . \quad . \quad (47a)$$

where, on the boundary of the finite four-dimensional region of integration which we have in view, the variations vanish.

We first have to show that the form (47a) is equivalent to the equations (47). For this purpose we regard  $H$  as a function of the  $g^{\mu\nu}$  and the  $g^{\mu\nu}_{,\sigma}$  ( $= \partial g^{\mu\nu} / \partial x_{\sigma}$ ).

Then in the first place

$$\begin{aligned} \delta H &= \Gamma_{\mu\beta}^{\alpha} \Gamma_{\nu\alpha}^{\beta} \delta g^{\mu\nu} + 2g^{\mu\nu} \Gamma_{\mu\beta}^{\alpha} \delta \Gamma_{\nu\alpha}^{\beta} \\ &= - \Gamma_{\mu\beta}^{\alpha} \Gamma_{\nu\alpha}^{\beta} \delta g^{\mu\nu} + 2\Gamma_{\mu\beta}^{\alpha} \delta(g^{\mu\nu} \Gamma_{\nu\alpha}^{\beta}). \end{aligned}$$

But

$$\delta(g^{\mu\nu} \Gamma_{\nu\alpha}^{\beta}) = -\frac{1}{2} \delta \left[ g^{\mu\nu} g^{\beta\lambda} \left( \frac{\partial g_{\nu\lambda}}{\partial x_{\alpha}} + \frac{\partial g_{\alpha\lambda}}{\partial x_{\nu}} - \frac{\partial g_{\alpha\nu}}{\partial x_{\lambda}} \right) \right].$$

The terms arising from the last two terms in round brackets are of different sign, and result from each other (since the denomination of the summation indices is immaterial) through interchange of the indices  $\mu$  and  $\beta$ . They cancel each other in the expression for  $\delta H$ , because they are multiplied by the

quantity  $I_{\mu\beta}^{\alpha}$ , which is symmetrical with respect to the indices  $\mu$  and  $\beta$ . Thus there remains only the first term in round brackets to be considered, so that, taking (31) into account, we obtain

$$\delta H = - I_{\mu\beta}^{\alpha} I_{\nu\alpha}^{\beta} \delta g^{\mu\nu} + I_{\mu\beta}^{\alpha} \delta g_{\alpha}^{\mu\beta}.$$

Thus

$$\left. \begin{aligned} \frac{\partial H}{\partial g^{\mu\nu}} &= - I_{\mu\beta}^{\alpha} I_{\nu\alpha}^{\beta} \\ \frac{\partial H}{\partial g_{\sigma}^{\mu\nu}} &= I_{\mu\nu}^{\sigma} \end{aligned} \right\} \quad . \quad . \quad . \quad (48)$$

Carrying out the variation in (47a), we get in the first place

$$\frac{\partial}{\partial x_{\alpha}} \left( \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} \right) - \frac{\partial H}{\partial g^{\mu\nu}} = 0, \quad . \quad . \quad . \quad (47b)$$

which, on account of (48), agrees with (47), as was to be proved.

If we multiply (47b) by  $g_{\sigma}^{\mu\nu}$ , then because

$$\frac{\partial g_{\sigma}^{\mu\nu}}{\partial x_{\alpha}} = \frac{\partial g_{\alpha}^{\mu\nu}}{\partial x_{\sigma}}$$

and, consequently,

$$g_{\sigma}^{\mu\nu} \frac{\partial}{\partial x_{\alpha}} \left( \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} \right) = \frac{\partial}{\partial x_{\alpha}} \left( g_{\sigma}^{\mu\nu} \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} \right) - \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} \frac{\partial g_{\alpha}^{\mu\nu}}{\partial x_{\sigma}},$$

we obtain the equation

$$\frac{\partial}{\partial x_{\alpha}} \left( g_{\sigma}^{\mu\nu} \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} \right) - \frac{\partial H}{\partial x_{\sigma}} = 0$$

or \*

$$\left. \begin{aligned} \frac{\partial t_{\sigma}^{\alpha}}{\partial x_{\alpha}} &= 0 \\ - 2\kappa t_{\sigma}^{\alpha} &= g_{\sigma}^{\mu\nu} \frac{\partial H}{\partial g_{\alpha}^{\mu\nu}} - \delta_{\sigma}^{\alpha} H \end{aligned} \right\} \quad . \quad . \quad . \quad (49)$$

where, on account of (48), the second equation of (47), and (34)

$$\kappa t_{\sigma}^{\alpha} = \frac{1}{2} \delta_{\sigma}^{\alpha} g^{\mu\nu} I_{\mu\beta}^{\lambda} I_{\nu\lambda}^{\beta} - g^{\mu\nu} I_{\mu\beta}^{\alpha} I_{\nu\sigma}^{\beta} \quad . \quad . \quad (50)$$

\* The reason for the introduction of the factor  $- 2\kappa$  will be apparent later.

It is to be noticed that  $t_\sigma^\alpha$  is not a tensor; on the other hand (49) applies to all systems of co-ordinates for which  $\sqrt{-g} = 1$ . This equation expresses the law of conservation of momentum and of energy for the gravitational field. Actually the integration of this equation over a three-dimensional volume  $V$  yields the four equations

$$\frac{d}{dx_\alpha} \int t_\sigma^\alpha dV = \int (lt_\sigma^1 + mt_\sigma^2 + nt_\sigma^3) dS. \quad (49a)$$

where  $l, m, n$  denote the direction-cosines of direction of the inward drawn normal at the element  $dS$  of the bounding surface (in the sense of Euclidean geometry). We recognize in this the expression of the laws of conservation in their usual form. The quantities  $t_\sigma^\alpha$  we call the "energy components" of the gravitational field.

I will now give equations (47) in a third form, which is particularly useful for a vivid grasp of our subject. By multiplication of the field equations (47) by  $g^{\nu\sigma}$  these are obtained in the "mixed" form. Note that

$$g^{\nu\sigma} \frac{\partial \Gamma_{\mu\nu}^\alpha}{\partial x_\alpha} = \frac{\partial}{\partial x_\alpha} (g^{\nu\sigma} \Gamma_{\mu\nu}^\alpha) - \frac{\partial g^{\nu\sigma}}{\partial x_\alpha} \Gamma_{\mu\nu}^\alpha,$$

which quantity, by reason of (34), is equal to

$$\frac{\partial}{\partial x_\alpha} (g^{\nu\sigma} \Gamma_{\mu\nu}^\alpha) - g^{\nu\beta} \Gamma_{\alpha\beta}^\sigma \Gamma_{\mu\nu}^\alpha - g^{\sigma\beta} \Gamma_{\beta\alpha}^\nu \Gamma_{\mu\nu}^\alpha,$$

or (with different symbols for the summation indices)

$$\frac{\partial}{\partial x_\alpha} (g^{\sigma\beta} \Gamma_{\mu\beta}^\alpha) - g^{\gamma\delta} \Gamma_{\gamma\beta}^\sigma \Gamma_{\delta\mu}^\beta - g^{\nu\sigma} \Gamma_{\mu\beta}^\alpha \Gamma_{\nu\alpha}^\beta.$$

The third term of this expression cancels with the one arising from the second term of the field equations (47); using relation (50), the second term may be written

$$\kappa(t_\mu^\sigma - \frac{1}{2}\delta_\mu^\sigma t),$$

where  $t = t_\alpha^\alpha$ . Thus instead of equations (47) we obtain

$$\left. \begin{aligned} \frac{\partial}{\partial x_\alpha} (g^{\sigma\beta} \Gamma_{\mu\beta}^\alpha) &= - \kappa(t_\mu^\sigma - \frac{1}{2}\delta_\mu^\sigma t) \\ \sqrt{-g} &= 1 \end{aligned} \right\} \quad (51)$$

### § 16. The General Form of the Field Equations of Gravitation

The field equations for matter-free space formulated in § 15 are to be compared with the field equation

$$\nabla^2\phi = 0$$

of Newton's theory. We require the equation corresponding to Poisson's equation

$$\nabla^2\phi = 4\pi\kappa\rho,$$

where  $\rho$  denotes the density of matter.

The special theory of relativity has led to the conclusion that inert mass is nothing more or less than energy, which finds its complete mathematical expression in a symmetrical tensor of second rank, the energy-tensor. Thus in the general theory of relativity we must introduce a corresponding energy-tensor of matter  $T_{\sigma}^{\alpha}$ , which, like the energy-components  $t_{\sigma}$  [equations (49) and (50)] of the gravitational field, will have mixed character, but will pertain to a symmetrical covariant tensor.\*

The system of equation (51) shows how this energy-tensor (corresponding to the density  $\rho$  in Poisson's equation) is to be introduced into the field equations of gravitation. For if we consider a complete system (e.g. the solar system), the total mass of the system, and therefore its total gravitating action as well, will depend on the total energy of the system, and therefore on the ponderable energy together with the gravitational energy. This will allow itself to be expressed by introducing into (51), in place of the energy-components of the gravitational field alone, the sums  $t_{\mu}^{\sigma} + T_{\mu}^{\sigma}$  of the energy-components of matter and of gravitational field. Thus instead of (51) we obtain the tensor equation

$$\left. \begin{aligned} \frac{\partial}{\partial x_{\alpha}}(g^{\sigma\beta}T_{\mu\beta}^{\alpha}) &= -\kappa[(t_{\mu}^{\sigma} + T_{\mu}^{\sigma}) - \frac{1}{2}\delta_{\mu}^{\sigma}(t + T)], \\ \sqrt{-g} &= 1 \end{aligned} \right\} \quad (52)$$

where we have set  $T = T_{\mu}^{\mu}$  (Laue's scalar). These are the

\*  $g_{\alpha\tau}T_{\sigma}^{\alpha} = T_{\sigma\tau}$  and  $g^{\sigma\beta}T_{\sigma}^{\alpha} = T^{\alpha\beta}$  are to be symmetrical tensors.

required general field equations of gravitation in mixed form. Working back from these, we have in place of (47)

$$\left. \begin{aligned} \frac{\partial}{\partial x_a} \Gamma_{\mu\nu}^a + \Gamma_{\mu\beta}^a \Gamma_{\nu a}^\beta &= -\kappa (T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T), \\ \sqrt{-g} &= 1 \end{aligned} \right\} \quad (53)$$

It must be admitted that this introduction of the energy-tensor of matter is not justified by the relativity postulate alone. For this reason we have here deduced it from the requirement that the energy of the gravitational field shall act gravitatively in the same way as any other kind of energy. But the strongest reason for the choice of these equations lies in their consequence, that the equations of conservation of momentum and energy, corresponding exactly to equations (49) and (49a), hold good for the components of the total energy. This will be shown in § 17.

### § 17. The Laws of Conservation in the General Case

Equation (52) may readily be transformed so that the second term on the right-hand side vanishes. Contract (52) with respect to the indices  $\mu$  and  $\sigma$ , and after multiplying the resulting equation by  $\frac{1}{2} \delta_\mu^\sigma$ , subtract it from equation (52). This gives

$$\frac{\partial}{\partial x_a} (g^{\sigma\beta} \Gamma_{\mu\beta}^a - \frac{1}{2} \delta_\mu^\sigma g^{\lambda\beta} \Gamma_{\lambda\beta}^a) = -\kappa (t_\mu^\sigma + T_\mu^\sigma). \quad (52a)$$

On this equation we perform the operation  $\partial/\partial x_\sigma$ . We have

$$\frac{\partial^2}{\partial x_a \partial x_\sigma} (g^\sigma \Gamma_{\beta\mu}^a) = -\frac{1}{2} \frac{\partial^2}{\partial x_a \partial x_\sigma} \left[ g^{\sigma\beta} g^{a\lambda} \left( \frac{\partial g_{\mu\lambda}}{\partial x_\beta} + \frac{\partial g_{\beta\lambda}}{\partial x_\mu} - \frac{\partial g_{\mu\beta}}{\partial x_\lambda} \right) \right].$$

The first and third terms of the round brackets yield contributions which cancel one another, as may be seen by interchanging, in the contribution of the third term, the summation indices  $\alpha$  and  $\sigma$  on the one hand, and  $\beta$  and  $\lambda$  on the other. The second term may be re-modelled by (31), so that we have

$$\frac{\partial^2}{\partial x_a \partial x_\sigma} (g^{\sigma\beta} \Gamma_{\mu\beta}^a) = \frac{1}{2} \frac{\partial^3 g^{a\beta}}{\partial x_a \partial x_\sigma \partial x_\mu} \quad (54)$$

The second term on the left-hand side of (52a) yields in the

first place

$$- \frac{1}{2} \frac{\partial^2}{\partial x_a \partial x_\mu} (g^{\lambda\beta} \Gamma_{\lambda\beta}^\alpha)$$

or

$$\frac{1}{2} \frac{\partial^2}{\partial x_a \partial x_\mu} \left[ g^{\lambda\beta} g^{\alpha\delta} \left( \frac{\partial g_{\delta\lambda}}{\partial x_\beta} + \frac{\partial g_{\delta\beta}}{\partial x_\lambda} - \frac{\partial g_{\lambda\beta}}{\partial x_\delta} \right) \right].$$

With the choice of co-ordinates which we have made, the term deriving from the last term in round brackets disappears by reason of (29). The other two may be combined, and together, by (31), they give

$$- \frac{1}{2} \frac{\partial^3 g^{\alpha\beta}}{\partial x_a \partial x_\beta \partial x_\mu},$$

so that in consideration of (54), we have the identity

$$\frac{\partial^2}{\partial x_a \partial x_\sigma} \left( g^{\rho\beta} \Gamma_{\mu\beta}^\sigma - \frac{1}{2} \delta_\mu^\sigma g^{\lambda\beta} \Gamma_{\lambda\beta}^\sigma \right) \equiv 0 \quad . \quad . \quad (55)$$

From (55) and (52a), it follows that

$$\frac{\partial(t_\mu^\sigma + T_\mu^\sigma)}{\partial x_\sigma} = 0. \quad . \quad . \quad (56)$$

Thus it results from our field equations of gravitation that the laws of conservation of momentum and energy are satisfied. This may be seen most easily from the consideration which leads to equation (49a); except that here, instead of the energy components  $t^\sigma$  of the gravitational field, we have to introduce the totality of the energy components of matter and gravitational field.

### § 18. The Laws of Momentum and Energy for Matter, as a Consequence of the Field Equations

Multiplying (53) by  $\partial g^{\mu\nu} / \partial x_\sigma$ , we obtain, by the method adopted in § 15, in view of the vanishing of

$$g_{\mu\nu} \frac{\partial g^{\mu\nu}}{\partial x_\sigma},$$

the equation

$$\frac{\partial t_\sigma^\alpha}{\partial x_\alpha} + \frac{1}{2} \frac{\partial g^{\mu\nu}}{\partial x_\sigma} T_{\mu\nu} = 0,$$

or, in view of (56),

$$\frac{\partial T_{\sigma}^{\alpha}}{\partial x_{\alpha}} + \frac{1}{2} \frac{\partial g^{\mu\nu}}{\partial x_{\sigma}} T_{\mu\nu} = 0 \quad . \quad . \quad . \quad (57)$$

Comparison with (41b) shows that with the choice of system of co-ordinates which we have made, this equation predicates nothing more or less than the vanishing of divergence of the material energy-tensor. Physically, the occurrence of the second term on the left-hand side shows that laws of conservation of momentum and energy do not apply in the strict sense for matter alone, or else that they apply only when the  $g^{\mu\nu}$  are constant, i.e. when the field intensities of gravitation vanish. This second term is an expression for momentum, and for energy, as transferred per unit of volume and time from the gravitational field to matter. This is brought out still more clearly by re-writing (57) in the sense of (41) as

$$\frac{\partial T_{\sigma}^{\alpha}}{\partial x_{\alpha}} = - \Gamma_{\alpha\sigma}^{\beta} T_{\beta}^{\alpha} \quad . \quad . \quad . \quad (57a)$$

The right side expresses the energetic effect of the gravitational field on matter.

Thus the field equations of gravitation contain four conditions which govern the course of material phenomena. They give the equations of material phenomena completely, if the latter is capable of being characterized by four differential equations independent of one another.\*

#### D. MATERIAL PHENOMENA

The mathematical aids developed in part B enable us forthwith to generalize the physical laws of matter (hydrodynamics, Maxwell's electrodynamics), as they are formulated in the special theory of relativity, so that they will fit in with the general theory of relativity. When this is done, the general principle of relativity does not indeed afford us a further limitation of possibilities; but it makes us acquainted with the influence of the gravitational field on all processes,

\* On this question cf. H. Hilbert, Nachr. d. K. Gesellsch. d. Wiss. zu Göttingen, Math.-phys. Klasse, 1915, p. 3.



without our having to introduce any new hypothesis whatever.

Hence it comes about that it is not necessary to introduce definite assumptions as to the physical nature of matter (in the narrower sense). In particular it may remain an open question whether the theory of the electromagnetic field in conjunction with that of the gravitational field furnishes a sufficient basis for the theory of matter or not. The general postulate of relativity is unable on principle to tell us anything about this. It must remain to be seen, during the working out of the theory, whether electromagnetics and the doctrine of gravitation are able in collaboration to perform what the former by itself is unable to do.

### § 19. Euler's Equations for a Frictionless Adiabatic Fluid

Let  $p$  and  $\rho$  be two scalars, the former of which we call the "pressure," the latter the "density" of a fluid; and let an equation subsist between them. Let the contravariant symmetrical tensor

$$T^{\alpha\beta} = -g^{\alpha\beta}p + \rho \frac{dx_\alpha}{ds} \frac{dx_\beta}{ds} . \quad (58)$$

be the contravariant energy-tensor of the fluid. To it belongs the covariant tensor

$$T_{\mu\nu} = -g_{\mu\nu}p + g_{\mu\alpha}g_{\nu\beta} \frac{dx_\alpha}{ds} \frac{dx_\beta}{ds} \rho, \quad (58a)$$

as well as the mixed tensor\*

$$T^\alpha_\sigma = -\delta^\alpha_\sigma p + g_{\sigma\beta} \frac{dx_\beta}{ds} \frac{dx_\alpha}{ds} \rho \quad (58b)$$

Inserting the right-hand side of (58b) in (57a), we obtain the Eulerian hydrodynamical equations of the general theory of relativity. They give, in theory, a complete solution of the problem of motion, since the four equations (57a), together

\* For an observer using a system of reference in the sense of the special theory of relativity for an infinitely small region, and moving with it, the density of energy  $T^4_4$  equals  $\rho - p$ . This gives the definition of  $\rho$ . Thus  $\rho$  is not constant for an incompressible fluid.

with the given equation between  $p$  and  $\rho$ , and the equation

$$g_{\alpha\beta} \frac{dx_\alpha}{ds} \frac{dx_\beta}{ds} = 1,$$

are sufficient,  $g_{\alpha\beta}$  being given, to define the six unknowns

$$p, \rho, \frac{dx_1}{ds}, \frac{dx_2}{ds}, \frac{dx_3}{ds}, \frac{dx_4}{ds}.$$

If the  $g_{\mu\nu}$  are also unknown, the equations (53) are brought in. These are eleven equations for defining the ten functions  $g_{\mu\nu}$ , so that these functions appear over-defined. We must remember, however, that the equations (57a) are already contained in the equations (53), so that the latter represent only seven independent equations. There is good reason for this lack of definition, in that the wide freedom of the choice of co-ordinates causes the problem to remain mathematically undefined to such a degree that three of the functions of space may be chosen at will.\*

## § 20. Maxwell's Electromagnetic Field Equations for Free Space

Let  $\phi_\nu$  be the components of a covariant vector—the electromagnetic potential vector. From them we form, in accordance with (36), the components  $F_{\rho\sigma}$  of the covariant six-vector of the electromagnetic field, in accordance with the system of equations

$$F_{\rho\sigma} = \frac{\partial \phi_\rho}{\partial x_\sigma} - \frac{\partial \phi_\sigma}{\partial x_\rho} \quad . \quad . \quad . \quad (59)$$

It follows from (59) that the system of equations

$$\frac{\partial F_{\rho\sigma}}{\partial x_\tau} + \frac{\partial F_{\sigma\tau}}{\partial x_\rho} + \frac{\partial F_{\tau\rho}}{\partial x_\sigma} = 0 \quad . \quad . \quad . \quad (60)$$

is satisfied, its left side being, by (37), an antisymmetrical tensor of the third rank. System (60) thus contains essentially four equations which are written out as follows:—

\* On the abandonment of the choice of co-ordinates with  $g = -1$ , there remain *four* functions of space with liberty of choice, corresponding to the four arbitrary functions at our disposal in the choice of co-ordinates.

$$\left. \begin{aligned} \frac{\partial F_{23}}{\partial x_4} + \frac{\partial F_{34}}{\partial x_2} + \frac{\partial F_{42}}{\partial x_3} &= 0 \\ \frac{\partial F_{34}}{\partial x_1} + \frac{\partial F_{41}}{\partial x_3} + \frac{\partial F_{13}}{\partial x_4} &= 0 \\ \frac{\partial F_{41}}{\partial x_2} + \frac{\partial F_{12}}{\partial x_4} + \frac{\partial F_{24}}{\partial x_1} &= 0 \\ \frac{\partial F_{12}}{\partial x_3} + \frac{\partial F_{23}}{\partial x_1} + \frac{\partial F_{31}}{\partial x_2} &= 0 \end{aligned} \right\} \quad (60a)$$

This system corresponds to the second of Maxwell's systems of equations. We recognize this at once by setting

$$\left. \begin{aligned} F_{23} &= H_x, & F_{14} &= E_x \\ F_{31} &= H_y, & F_{24} &= E_y \\ F_{12} &= H_z, & F_{34} &= E_z \end{aligned} \right\} \quad (61)$$

Then in place of (60a) we may set, in the usual notation of three-dimensional vector analysis,

$$\left. \begin{aligned} -\frac{\partial \mathbf{H}}{\partial t} &= \text{curl } \mathbf{E} \\ \text{div } \mathbf{H} &= 0 \end{aligned} \right\} \quad (60b)$$

We obtain Maxwell's first system by generalizing the form given by Minkowski. We introduce the contravariant six-vector associated with  $F^{\alpha\beta}$

$$F^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta} \quad (62)$$

and also the contravariant vector  $J^\mu$  of the density of the electric current. Then, taking (40) into consideration, the following equations will be invariant for any substitution whose invariant is unity (in agreement with the chosen co-ordinates) :—

$$\frac{\partial}{\partial x_\nu} F^{\mu\nu} = J^\mu \quad (63)$$

Let

$$\left. \begin{aligned} F^{23} &= H'_x, & F^{14} &= -E'_x \\ F^{31} &= H'_y, & F^{24} &= -E'_y \\ F^{12} &= H'_z, & F^{34} &= -E'_z \end{aligned} \right\} \quad (64)$$

which quantities are equal to the quantities  $H_x \dots E_z$  in

the special case of the restricted theory of relativity ; and in addition

$$J^1 = j_x, J^2 = j_y, J^3 = j_z, J^4 = \rho,$$

we obtain in place of (63)

$$\left. \begin{aligned} \frac{\partial \mathbf{E}'}{\partial t} + \mathbf{j} &= \text{curl } \mathbf{H}' \\ \text{div } \mathbf{E}' &= \rho \end{aligned} \right\} \quad . \quad . \quad . \quad (63a)$$

The equations (60), (62), and (63) thus form the generalization of Maxwell's field equations for free space, with the convention which we have established with respect to the choice of co-ordinates.

*The Energy-components of the Electromagnetic Field.*—We form the inner product

$$\kappa_\sigma = F_{\sigma\mu} J^\mu \quad . \quad . \quad . \quad (65)$$

By (61) its components, written in the three-dimensional manner, are

$$\left. \begin{aligned} \kappa_1 &= \rho E_x + [\mathbf{j} \cdot \mathbf{E}]^x \\ &\vdots \quad \vdots \quad \vdots \quad \vdots \\ \kappa_4 &= -(\mathbf{j} \cdot \mathbf{E}) \end{aligned} \right\} \quad . \quad . \quad . \quad (65a)$$

$\kappa_\sigma$  is a covariant vector the components of which are equal to the negative momentum, or, respectively, the energy, which is transferred from the electric masses to the electromagnetic field per unit of time and volume. If the electric masses are free, that is, under the sole influence of the electromagnetic field, the covariant vector  $\kappa_\sigma$  will vanish.

To obtain the energy-components  $T_\sigma^\nu$  of the electromagnetic field, we need only give to equation  $\kappa_\sigma = 0$  the form of equation (57). From (63) and (65) we have in the first place

$$\kappa_\sigma = F_{\sigma\mu} \frac{\partial F^{\mu\nu}}{\partial x_\nu} = \frac{\partial}{\partial x_\nu} (F_{\sigma\mu} F^{\mu\nu}) - F^{\mu\rho} \frac{\partial F_{\sigma\mu}}{\partial x_\nu}.$$

The second term of the right-hand side, by reason of (60), permits the transformation

$$F^{\mu\nu} \frac{\partial F_{\sigma\mu}}{\partial x_\nu} = -\frac{1}{2} F^{\mu\nu} \frac{\partial F_{\mu\nu}}{\partial x_\sigma} = -\frac{1}{2} g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta} \frac{\partial F_{\mu\nu}}{\partial x_\sigma},$$

which latter expression may, for reasons of symmetry, also be written in the form

$$- \frac{1}{4} \left[ g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta} \frac{\partial F_{\mu\nu}}{\partial x_\sigma} + g^{\mu\alpha} g^{\nu\beta} \frac{\partial F_{\alpha\beta}}{\partial x_\sigma} F_{\mu\nu} \right].$$

But for this we may set

$$- \frac{1}{4} \frac{\partial}{\partial x_\sigma} (g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta} F_{\mu\nu}) + \frac{1}{4} F_{\alpha\beta} F_{\mu\nu} \frac{\partial}{\partial x_\sigma} (g^{\mu\alpha} g^{\nu\beta}).$$

The first of these terms is written more briefly

$$- \frac{1}{4} \frac{\partial}{\partial x_\sigma} (F^{\mu\nu} F_{\mu\nu});$$

the second, after the differentiation is carried out, and after some reduction, results in

$$- \frac{1}{2} F^{\mu\tau} F_{\mu\nu} g^{\nu\rho} \frac{\partial g_{\sigma\tau}}{\partial x_\sigma}.$$

Taking all three terms together we obtain the relation

$$\kappa_\sigma = \frac{\partial T_\sigma^\nu}{\partial x_\nu} - \frac{1}{2} g^{\tau\mu} \frac{\partial g_{\mu\nu}}{\partial x_\sigma} T_\tau^\nu. \quad (66)$$

where

$$T_\sigma^\nu = - F_{\sigma\alpha} F^{\nu\alpha} + \frac{1}{4} \delta_\sigma^\nu F_{\alpha\beta} F^{\alpha\beta}.$$

Equation (66), if  $\kappa_\sigma$  vanishes, is, on account of (30), equivalent to (57) or (57a) respectively. Therefore the  $T_\sigma^\nu$  are the energy-components of the electromagnetic field. With the help of (61) and (64), it is easy to show that these energy-components of the electromagnetic field in the case of the special theory of relativity give the well-known Maxwell-Poynting expressions.

We have now deduced the general laws which are satisfied by the gravitational field and matter, by consistently using a system of co-ordinates for which  $\sqrt{-g} = 1$ . We have thereby achieved a considerable simplification of formulæ and calculations, without failing to comply with the requirement of general covariance; for we have drawn our equations from generally covariant equations by specializing the system of co-ordinates.

Still the question is not without a formal interest, whether with a correspondingly generalized definition of the energy-components of gravitational field and matter, even without specializing the system of co-ordinates, it is possible to formulate laws of conservation in the form of equation (56), and field equations of gravitation of the same nature as (52) or (52a), in such a manner that on the left we have a divergence (in the ordinary sense), and on the right the sum of the energy-components of matter and gravitation. I have found that in both cases this is actually so. But I do not think that the communication of my somewhat extensive reflexions on this subject would be worth while, because after all they do not give us anything that is materially new.

## E

## § 21. Newton's Theory as a First Approximation

As has already been mentioned more than once, the special theory of relativity as a special case of the general theory is characterized by the  $g_{\mu\nu}$  having the constant values (4). From what has already been said, this means complete neglect of the effects of gravitation. We arrive at a closer approximation to reality by considering the case where the  $g_{\mu\nu}$  differ from the values of (4) by quantities which are small compared with 1, and neglecting small quantities of second and higher order. (First point of view of approximation.)

It is further to be assumed that in the space-time territory under consideration the  $g_{\mu\nu}$  at spatial infinity, with a suitable choice of co-ordinates, tend toward the values (4); i.e. we are considering gravitational fields which may be regarded as generated exclusively by matter in the finite region.

It might be thought that these approximations must lead us to Newton's theory. But to that end we still need to approximate the fundamental equations from a second point of view. We give our attention to the motion of a material point in accordance with the equations (16). In the case of the special theory of relativity the components

$$\frac{dx_1}{ds}, \frac{dx_2}{ds}, \frac{dx_3}{ds}$$

may take on any values. This signifies that any velocity

$$v = \sqrt{\left(\frac{dx_1}{dx_4}\right)^2 + \left(\frac{dx_2}{dx_4}\right)^2 + \left(\frac{dx_3}{dx_4}\right)^2}$$

may occur, which is less than the velocity of light *in vacuo*. If we restrict ourselves to the case which almost exclusively offers itself to our experience, of  $v$  being small as compared with the velocity of light, this denotes that the components

$$\frac{dx_1}{ds}, \frac{dx_2}{ds}, \frac{dx_3}{ds}$$

are to be treated as small quantities, while  $dx_4/ds$ , to the second order of small quantities, is equal to one. (Second point of view of approximation.)

Now we remark that from the first point of view of approximation the magnitudes  $\Gamma_{\mu\nu}^\tau$  are all small magnitudes of at least the first order. A glance at (46) thus shows that in this equation, from the second point of view of approximation, we have to consider only terms for which  $\mu = \nu = 4$ . Restricting ourselves to terms of lowest order we first obtain in place of (46) the equations

$$\frac{d^2 x_\tau}{ds^2} = \Gamma_{44}^\tau$$

where we have set  $ds = dx_4 = dt$ ; or with restriction to terms which from the first point of view of approximation are of first order:—

$$\frac{d^2 x_\tau}{dt^2} = [44, \tau] \quad (\tau = 1, 2, 3)$$

$$\frac{d^2 x_4}{dt^2} = -[44, 4].$$

If in addition we suppose the gravitational field to be a quasi-static field, by confining ourselves to the case where the motion of the matter generating the gravitational field is but slow (in comparison with the velocity of the propagation of light), we may neglect on the right-hand side differentiations with respect to the time in comparison with those with respect to the space co-ordinates, so that we have

$$\frac{d^2 x_\tau}{dt^2} = -\frac{1}{2} \frac{\partial g_{44}}{\partial x_\tau} \quad (\tau = 1, 2, 3) \quad (67)$$

This is the equation of motion of the material point according to Newton's theory, in which  $\frac{1}{2}g_{44}$  plays the part of the gravitational potential. What is remarkable in this result is that the component  $g_{44}$  of the fundamental tensor alone defines, to a first approximation, the motion of the material point.

We now turn to the field equations (53). Here we have to take into consideration that the energy-tensor of "matter" is almost exclusively defined by the density of matter in the narrower sense, i.e. by the second term of the right-hand side of (58) [or, respectively, (58a) or (58b)]. If we form the approximation in question, all the components vanish with the one exception of  $T_{44} = \rho = T$ . On the left-hand side of (53) the second term is a small quantity of second order; the first yields, to the approximation in question,

$$\frac{\partial}{\partial x_1}[\mu\nu, 1] + \frac{\partial}{\partial x_2}[\mu\nu, 2] + \frac{\partial}{\partial x_3}[\mu\nu, 3] - \frac{\partial}{\partial x_4}[\mu\nu, 4].$$

For  $\mu = \nu = 4$ , this gives, with the omission of terms differentiated with respect to time,

$$-\frac{1}{2} \left( \frac{\partial^2 g_{44}}{\partial x_1^2} + \frac{\partial^2 g_{44}}{\partial x_2^2} + \frac{\partial^2 g_{44}}{\partial x_3^2} \right) = -\frac{1}{2} \nabla^2 g_{44}.$$

The last of equations (53) thus yields

$$\nabla^2 g_{44} = \kappa \rho \quad (68)$$

The equations (67) and (68) together are equivalent to Newton's law of gravitation.

By (67) and (68) the expression for the gravitational potential becomes

$$-\frac{\kappa}{8\pi} \int \frac{\rho d\tau}{r} \quad (68a)$$

while Newton's theory, with the unit of time which we have chosen, gives

$$-\frac{K}{c^2} \int \frac{\rho d\tau}{r}$$



in which  $K$  denotes the constant  $6.7 \times 10^{-8}$ , usually called the constant of gravitation. By comparison we obtain

$$\kappa = \frac{8\pi K}{c^2} = 1.87 \times 10^{-27} \quad . \quad . \quad (69)$$

## § 22. Behaviour of Rods and Clocks in the Static Gravitational Field. Bending of Light-rays. Motion of the Perihelion of a Planetary Orbit

To arrive at Newton's theory as a first approximation we had to calculate only one component,  $g_{44}$ , of the ten  $g_{\mu\nu}$  of the gravitational field, since this component alone enters into the first approximation, (67), of the equation for the motion of the material point in the gravitational field. From this, however, it is already apparent that other components of the  $g_{\mu\nu}$  must differ from the values given in (4) by small quantities of the first order. This is required by the condition  $g = -1$ .

For a field-producing point mass at the origin of co-ordinates, we obtain, to the first approximation, the radially symmetrical solution

$$\left. \begin{aligned} g_{\rho\sigma} &= -\delta_{\rho\sigma} - \alpha \frac{x_\rho x_\sigma}{r^3} \quad (\rho, \sigma = 1, 2, 3) \\ g_{\rho 4} &= g_{4\rho} = 0 \quad (\rho = 1, 2, 3) \\ g_{44} &= 1 - \frac{\alpha}{r} \end{aligned} \right\} \quad . \quad (70)$$

where  $\delta_{\rho\sigma}$  is 1 or 0, respectively, accordingly as  $\rho = \sigma$  or  $\rho \neq \sigma$ , and  $r$  is the quantity  $+\sqrt{x_1^2 + x_2^2 + x_3^2}$ . On account of (68a)

$$\alpha = \frac{\kappa M}{4\pi}, \quad . \quad . \quad . \quad (70a)$$

if  $M$  denotes the field-producing mass. It is easy to verify that the field equations (outside the mass) are satisfied to the first order of small quantities.

We now examine the influence exerted by the field of the mass  $M$  upon the metrical properties of space. The relation

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu$$

always holds between the "locally" (§ 4) measured lengths and times  $ds$  on the one hand, and the differences of co-ordinates  $dx_\nu$ , on the other hand.

For a unit-measure of length laid "parallel" to the axis of  $x$ , for example, we should have to set  $ds^2 = -1$ ;  $dx_2 = dx_3 = dx_4 = 0$ . Therefore  $-1 = g_{11}dx_1^2$ . If, in addition, the unit-measure lies on the axis of  $x$ , the first of equations (70) gives

$$g_{11} = -\left(1 + \frac{a}{r}\right).$$

From these two relations it follows that, correct to a first order of small quantities,

$$dx = 1 - \frac{a}{2r} \quad . \quad . \quad . \quad (71)$$

The unit measuring-rod thus appears a little shortened in relation to the system of co-ordinates by the presence of the gravitational field, if the rod is laid along a radius.

In an analogous manner we obtain the length of co-ordinates in tangential direction if, for example, we set

$$ds^2 = -1; dx_1 = dx_3 = dx_4 = 0; x_1 = r, x_2 = x_3 = 0.$$

The result is

$$-1 = g_{22}dx_2^2 = -dx_2^2 \quad . \quad . \quad . \quad (71a)$$

With the tangential position, therefore, the gravitational field of the point of mass has no influence on the length of a rod.

Thus Euclidean geometry does not hold even to a first approximation in the gravitational field, if we wish to take one and the same rod, independently of its place and orientation, as a realization of the same interval; although, to be sure, a glance at (70a) and (69) shows that the deviations to be expected are much too slight to be noticeable in measurements of the earth's surface.

Further, let us examine the rate of a unit clock, which is arranged to be at rest in a static gravitational field. Here we have for a clock period  $ds = 1$ ;  $dx_1 = dx_2 = dx_3 = 0$ . Therefore

$$1 = g_{44}dx_4^2;$$

$$dx_4 = \frac{1}{\sqrt{g_{44}}} = \frac{1}{\sqrt{(1 + (g_{44} - 1))}} = 1 - \frac{1}{2}(g_{44} - 1)$$

or

$$dx_4 = 1 + \frac{\kappa}{8\pi} \int \frac{\rho dr}{r} \quad . \quad . \quad . \quad (72)$$

Thus the clock goes more slowly if set up in the neighbourhood of ponderable masses. From this it follows that the spectral lines of light reaching us from the surface of large stars must appear displaced towards the red end of the spectrum.\*

We now examine the course of light-rays in the static gravitational field. By the special theory of relativity the velocity of light is given by the equation

$$-dx_1^2 - dx_2^2 - dx_3^2 + dx_4^2 = 0$$

and therefore by the general theory of relativity by the equation

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu = 0 \quad . \quad . \quad . \quad (73)$$

If the direction, i.e. the ratio  $dx_1 : dx_2 : dx_3$  is given, equation (73) gives the quantities

$$\frac{dx_1}{dx_4}, \frac{dx_2}{dx_4}, \frac{dx_3}{dx_4}$$

and accordingly the velocity

$$\sqrt{\left(\frac{dx_1}{dx_4}\right)^2 + \left(\frac{dx_2}{dx_4}\right)^2 + \left(\frac{dx_3}{dx_4}\right)^2} = \gamma$$

defined in the sense of Euclidean geometry. We easily recognize that the course of the light-rays must be bent with regard to the system of co-ordinates, if the  $g_{\mu\nu}$  are not constant. If  $n$  is a direction perpendicular to the propagation of light, the Huyghens principle shows that the light-ray, envisaged in the plane  $(\gamma, n)$ , has the curvature  $-\partial\gamma/\partial n$ .

We examine the curvature undergone by a ray of light passing by a mass  $M$  at the distance  $\Delta$ . If we choose the system of co-ordinates in agreement with the accompanying diagram, the total bending of the ray (calculated positively if

\* According to E. Freundlich, spectroscopical observations on fixed stars of certain types indicate the existence of an effect of this kind, but a crucial test of this consequence has not yet been made.

concave towards the origin) is given in sufficient approximation by

$$B = \int_{-\infty}^{+\infty} \frac{\partial \gamma}{\partial x_1} dx_2,$$

while (73) and (70) give

$$\gamma = \sqrt{\left(-\frac{g_{44}}{g_{22}}\right)} = 1 - \frac{a}{2r} \left(1 + \frac{x_2^2}{r^2}\right).$$

Carrying out the calculation, this gives

$$B = \frac{2a}{\Delta} = \frac{\kappa M}{2\pi \Delta}. \quad (74)$$




FIG. 8.

According to this, a ray of light going past the sun undergoes a deflexion of  $1.7''$ ; and a ray going past the planet Jupiter a deflexion of about  $.02''$ .

If we calculate the gravitational field to a higher degree of approximation, and likewise with corresponding accuracy the orbital motion of a material point of relatively infinitely small mass, we find a deviation of the following kind from the Kepler-Newton laws of planetary motion. The orbital ellipse of a planet undergoes a slow rotation, in the direction of motion, of amount

$$e = 24\pi^3 \frac{a^3}{T^2 c^2 (1 - e^2)} \quad (75)$$

per revolution. In this formula  $a$  denotes the axis,  $c$  the velocity of light in the usual measure,  $e$  the eccentricity,  $T$  the time of revolution in second

.. Calculation gives for the planet Mercury a shift of orbit of 43" per century, corresponding exactly to observation (Leverrier); for the astronomers have found in the motion of the perihelion of this planet, after subtracting the effect of disturbances by other planets, an inexplicable shift of this magnitude.

\* For the calculation I refer to the original paper in *Sitzungsber. d. Preuss. Akad. d. Wiss.*, 1915, p. 831; *ibid.*, 1916, p. 189.

# HAMILTON'S PRINCIPLE AND THE GENERAL THEORY OF RELATIVITY

BY

A. EINSTEIN

*Translated from "Hamiltonsches Princip und allgemeine  
Relativitätstheorie," Sitzungsberichte der Preussi-  
schen Akad. d. Wissenschaften, 1916.*



# HAMILTON'S PRINCIPLE AND THE GENERAL THEORY OF RELATIVITY

By A. EINSTEIN

THE general theory of relativity has recently been given in a particularly clear form by H. A. Lorentz and D. Hilbert,\* who have deduced its equations from one single principle of variation. The same thing will be done in the present paper. But my purpose here is to present the fundamental connexions in as perspicuous a manner as possible, and in as general terms as is permissible from the point of view of the general theory of relativity. In particular we shall make as few specializing assumptions as possible, in marked contrast to Hilbert's treatment of the subject. On the other hand, in antithesis to my own most recent treatment of the subject, there is to be complete liberty in the choice of the system of co-ordinates.

## § 1. The Principle of Variation and the Field-equations of Gravitation and Matter

Let the gravitational field be described as usual by the tensor† of the  $g_{\mu\nu}$  (or the  $g^{\mu\nu}$ ); and matter, including the electromagnetic field, by any number of space-time functions  $q_{(\rho)}$ . How these functions may be characterized in the theory of invariants does not concern us. Further, let  $\mathfrak{S}$  be a function of the

$$g^{\mu\nu}, g^{\mu\nu}_{,\sigma} \left( = \frac{\partial g^{\mu\nu}}{\partial x_{\sigma}} \right) \text{ and } g^{\mu\nu}_{,\sigma\tau} \left( = \frac{\partial^2 g^{\mu\nu}}{\partial x_{\sigma} \partial x_{\tau}} \right), \text{ the } q_{(\rho)} \text{ and } q_{(\rho),\alpha} \left( = \frac{\partial q_{(\rho)}}{\partial x_{\alpha}} \right).$$

\* Four papers by Lorentz in the Publications of the Koninkl. Akad. van Wetensch. te Amsterdam, 1915 and 1916; D. Hilbert, Göttinger Nachr., 1915, Part 3.

† No use is made for the present of the tensor character of the  $g_{\mu\nu}$ .



The principle of variation

$$\delta \int \mathfrak{H} d\tau = 0 \quad . \quad . \quad . \quad (1)$$

then gives us as many differential equations as there are functions  $g_{\mu\nu}$  and  $q_{(\rho)}$  to be defined, if the  $g^{\mu\nu}$  and  $q_{(\rho)}$  are varied independently of one another, and in such a way that at the limits of integration the  $\delta q_{(\rho)}$ ,  $\delta g^{\mu\nu}$ , and  $\frac{\partial}{\partial x_\sigma} (\delta g^{\mu\nu})$  all vanish.

We will now assume that  $\mathfrak{H}$  is linear in the  $g_{\sigma\tau}$ , and that the coefficients of the  $g_{\sigma\tau}^{\mu\nu}$  depend only on the  $g^{\mu\nu}$ . We may then replace the principle of variation (1) by one which is more convenient for us. For by appropriate partial integration we obtain

$$\int \mathfrak{H} d\tau = \int \mathfrak{H}^* d\tau + F \quad . \quad . \quad . \quad (2)$$

where  $F$  denotes an integral over the boundary of the domain in question, and  $\mathfrak{H}^*$  depends only on the  $g^{\mu\nu}$ ,  $g_{\sigma\tau}^{\mu\nu}$ ,  $q_{(\rho)}$ ,  $q_{(\rho)\alpha}$ , and no longer on the  $g_{\sigma\tau}^{\mu\nu}$ . From (2) we obtain, for such variations as are of interest to us,

$$\delta \int \mathfrak{H} d\tau = \delta \int \mathfrak{H}^* d\tau, \quad . \quad . \quad . \quad (3)$$

so that we may replace our principle of variation (1) by the more convenient form

$$\delta \int \mathfrak{H}^* d\tau = 0. \quad . \quad . \quad . \quad (1a)$$

By carrying out the variation of the  $g^{\mu\nu}$  and the  $q_{(\rho)}$  we obtain, as field-equations of gravitation and matter, the equations †

$$\frac{\partial}{\partial x_\alpha} \left( \frac{\partial \mathfrak{H}^*}{\partial g_\alpha^{\mu\nu}} \right) - \frac{\partial \mathfrak{H}^*}{\partial g^{\mu\nu}} = 0 \quad . \quad . \quad . \quad (4)$$

$$\frac{\partial}{\partial x_\alpha} \left( \frac{\partial \mathfrak{H}^*}{\partial q_{(\rho)\alpha}} \right) - \frac{\partial \mathfrak{H}^*}{\partial q_{(\rho)}} = 0 \quad . \quad . \quad . \quad (5)$$

† For brevity the summation symbols are omitted in the formulae. Indices occurring twice in a term are always to be taken as summed. Thus in (4), for example,  $\frac{\partial}{\partial x_\alpha} \left( \frac{\partial \mathfrak{H}^*}{\partial g_\alpha^{\mu\nu}} \right)$  denotes the term  $\sum_\alpha \frac{\partial}{\partial x_\alpha} \left( \frac{\partial \mathfrak{H}^*}{\partial g_\alpha^{\mu\nu}} \right)$ .

## § 2. Separate Existence of the Gravitational Field

If we make no restrictive assumption as to the manner in which  $\mathfrak{H}$  depends on the  $g^{\mu\nu}$ ,  $g_{\sigma}^{\mu\nu}$ ,  $g_{\sigma\tau}^{\mu\nu}$ ,  $q_{(\rho)}$ ,  $q_{(\rho)\alpha}$ , the energy-components cannot be divided into two parts, one belonging to the gravitational field, the other to matter. To ensure this feature of the theory, we make the following assumption

$$\mathfrak{H} = \mathfrak{G} + \mathfrak{M} \quad . \quad . \quad . \quad (6)$$

where  $\mathfrak{G}$  is to depend only on the  $g^{\mu\nu}$ ,  $g_{\sigma}^{\mu\nu}$ ,  $g_{\sigma\tau}^{\mu\nu}$ , and  $\mathfrak{M}$  only on  $g^{\mu\nu}$ ,  $q_{(\rho)}$ ,  $q_{(\rho)\alpha}$ . Equations (4), (4a) then assume the form

$$\frac{\partial}{\partial x_{\alpha}} \left( \frac{\partial \mathfrak{G}^*}{\partial g_{\alpha}^{\mu\nu}} \right) - \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\nu}} = \frac{\partial \mathfrak{M}}{\partial g^{\mu\nu}} \quad . \quad . \quad . \quad (7)$$

$$\frac{\partial}{\partial x_{\alpha}} \left( \frac{\partial \mathfrak{M}}{\partial q_{(\rho)\alpha}} \right) - \frac{\partial \mathfrak{M}}{\partial q_{(\rho)}} = 0 \quad . \quad . \quad . \quad (8)$$

Here  $\mathfrak{G}^*$  stands in the same relation to  $\mathfrak{G}$  as  $\mathfrak{H}^*$  to  $\mathfrak{H}$ .

It is to be noted carefully that equations (8) or (5) would have to give way to others, if we were to assume  $\mathfrak{M}$  or  $\mathfrak{H}$  to be also dependent on derivatives of the  $q_{(\rho)}$  of order higher than the first. Likewise it might be imaginable that the  $q_{(\rho)}$  would have to be taken, not as independent of one another, but as connected by conditional equations. All this is of no importance for the following developments, as these are based solely on the equations (7), which have been found by varying our integral with respect to the  $g^{\mu\nu}$ .

## § 3. Properties of the Field Equations of Gravitation Conditioned by the Theory of Invariants

We now introduce the assumption that

$$ds^2 = g_{\mu\nu} dx_{\mu} dx_{\nu} \quad . \quad . \quad . \quad (9)$$

is an invariant. This determines the transformational character of the  $g_{\mu\nu}$ . As to the transformational character of the  $q_{(\rho)}$ , which describe matter, we make no supposition. On the other hand, let the functions  $H = \frac{\mathfrak{H}}{\sqrt{-g}}$ , as well as

$G = \frac{\Theta}{\sqrt{-g}}$ , and  $M = \frac{\mathfrak{M}}{\sqrt{-g}}$ , be invariants in relation to any substitutions of space-time co-ordinates. From these assumptions follows the general covariance of the equations (7) and (8), deduced from (1). It further follows that  $G$  (apart from a constant factor) must be equal to the scalar of Riemann's tensor of curvature; because there is no other invariant with the properties required for  $G$ .† Thereby  $\Theta^*$  is also perfectly determined, and consequently the left-hand side of field equation (7) as well.‡

From the general postulate of relativity there follow certain properties of the function  $\Theta^*$  which we shall now deduce. For this purpose we carry through an infinitesimal transformation of the co-ordinates, by setting

$$x'_\nu = x_\nu + \Delta x_\nu. \quad (10)$$

where the  $\Delta x_\nu$  are arbitrary, infinitely small functions of the co-ordinates, and  $x'_\nu$  are the co-ordinates, in the new system, of the world-point having the co-ordinates  $x_\nu$  in the original system. As for the co-ordinates, so too for any other magnitude  $\psi$ , a law of transformation holds good, of the type

$$\psi' = \psi + \Delta\psi,$$

where  $\Delta\psi$  must always be expressible by the  $\Delta x_\nu$ . From the covariant property of the  $g^{\mu\nu}$  we easily deduce for the  $g^{\mu\nu}$  and  $g^{\mu\nu}_\sigma$  the laws of transformation

$$\Delta g^{\mu\nu} = g^{\mu\alpha} \frac{\partial(\Delta x_\nu)}{\partial x_\alpha} + g^{\nu\alpha} \frac{\partial(\Delta x_\mu)}{\partial x_\alpha} \quad (11)$$

$$\Delta g^{\mu\nu}_\sigma = \frac{\partial(\Delta g^{\mu\nu})}{\partial x_\sigma} - g^{\mu\nu}_\alpha \frac{\partial(\Delta x_\alpha)}{\partial x_\sigma} \quad (12)$$

Since  $\Theta^*$  depends only on the  $g^{\mu\nu}$  and  $g^{\mu\nu}_\sigma$ , it is possible, with the help of (11) and (12), to calculate  $\Delta\Theta^*$ . We thus obtain the equation

$$\sqrt{-g} \Delta \left( \frac{\Theta^*}{\sqrt{-g}} \right) = S^\nu_\sigma \frac{\partial(\Delta x_\sigma)}{\partial x_\nu} + 2 \frac{\partial\Theta^*}{\partial g^{\mu\sigma}} g^{\mu\nu} \frac{\partial^2 \Delta x_\sigma}{\partial x_\nu \partial x_\alpha}, \quad (13)$$

† Herein is to be found the reason why the general postulate of relativity leads to a very definite theory of gravitation.

‡ By performing partial integration we obtain

$$\Theta^* = \sqrt{-g} g^{\mu\nu} [\{\mu\alpha, \beta\} \{\nu\beta, \alpha\} - \{\mu\nu, \alpha\} \{\alpha\beta, \beta\}].$$

where for brevity we have set

$$S_{\sigma}^{\nu} = 2 \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\sigma}} g^{\mu\nu} + 2 \frac{\partial \mathfrak{G}^*}{\partial g_{\alpha}^{\mu\sigma}} g_{\alpha}^{\mu\nu} + \mathfrak{G}^* \delta_{\sigma}^{\nu} - \frac{\partial \mathfrak{G}^*}{\partial g_{\nu}^{\mu\alpha}} g_{\sigma}^{\mu\alpha}. \quad (14)$$

From these two equations we draw two inferences which are important for what follows. We know that  $\frac{\mathfrak{G}}{\sqrt{-g}}$  is an invariant with respect to any substitution, but we do not know this of  $\frac{\mathfrak{G}^*}{\sqrt{-g}}$ . It is easy to demonstrate, however, that the latter quantity is an invariant with respect to any *linear* substitutions of the co-ordinates. Hence it follows that the right side of (13) must always vanish if all  $\frac{\partial^2 \Delta x_{\sigma}}{\partial x_{\nu} \partial x_{\alpha}}$  vanish. Consequently  $\mathfrak{G}^*$  must satisfy the identity

$$S_{\sigma}^{\nu} \equiv 0 \quad . \quad . \quad . \quad . \quad (15)$$

If, further, we choose the  $\Delta x_{\sigma}$ , so that they differ from zero only in the interior of a given domain, but in infinitesimal proximity to the boundary they vanish, then, with the transformation in question, the value of the boundary integral occurring in equation (2) does not change. Therefore  $\Delta F = 0$ , and, in consequence,†

$$\Delta \int \mathfrak{G} d\tau = \Delta \int \mathfrak{G}^* d\tau.$$

But the left-hand side of the equation must vanish, since both  $\frac{\mathfrak{G}}{\sqrt{-g}}$  and  $\sqrt{-g} d\tau$  are invariants. Consequently the right-hand side also vanishes. Thus, taking (14), (15), and (16) into consideration, we obtain, in the first place, the equation

$$\int \frac{\partial \mathfrak{G}^*}{\partial g_{\alpha}^{\mu\sigma}} g^{\mu\nu} \frac{\partial^2 (\Delta x_{\sigma})}{\partial x_{\nu} \partial x_{\alpha}} d\tau = 0 \quad . \quad . \quad . \quad (16)$$

Transforming this equation by two partial integrations, and having regard to the liberty of choice of the  $\Delta x_{\sigma}$ , we obtain

† By the introduction of the quantities  $\mathfrak{G}$  and  $\mathfrak{G}^*$  instead of  $\mathfrak{G}$  and  $\mathfrak{G}^*$ .

the identity

$$\frac{\partial^2}{\partial x_\nu \partial x_\alpha} \left( g^{\mu\nu} \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\sigma}} \right) \equiv 0 \quad . \quad . \quad . \quad (17)$$

From the two identities (16) and (17), which result from the invariance of  $\frac{\mathfrak{G}}{\sqrt{-g}}$ , and therefore from the postulate of general relativity, we now have to draw conclusions.

We first transform the field equations (7) of gravitation by mixed multiplication by  $g^{\mu\sigma}$ . We then obtain (by interchanging the indices  $\sigma$  and  $\nu$ ), as equivalents of the field equations (7), the equations

$$\frac{\partial}{\partial x_\alpha} \left( g^{\mu\nu} \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\sigma}} \right) = - (\mathfrak{T}_\sigma^\nu + t_\sigma^\nu) \quad . \quad . \quad . \quad (18)$$

where we have set

$$\mathfrak{T}_\sigma^\nu = - \frac{\partial \mathfrak{M}}{\partial g^{\mu\sigma}} g^{\mu\nu} \quad . \quad . \quad . \quad . \quad . \quad . \quad (19)$$

$$t_\sigma^\nu = - \left( \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\sigma}} g^{\mu\nu} + \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\sigma}} g^{\mu\nu} \right) = \frac{1}{2} \left( \mathfrak{G}^* \delta_\sigma^\nu - \frac{\partial \mathfrak{G}^*}{\partial g^{\mu\alpha}} g_\sigma^{\mu\alpha} \right) \quad (20)$$

The last expression for  $t_\sigma^\nu$  is vindicated by (14) and (15). By differentiation of (18) with respect to  $x_\nu$ , and summation for  $\nu$ , there follows, in view of (17),

$$\frac{\partial}{\partial x_\nu} (\mathfrak{T}_\sigma^\nu + t_\sigma^\nu) = 0 \quad . \quad . \quad . \quad (21)$$

Equation (21) expresses the conservation of momentum and energy. We call  $\mathfrak{T}_\sigma^\nu$  the components of the energy of matter,  $t_\sigma^\nu$  the components of the energy of the gravitational field.

Having regard to (20), there follows from the field equations (7) of gravitation, by multiplication by  $g_\sigma^{\mu\nu}$  and summation with respect to  $\mu$  and  $\nu$ ,

$$\frac{\partial t_\sigma^\nu}{\partial x_\nu} + \frac{1}{2} g_\sigma^{\mu\nu} \frac{\partial \mathfrak{M}}{\partial g^{\mu\nu}} = 0,$$

or, in view of (19) and (21),

$$\frac{\partial \mathfrak{T}_{\sigma}^{\nu}}{\partial x_{\nu}} + \frac{1}{2} g_{\sigma}^{\mu\nu} \mathfrak{T}_{\mu\nu} = 0 \quad . \quad . \quad . \quad (22)$$

where  $\mathfrak{T}_{\mu\nu}$  denotes the quantities  $g_{\nu\sigma} \mathfrak{T}_{\mu}^{\sigma}$ . These are four equations which the energy-components of matter have to satisfy.

It is to be emphasized that the (generally covariant) laws of conservation (21) and (22) are deduced from the field equations (7) of gravitation, in combination with the postulate of general covariance (relativity) *alone*, without using the field equations (8) for material phenomena.



# COSMOLOGICAL CONSIDERATIONS ON THE GENERAL THEORY OF RELATIVITY

BY

A. EINSTEIN

*Translated from "Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie," Sitzungsberichte der Preussischen Akad. d. Wissenschaften, 1917.*





# COSMOLOGICAL CONSIDERATIONS ON THE GENERAL THEORY OF RELATIVITY

By A. EINSTEIN

IT is well known that Poisson's equation

$$\nabla^2\phi = 4\pi K\rho \quad (1)$$

in combination with the equations of motion of a material point is not as yet a perfect substitute for Newton's theory of action at a distance. There is still to be taken into account the condition that at spatial infinity the potential  $\phi$  tends toward a fixed limiting value. There is an analogous state of things in the theory of gravitation in general relativity. Here, too, we must supplement the differential equations by limiting conditions at spatial infinity, if we really have to regard the universe as being of infinite spatial extent.

In my treatment of the planetary problem I chose these limiting conditions in the form of the following assumption : it is possible to select a system of reference so that at spatial infinity all the gravitational potentials  $g_{\mu\nu}$  become constant. But it is by no means evident *a priori* that we may lay down the same limiting conditions when we wish to take larger portions of the physical universe into consideration. In the following pages the reflexions will be given which, up to the present, I have made on this fundamentally important question.

## § 1. The Newtonian Theory

It is well known that Newton's limiting condition of the constant limit for  $\phi$  at spatial infinity leads to the view that the density of matter becomes zero at infinity. For we imagine that there may be a place in universal space round about which the gravitational field of matter, viewed on a large scale, possesses spherical symmetry. It then follows from Poisson's equation that, in order that  $\phi$  may tend to a

## 178 COSMOLOGICAL CONSIDERATIONS

limit at infinity, the mean density  $\rho$  must decrease toward zero more rapidly than  $1/r^2$  as the distance  $r$  from the centre increases.\* In this sense, therefore, the universe according to Newton is finite, although it may possess an infinitely great total mass.

From this it follows in the first place that the radiation emitted by the heavenly bodies will, in part, leave the Newtonian system of the universe, passing radially outwards, to become ineffective and lost in the infinite. May not entire heavenly bodies fare likewise? It is hardly possible to give a negative answer to this question. For it follows from the assumption of a finite limit for  $\phi$  at spatial infinity that a heavenly body with finite kinetic energy is able to reach spatial infinity by overcoming the Newtonian forces of attraction. By statistical mechanics this case must occur from time to time, as long as the total energy of the stellar system—transferred to one single star—is great enough to send that star on its journey to infinity, whence it never can return.

We might try to avoid this peculiar difficulty by assuming a very high value for the limiting potential at infinity. That would be a possible way, if the value of the gravitational potential were not itself necessarily conditioned by the heavenly bodies. The truth is that we are compelled to regard the occurrence of any great differences of potential of the gravitational field as contradicting the facts. These differences must really be of so low an order of magnitude that the stellar velocities generated by them do not exceed the velocities actually observed.

If we apply Boltzmann's law of distribution for gas molecules to the stars, by comparing the stellar system with a gas in thermal equilibrium, we find that the Newtonian stellar system cannot exist at all. For there is a finite ratio of densities corresponding to the finite difference of potential between the centre and spatial infinity. A vanishing of the density at infinity thus implies a vanishing of the density at the centre.

\*  $\rho$  is the mean density of matter, calculated for a region which is large as compared with the distance between neighbouring fixed stars, but small in comparison with the dimensions of the whole stellar system.

It seems hardly possible to surmount these difficulties on the basis of the Newtonian theory. We may ask ourselves the question whether they can be removed by a modification of the Newtonian theory. First of all we will indicate a method which does not in itself claim to be taken seriously; it merely serves as a foil for what is to follow. In place of Poisson's equation we write

$$\nabla^2\phi - \lambda\phi = 4\pi\kappa\rho \quad . \quad . \quad . \quad (2)$$

where  $\lambda$  denotes a universal constant. If  $\rho_0$  be the uniform density of a distribution of mass, then

$$\phi = -\frac{4\pi\kappa}{\lambda}\rho_0 \quad . \quad . \quad . \quad (3)$$

is a solution of equation (2). This solution would correspond to the case in which the matter of the fixed stars was distributed uniformly through space, if the density  $\rho_0$  is equal to the actual mean density of the matter in the universe. The solution then corresponds to an infinite extension of the central space, filled uniformly with matter. If, without making any change in the mean density, we imagine matter to be non-uniformly distributed locally, there will be, over and above the  $\phi$  with the constant value of equation (3), an additional  $\phi$ , which in the neighbourhood of denser masses will so much the more resemble the Newtonian field as  $\lambda\phi$  is smaller in comparison with  $4\pi\kappa\rho$ .

A universe so constituted would have, with respect to its gravitational field, no centre. A decrease of density in spatial infinity would not have to be assumed, but both the mean potential and mean density would remain constant to infinity. The conflict with statistical mechanics which we found in the case of the Newtonian theory is not repeated. With a definite but extremely small density, matter is in equilibrium, without any internal material forces (pressures) being required to maintain equilibrium.

## § 2. The Boundary Conditions According to the General Theory of Relativity

In the present paragraph I shall conduct the reader over the road that I have myself travelled, rather a rough and winding road, because otherwise I cannot hope that he will

## 180 COSMOLOGICAL CONSIDERATIONS

take much interest in the result at the end of the journey. The conclusion I shall arrive at is that the field equations of gravitation which I have championed hitherto still need a slight modification, so that on the basis of the general theory of relativity those fundamental difficulties may be avoided which have been set forth in § 1 as confronting the Newtonian theory. This modification corresponds perfectly to the transition from Poisson's equation (1) to equation (2) of § 1. We finally infer that boundary conditions in spatial infinity fall away altogether, because the universal continuum in respect of its spatial dimensions is to be viewed as a self-contained continuum of finite spatial (three-dimensional) volume.

The opinion which I entertained until recently, as to the limiting conditions to be laid down in spatial infinity, took its stand on the following considerations. In a consistent theory of relativity there can be no inertia *relatively to "space,"* but only an inertia of masses *relatively to one another*. If, therefore, I have a mass at a sufficient distance from all other masses in the universe, its inertia must fall to zero. We will try to formulate this condition mathematically.

According to the general theory of relativity the negative momentum is given by the first three components, the energy by the last component of the covariant tensor multiplied by  $\sqrt{-g}$

$$m\sqrt{-g} \quad g_{\mu\alpha} \frac{dx_\alpha}{ds} \quad . \quad . \quad . \quad . \quad (4)$$

where, as always, we set

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu \quad . \quad . \quad . \quad . \quad (5)$$

In the particularly perspicuous case of the possibility of choosing the system of co-ordinates so that the gravitational field at every point is spatially isotropic, we have more simply

$$ds^2 = -A(dx_1^2 + dx_2^2 + dx_3^2) + Bdx_4^2.$$

If, moreover, at the same time

$$\sqrt{-g} = 1 = \sqrt{A^3B}$$

we obtain from (4), to a first approximation for small velocities,

$$m \frac{A}{\sqrt{B}} \frac{dx_1}{dx_4}, m \frac{A}{\sqrt{B}} \frac{dx_2}{dx_4}, m \frac{A}{\sqrt{B}} \frac{dx_3}{dx_4}$$

for the components of momentum, and for the energy (in the static case)

$$m\sqrt{B}.$$

From the expressions for the momentum, it follows that  $m\frac{A}{\sqrt{B}}$  plays the part of the rest mass. As  $m$  is a constant peculiar to the point of mass, independently of its position, this expression, if we retain the condition  $\sqrt{g} = 1$  at spatial infinity, can vanish only when  $A$  diminishes to zero, while  $B$  increases to infinity. It seems, therefore, that such a degeneration of the co-efficients  $g_{\mu\nu}$  is required by the postulate of relativity of all inertia. This requirement implies that the potential energy  $m\sqrt{B}$  becomes infinitely great at infinity. Thus a point of mass can never leave the system; and a more detailed investigation shows that the same thing applies to light-rays. A system of the universe with such behaviour of the gravitational potentials at infinity would not therefore run the risk of wasting away which was mooted just now in connexion with the Newtonian theory.

I wish to point out that the simplifying assumptions as to the gravitational potentials on which this reasoning is based, have been introduced merely for the sake of lucidity. It is possible to find general formulations for the behaviour of the  $g_{\mu\nu}$  at infinity which express the essentials of the question without further restrictive assumptions.

At this stage, with the kind assistance of the mathematician J. Grommer, I investigated centrally symmetrical, static gravitational fields, degenerating at infinity in the way mentioned. The gravitational potentials  $g_{\mu\nu}$  were applied, and from them the energy-tensor  $T_{\mu\nu}$  of matter was calculated on the basis of the field equations of gravitation. But here it proved that for the system of the fixed stars no boundary conditions of the kind can come into question at all, as was also rightly emphasized by the astronomer de Sitter recently.

For the contravariant energy-tensor  $T^{\mu\nu}$  of ponderable matter is given by

$$T^{\mu\nu} = \rho \frac{dx_\mu}{ds} \frac{dx_\nu}{ds},$$

where  $\rho$  is the density of matter in natural measure. With

## 182 COSMOLOGICAL CONSIDERATIONS

an appropriate choice of the system of co-ordinates the stellar velocities are very small in comparison with that of light. We may, therefore, substitute  $\sqrt{g_{44}} dx_4$  for  $ds$ . This shows us that all components of  $T^{\mu\nu}$  must be very small in comparison with the last component  $T^{44}$ . But it was quite impossible to reconcile this condition with the chosen boundary conditions. In the retrospect this result does not appear astonishing. The fact of the small velocities of the stars allows the conclusion that wherever there are fixed stars, the gravitational potential (in our case  $\sqrt{B}$ ) can never be much greater than here on earth. This follows from statistical reasoning, exactly as in the case of the Newtonian theory. At any rate, our calculations have convinced me that such conditions of degeneration for the  $g_{\mu\nu}$  in spatial infinity may not be postulated.

After the failure of this attempt, two possibilities next present themselves.

(a) We may require, as in the problem of the planets, that, with a suitable choice of the system of reference, the  $g_{\mu\nu}$  in spatial infinity approximate to the values

$$\begin{array}{cccc} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{array}$$

(b) We may refrain entirely from laying down boundary conditions for spatial infinity claiming general validity; but at the spatial limit of the domain under consideration we have to give the  $g_{\mu\nu}$  separately in each individual case, as hitherto we were accustomed to give the initial conditions for time separately.

The possibility (b) holds out no hope of solving the problem, but amounts to giving it up. This is an incontestable position, which is taken up at the present time by de Sitter.\* But I must confess that such a complete resignation in this fundamental question is for me a difficult thing. I should not make up my mind to it until every effort to make headway toward a satisfactory view had proved to be vain.

Possibility (a) is unsatisfactory in more respects than one.

\* de Sitter, Akad. van Wetensch. te Amsterdam, 8 Nov., 1916.

In the first place those boundary conditions pre-suppose a definite choice of the system of reference, which is contrary to the spirit of the relativity principle. Secondly, if we adopt this view, we fail to comply with the requirement of the relativity of inertia. For the inertia of a material point of mass  $m$  (in natural measure) depends upon the  $g_{\mu\nu}$ ; but these differ but little from their postulated values, as given above, for spatial infinity. Thus inertia would indeed be *influenced*, but would not be *conditioned* by matter (present in finite space). If only one single point of mass were present, according to this view, it would possess inertia, and in fact an inertia almost as great as when it is surrounded by the other masses of the actual universe. Finally, those statistical objections must be raised against this view which were mentioned in respect of the Newtonian theory.

From what has now been said it will be seen that I have not succeeded in formulating boundary conditions for spatial infinity. Nevertheless, there is still a possible way out, without resigning as suggested under (b). For if it were possible to regard the universe as a continuum which is *finite (closed) with respect to its spatial dimensions*, we should have no need at all of any such boundary conditions. We shall proceed to show that both the general postulate of relativity and the fact of the small stellar velocities are compatible with the hypothesis of a spatially finite universe; though certainly, in order to carry through this idea, we need a generalizing modification of the field equations of gravitation.

### § 3. The Spatially Finite Universe with a Uniform Distribution of Matter

According to the general theory of relativity the metrical character (curvature) of the four-dimensional space-time continuum is defined at every point by the matter at that point and the state of that matter. Therefore, on account of the lack of uniformity in the distribution of matter, the metrical structure of this continuum must necessarily be extremely complicated. But if we are concerned with the structure only on a large scale, we may represent matter to ourselves as being uniformly distributed over enormous spaces, so that its density of distribution is a variable function which varies



## 184 COSMOLOGICAL CONSIDERATIONS

extremely slowly. Thus our procedure will somewhat resemble that of the geodesists who, by means of an ellipsoid, approximate to the shape of the earth's surface, which on a small scale is extremely complicated.

The most important fact that we draw from experience as to the distribution of matter is that the relative velocities of the stars are very small as compared with the velocity of light. So I think that for the present we may base our reasoning upon the following approximative assumption. There is a system of reference relatively to which matter may be looked upon as being permanently at rest. With respect to this system, therefore, the contravariant energy-tensor  $T^{\mu\nu}$  of matter is, by reason of (5), of the simple form

$$\left. \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \rho \end{array} \right\} \dots \dots \dots (6)$$

The scalar  $\rho$  of the (mean) density of distribution may be *a priori* a function of the space co-ordinates. But if we assume the universe to be spatially finite, we are prompted to the hypothesis that  $\rho$  is to be independent of locality. On this hypothesis we base the following considerations.

As concerns the gravitational field, it follows from the equation of motion of the material point

$$\frac{d^2 x_\nu}{ds^2} + \{a\beta, \nu\} \frac{dx_a}{ds} \frac{dx_\beta}{ds} = 0$$

that a material point in a static gravitational field can remain at rest only when  $g_{44}$  is independent of locality. Since, further, we presuppose independence of the time co-ordinate  $x_4$  for all magnitudes, we may demand for the required solution that, for all  $x_\nu$ ,

$$g_{44} = 1 \quad \dots \dots \dots (7)$$

Further, as always with static problems, we shall have to set

$$g_{14} = g_{24} = g_{34} = 0 \quad \dots \dots \dots (8)$$

It remains now to determine those components of the gravitational potential which define the purely spatial-geometrical relations of our continuum ( $g_{11}, g_{12}, \dots g_{33}$ ). From

our assumption as to the uniformity of distribution of the masses generating the field, it follows that the curvature of the required space must be constant. With this distribution of mass, therefore, the required finite continuum of the  $x_1, x_2, x_3$ , with constant  $x_4$ , will be a spherical space.

We arrive at such a space, for example, in the following way. We start from a Euclidean space of four dimensions,  $\xi_1, \xi_2, \xi_3, \xi_4$ , with a linear element  $d\sigma$ ; let, therefore,

$$d\sigma^2 = d\xi_1^2 + d\xi_2^2 + d\xi_3^2 + d\xi_4^2. \quad (9)$$

In this space we consider the hyper-surface

$$R^2 = \xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2, \quad (10)$$

where  $R$  denotes a constant. The points of this hyper-surface form a three-dimensional continuum, a spherical space of radius of curvature  $R$ .

The four-dimensional Euclidean space with which we started serves only for a convenient definition of our hyper-surface. Only those points of the hyper-surface are of interest to us which have metrical properties in agreement with those of physical space with a uniform distribution of matter. For the description of this three-dimensional continuum we may employ the co-ordinates  $\xi_1, \xi_2, \xi_3$  (the projection upon the hyper-plane  $\xi_4 = 0$ ) since, by reason of (10),  $\xi_4$  can be expressed in terms of  $\xi_1, \xi_2, \xi_3$ . Eliminating  $\xi_4$  from (9), we obtain for the linear element of the spherical space the expression

$$\left. \begin{aligned} d\sigma^2 &= \gamma_{\mu\nu} d\xi_\mu d\xi_\nu \\ \gamma_{\mu\nu} &= \delta_{\mu\nu} + \frac{\xi_\mu \xi_\nu}{R^2 - \rho^2} \end{aligned} \right\} \quad (11)$$

where  $\delta_{\mu\nu} = 1$ , if  $\mu = \nu$ ;  $\delta_{\mu\nu} = 0$ , if  $\mu \neq \nu$ , and  $\rho^2 = \xi_1^2 + \xi_2^2 + \xi_3^2$ . The co-ordinates chosen are convenient when it is a question of examining the environment of one of the two points  $\xi_1 = \xi_2 = \xi_3 = 0$ .

Now the linear element of the required four-dimensional space-time universe is also given us. For the potential  $g_{\mu\nu}$ , both indices of which differ from 4, we have to set

$$g_{\mu\nu} = - \left( \delta_{\mu\nu} + \frac{x_\mu x_\nu}{R^2 - (x_1^2 + x_2^2 + x_3^2)} \right) \quad (12)$$

## 186 COSMOLOGICAL CONSIDERATIONS

which equation, in combination with (7) and (8), perfectly defines the behaviour of measuring-rods, clocks, and light-rays.

### § 4. On an Additional Term for the Field Equations of Gravitation

My proposed field equations of gravitation for any chosen system of co-ordinates run as follows:—

$$\left. \begin{aligned} G_{\mu\nu} &= -\kappa(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T), \\ G_{\mu\nu} &= -\frac{\delta}{\delta x_a}\{\mu\nu, \alpha\} + \{\mu\alpha, \beta\}\{\nu\beta, \alpha\} \\ &\quad + \frac{\partial^2 \log \sqrt{-g}}{\partial x_\mu \partial x_\nu} - \{\mu\nu, \alpha\} \frac{\partial \log \sqrt{-g}}{\partial x_\alpha} \end{aligned} \right\} \quad (13)$$

The system of equations (13) is by no means satisfied when we insert for the  $g_{\mu\nu}$  the values given in (7), (8), and (12), and for the (contravariant) energy-tensor of matter the values indicated in (6). It will be shown in the next paragraph how this calculation may conveniently be made. So that, if it were certain that the field equations (13) which I have hitherto employed were the only ones compatible with the postulate of general relativity, we should probably have to conclude that the theory of relativity does not admit the hypothesis of a spatially finite universe.

However, the system of equations (14) allows a readily suggested extension which is compatible with the relativity postulate, and is perfectly analogous to the extension of Poisson's equation given by equation (2). For on the left-hand side of field equation (13) we may add the fundamental tensor  $g_{\mu\nu}$ , multiplied by a universal constant,  $-\lambda$ , at present unknown, without destroying the general covariance. In place of field equation (13) we write

$$G_{\mu\nu} - \lambda g_{\mu\nu} = -\kappa(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T) \quad (13a)$$

This field equation, with  $\lambda$  sufficiently small, is in any case also compatible with the facts of experience derived from the solar system. It also satisfies laws of conservation of momentum and energy, because we arrive at (13a) in place of (13) by introducing into Hamilton's principle, instead of the scalar of Riemann's tensor, this scalar increased by a

universal constant; and Hamilton's principle, of course, guarantees the validity of laws of conservation. It will be shown in § 5 that field equation (13a) is compatible with our conjectures on field and matter.

### § 5. Calculation and Result

Since all points of our continuum are on an equal footing, it is sufficient to carry through the calculation for *one* point, e.g. for one of the two points with the co-ordinates

$$x_1 = x_2 = x_3 = x_4 = 0.$$

Then for the  $g_{\mu\nu}$  in (13a) we have to insert the values

$$\begin{array}{cccc} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{array}$$

wherever they appear differentiated only once or not at all. We thus obtain in the first place

$$G_{\mu\nu} = \frac{\partial}{\partial x_1}[\mu\nu, 1] + \frac{\partial}{\partial x_2}[\mu\nu, 2] + \frac{\partial}{\partial x_3}[\mu\nu, 3] + \frac{\partial^2 \log \sqrt{-g}}{\partial x_\mu \partial x_\nu}.$$

From this we readily discover, taking (7), (8), and (13) into account, that all equations (13a) are satisfied if the two relations

$$-\frac{2}{R^3} + \lambda = -\frac{\kappa\rho}{2}, \quad -\lambda = -\frac{\kappa\rho}{2},$$

or

$$\lambda = \frac{\kappa\rho}{2} = \frac{1}{R^3} \quad . \quad . \quad . \quad (14)$$

are fulfilled.

Thus the newly introduced universal constant  $\lambda$  defines both the mean density of distribution  $\rho$  which can remain in equilibrium and also the radius  $R$  and the volume  $2\pi^2 R^3$  of spherical space. The total mass  $M$  of the universe, according to our view, is finite, and is in fact

$$M = \rho \cdot 2\pi^2 R^3 = 4\pi^2 \frac{R}{\kappa} = \pi^2 \sqrt{\frac{32}{\kappa^3 \rho}} \quad . \quad . \quad (15)$$

Thus the theoretical view of the actual universe, if it is in correspondence with our reasoning, is the following. The

## 188 COSMOLOGICAL CONSIDERATIONS

curvature of space is variable in time and place, according to the distribution of matter, but we may roughly approximate to it by means of a spherical space. At any rate, this view is logically consistent, and from the standpoint of the general theory of relativity lies nearest at hand; whether, from the standpoint of present astronomical knowledge, it is tenable, will not here be discussed. In order to arrive at this consistent view, we admittedly had to introduce an extension of the field equations of gravitation which is not justified by our actual knowledge of gravitation. It is to be emphasized, however, that a positive curvature of space is given by our results, even if the supplementary term is not introduced. That term is necessary only for the purpose of making possible a quasi-static distribution of matter, as required by the fact of the small velocities of the stars.

DO GRAVITATIONAL FIELDS PLAY AN  
ESSENTIAL PART IN THE STRUC-  
TURE OF THE ELEMENTARY PAR-  
TICLES OF MATTER?

BY

A. EINSTEIN

*Translated from "Spielen Gravitationsfelder im Aufßer der  
materiellen Elementarteilchen eine wesentliche Rolle?"  
Sitzungsberichte der Preussischen Akad. d. Wissen-  
schaften, 1919.*



# DO GRAVITATIONAL FIELDS PLAY AN ESSENTIAL PART IN THE STRUCTURE OF THE ELEMENTARY PARTICLES OF MATTER?

By A. EINSTEIN

**N**EITHER the Newtonian nor the relativistic theory of gravitation has so far led to any advance in the theory of the constitution of matter. In view of this fact it will be shown in the following pages that there are reasons for thinking that the elementary formations which go to make up the atom are held together by gravitational forces.

## § 1. Defects of the Present View

Great pains have been taken to elaborate a theory which will account for the equilibrium of the electricity constituting the electron. G. Mie, in particular, has devoted deep researches to this question. His theory, which has found considerable support among theoretical physicists, is based mainly on the introduction into the energy-tensor of supplementary terms depending on the components of the electro-dynamic potential, in addition to the energy terms of the Maxwell-Lorentz theory. These new terms, which in outside space are unimportant, are nevertheless effective in the interior of the electrons in maintaining equilibrium against the electric forces of repulsion. In spite of the beauty of the formal structure of this theory, as erected by Mie, Hilbert, and Weyl, its physical results have hitherto been unsatisfactory. On the one hand the multiplicity of possibilities is discouraging, and on the other hand those additional terms have not as yet allowed themselves to be framed in such a simple form that the solution could be satisfactory.



So far the general theory of relativity has made no change in this state of the question. If we for the moment disregard the additional cosmological term, the field equations take the form

$$G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G = -\kappa T_{\mu\nu} \quad . \quad . \quad . \quad (1)$$

where  $G_{\mu\nu}$  denotes the contracted Riemann tensor of curvature,  $G$  the scalar of curvature formed by repeated contraction, and  $T_{\mu\nu}$  the energy-tensor of "matter." The assumption that the  $T_{\mu\nu}$  do *not* depend on the derivatives of the  $g_{\mu\nu}$  is in keeping with the historical development of these equations. For these quantities are, of course, the energy-components in the sense of the special theory of relativity, in which variable  $g_{\mu\nu}$  do not occur. The second term on the left-hand side of the equation is so chosen that the divergence of the left-hand side of (1) vanishes identically, so that taking the divergence of (1), we obtain the equation

$$\frac{\partial \mathfrak{T}_\mu^\sigma}{\partial x_\sigma} + \frac{1}{2}g_{\mu}^{\sigma\tau}\mathfrak{T}_{\sigma\tau} = 0 \quad . \quad . \quad . \quad (2)$$

which in the limiting case of the special theory of relativity gives the complete equations of conservation

$$\frac{\partial T_{\mu\nu}}{\partial x_\nu} = 0.$$

Therein lies the physical foundation for the second term of the left-hand side of (1). It is by no means settled *a priori* that a limiting transition of this kind has any possible meaning. For if gravitational fields do play an essential part in the structure of the particles of matter, the transition to the limiting case of constant  $g_{\mu\nu}$  would, for them, lose its justification, for indeed, with constant  $g_{\mu\nu}$  there could not be any particles of matter. So if we wish to contemplate the possibility that gravitation may take part in the structure of the fields which constitute the corpuscles, we cannot regard equation (1) as confirmed.

Placing in (1) the Maxwell-Lorentz energy-components of the electromagnetic field  $\phi_{\mu\nu}$ ,

$$T_{\mu\nu} = \frac{1}{2}g_{\mu\nu}\phi_{\sigma\tau}\phi^{\sigma\tau} - \phi_{\mu\sigma}\phi_{\nu\tau}g^{\sigma\tau}, \quad . \quad . \quad . \quad (3)$$

we obtain for (2), by taking the divergence, and after some reduction,\*

$$\phi_{\mu\sigma} \mathfrak{S}^\sigma = 0 \quad . \quad . \quad . \quad (4)$$

where, for brevity, we have set

$$\frac{\partial}{\partial x_r} (\sqrt{-g} \phi_{\mu\nu} g^{\mu\sigma} g^{\nu r}) = \frac{\partial f^{\sigma r}}{\partial x_r} = \mathfrak{S}^\sigma \quad . \quad . \quad (5)$$

In the calculation we have employed the second of Maxwell's systems of equations

$$\frac{\partial \phi_{\mu\nu}}{\partial x_\rho} + \frac{\partial \phi_{\nu\rho}}{\partial x_\mu} + \frac{\partial \phi_{\rho\mu}}{\partial x_\nu} = 0 \quad . \quad . \quad . \quad (6)$$

We see from (4) that the current-density  $\mathfrak{S}^\sigma$  must everywhere vanish. Therefore, by equation (1), we cannot arrive at a theory of the electron by restricting ourselves to the electromagnetic components of the Maxwell-Lorentz theory, as has long been known. Thus if we hold to (1) we are driven on to the path of Mie's theory.†

Not only the problem of matter, but the cosmological problem as well, leads to doubt as to equation (1). As I have shown in the previous paper, the general theory of relativity requires that the universe be spatially finite. But this view of the universe necessitated an extension of equations (1), with the introduction of a new universal constant  $\lambda$ , standing in a fixed relation to the total mass of the universe (or, respectively, to the equilibrium density of matter). This is gravely detrimental to the formal beauty of the theory.

## § 2. The Field Equations Freed of Scalars

The difficulties set forth above are removed by setting in place of field equations (1) the field equations

$$G_{\mu\nu} - \frac{1}{2} g_{\mu\nu} G = -\kappa T_{\mu\nu} \quad . \quad . \quad . \quad (1a)$$

where  $T_{\mu\nu}$  denotes the energy-tensor of the electromagnetic field given by (3).

The formal justification for the factor  $-\frac{1}{2}$  in the second

\* Cf. e.g. A. Einstein, Sitzungsber. d. Preuss. Akad. d. Wiss., 1916, pp. 187, 188.

† Cf. D. Hilbert, Göttinger Nachr., 20 Nov., 1915.

term of this equation lies in its causing the scalar of the left-hand side,

$$g^{\mu\nu}(G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G),$$

to vanish identically, as the scalar  $g^{\mu\nu}T_{\mu\nu}$  of the right-hand side does by reason of (3). If we had reasoned on the basis of equations (1) instead of (1a), we should, on the contrary, have obtained the condition  $G = 0$ , which would have to hold good everywhere for the  $g_{\mu\nu}$ , independently of the electric field. It is clear that the system of equations [(1a), (3)] is a consequence of the system [(1), (3)], but not conversely.

We might at first sight feel doubtful whether (1a) together with (6) sufficiently define the entire field. In a generally relativistic theory we need  $n - 4$  differential equations, independent of one another, for the definition of  $n$  independent variables, since in the solution, on account of the liberty of choice of the co-ordinates, four quite arbitrary functions of all co-ordinates must naturally occur. Thus to define the sixteen independent quantities  $g_{\mu\nu}$  and  $\phi_{\mu\nu}$  we require twelve equations, all independent of one another. But as it happens, nine of the equations (1a), and three of the equations (6) are independent of one another.

Forming the divergence of (1a), and taking into account that the divergence of  $G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G$  vanishes, we obtain

$$\phi_{\sigma\alpha}J^{\alpha} + \frac{1}{4\kappa} \frac{\partial G}{\partial x_{\sigma}} = 0 \quad . \quad . \quad . \quad (4a)$$

From this we recognize first of all that the scalar of curvature  $G$  in the four-dimensional domains in which the density of electricity vanishes, is constant. If we assume that all these parts of space are connected, and therefore that the density of electricity differs from zero only in separate "world-threads," then the scalar of curvature, everywhere outside these world-threads, possesses a constant value  $G_0$ . But equation (4a) also allows an important conclusion as to the behaviour of  $G$  within the domains having a density of electricity other than zero. If, as is customary, we regard electricity as a moving density of charge, by setting

$$J^{\sigma} = \frac{3^{\sigma}}{\sqrt{-g}} = \rho \frac{dx_{\sigma}}{ds}, \quad . \quad . \quad . \quad (7)$$

we obtain from (4a) by inner multiplication by  $J^\sigma$ , on account of the antisymmetry of  $\phi_{\mu\nu}$ , the relation

$$\frac{\partial G}{\partial x_\sigma} \frac{dx_\sigma}{ds} = 0 \quad . \quad . \quad . \quad . \quad (8)$$

Thus the scalar of curvature is constant on every world-line of the motion of electricity. Equation (4a) can be interpreted in a graphic manner by the statement: The scalar of curvature plays the part of a negative pressure which, outside of the electric corpuscles, has a constant value  $G_0$ . In the interior of every corpuscle there subsists a negative pressure (positive  $G - G_0$ ) the fall of which maintains the electrodynamic force in equilibrium. The minimum of pressure, or, respectively, the maximum of the scalar of curvature, does not change with time in the interior of the corpuscle.

We now write the field equations (1a) in the form

$$(G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G) + \frac{1}{2}g_{\mu\nu}G_0 = -\kappa\left(T_{\mu\nu} + \frac{1}{4\kappa}g_{\mu\nu}(G - G_0)\right) \quad (9)$$

On the other hand, we transform the equations supplied with the cosmological term as already given

$$G_{\mu\nu} - \lambda g_{\mu\nu} = -\kappa(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T).$$

Subtracting the scalar equation multiplied by  $\frac{1}{2}$ , we next obtain

$$(G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G) + g_{\mu\nu}\lambda = -\kappa T_{\mu\nu}.$$

Now in regions where only electrical and gravitational fields are present, the right-hand side of this equation vanishes. For such regions we obtain, by forming the scalar,

$$-G + 4\lambda = 0.$$

In such regions, therefore, the scalar of curvature is constant, so that  $\lambda$  may be replaced by  $\frac{1}{4}G_0$ . Thus we may write the earlier field equation (1) in the form

$$G_{\mu\nu} - \frac{1}{2}g_{\mu\nu}G + \frac{1}{2}g_{\mu\nu}G_0 = -\kappa T_{\mu\nu} \quad . \quad . \quad (10)$$

Comparing (9) with (10), we see that there is no difference between the new field equations and the earlier ones, except that instead of  $T_{\mu\nu}$  as tensor of "gravitating mass" there now

occurs  $T_{\mu\nu} + \frac{1}{4\kappa} g_{\mu\nu}(G - G_0)$  which is independent of the scalar of curvature. But the new formulation has this great advantage, that the quantity  $\lambda$  appears in the fundamental equations as a constant of integration, and no longer as a universal constant peculiar to the fundamental law.

### § 3. On the Cosmological Question

The last result already permits the surmise that with our new formulation the universe may be regarded as spatially finite, without any necessity for an additional hypothesis. As in the preceding paper I shall again show that with a uniform distribution of matter, a spherical world is compatible with the equations.

In the first place we set

$$ds^2 = -\gamma_{ik} dx_i dx_k + dx_4^2 \quad (i, k = 1, 2, 3) \quad (11)$$

Then if  $P_{ik}$  and  $P$  are, respectively, the curvature tensor of the second rank and the curvature scalar in three-dimensional space, we have

$$\begin{aligned} G_{ik} &= P_{ik} \quad (i, k = 1, 2, 3) \\ G_{i4} &= G_{4i} = G_{44} = 0 \\ G &= -P \\ -g &= \gamma. \end{aligned}$$

It therefore follows for our case that

$$\begin{aligned} G_{ik} - \frac{1}{2} g_{ik} G &= P_{ik} - \frac{1}{2} \gamma_{ik} P \quad (i, k = 1, 2, 3) \\ G_{44} - \frac{1}{2} g_{44} G &= \frac{1}{2} P. \end{aligned}$$

We pursue our reflexions, from this point on, in two ways. Firstly, with the support of equation (1a). Here  $T_{\mu\nu}$  denotes the energy-tensor of the electro-magnetic field, arising from the electrical particles constituting matter. For this field we have everywhere

$$\mathfrak{F}_1^1 + \mathfrak{F}_2^2 + \mathfrak{F}_3^3 + \mathfrak{F}_4^4 = 0.$$

The individual  $\mathfrak{F}_\mu^\nu$  are quantities which vary rapidly with position; but for our purpose we no doubt may replace them by their mean values. We therefore have to choose

$$\left. \begin{aligned} \mathfrak{F}_1^1 &= \mathfrak{F}_2^2 = \mathfrak{F}_3^3 = -\frac{1}{3} \mathfrak{F}_4^4 = \text{const.} \\ \mathfrak{F}_\mu^\nu &= 0 \quad (\text{for } \mu \neq \nu), \end{aligned} \right\} \quad (12)$$

and therefore

$$T_{ik} = \frac{1}{3} \frac{\mathfrak{T}_i^4}{\sqrt{\gamma}} \gamma_{ik}, \quad T_{44} = \frac{\mathfrak{T}_4^4}{\sqrt{\gamma}}.$$

In consideration of what has been shown hitherto, we obtain in place of (1a)

$$P_{ik} - \frac{1}{2} \gamma_{ik} P = - \frac{1}{3} \gamma_{ik} \frac{\kappa \mathfrak{T}_i^4}{\sqrt{\gamma}}. \quad (13)$$

$$\frac{1}{2} P = - \frac{\kappa \mathfrak{T}_4^4}{\sqrt{\gamma}}. \quad (14)$$

The scalar of equation (13) agrees with (14). It is on this account that our fundamental equations permit the idea of a spherical universe. For from (13) and (14) follows

$$P_{ik} + \frac{4}{3} \frac{\kappa \mathfrak{T}_i^4}{\sqrt{\gamma}} \gamma_{ik} = 0. \quad (15)$$

and it is known\* that this system is satisfied by a (three-dimensional) spherical universe.

But we may also base our reflexions on the equations (9). On the right-hand side of (9) stand those terms which, from the phenomenological point of view, are to be replaced by the energy-tensor of matter; that is, they are to be replaced by

$$\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \rho \end{array}$$

where  $\rho$  denotes the mean density of matter assumed to be at rest. We thus obtain the equations

$$P_{ik} - \frac{1}{2} \gamma_{ik} P - \frac{1}{2} \gamma_{ik} G_0 = 0. \quad (16)$$

$$\frac{1}{2} P + \frac{1}{2} G_0 = - \kappa \rho. \quad (17)$$

From the scalar of equation (16) and from (17) we obtain

$$G_0 = - \frac{2}{3} P = 2\kappa\rho, \quad (18)$$

and consequently from (16)

$$P_{ik} - \kappa\rho\gamma_{ik} = 0. \quad (19)$$

\* Cf. H. Weyl, "Raum, Zeit, Materie," § 88.

which equation, with the exception of the expression for the co-efficient, agrees with (15). By comparison we obtain

$$\mathfrak{E}_i^4 = \frac{1}{2} \rho \sqrt{\gamma}. \quad . \quad . \quad . \quad . \quad (20)$$

This equation signifies that of the energy constituting matter three-quarters is to be ascribed to the electromagnetic field, and one-quarter to the gravitational field.

#### § 4. Concluding Remarks

The above reflexions show the possibility of a theoretical construction of matter out of gravitational field and electromagnetic field alone, without the introduction of hypothetical supplementary terms on the lines of Mie's theory. This possibility appears particularly promising in that it frees us from the necessity of introducing a special constant  $\lambda$  for the solution of the cosmological problem. On the other hand, there is a peculiar difficulty. For, if we specialize (1) for the spherically symmetrical static case we obtain one equation too few for defining the  $g_{\mu\nu}$  and  $\phi_{\mu\nu}$ , with the result that *any spherically symmetrical distribution* of electricity appears capable of remaining in equilibrium. Thus the problem of the constitution of the elementary quanta cannot yet be solved on the immediate basis of the given field equations.

# GRAVITATION AND ELECTRICITY

BY

H. WEYL

*Translated from "Gravitation und Elektrizität," Sitzungs-  
berichte der Preussischen Akad. d. Wissenschaften, 1918.*





# GRAVITATION AND ELECTRICITY\*

By H. WEYL

ACCORDING to Riemann,<sup>†</sup> geometry is based upon the following two facts:—

1. *Space is a Three-dimensional Continuum.*—The manifold of its points may therefore be consistently represented by the values of three co-ordinates  $x_1, x_2, x_3$ .

2. (*Pythagorean Theorem*).—The square of the distance  $ds$  between two infinitely proximate points

$P = (x_1, x_2, x_3)$  and  $P' = (x_1 + dx_1, x_2 + dx_2, x_3 + dx_3)$  (1) (any co-ordinates being employed) is a quadratic form of the relative co-ordinates  $dx_\mu$ :—

$$ds^2 = \sum_{\mu\nu} g_{\mu\nu} dx_\mu dx_\nu, \quad (g_{\mu\nu} = g_{\nu\mu}) \quad . \quad . \quad (2)$$

The second of these facts may be briefly stated by saying that space is a *metrical* continuum. In complete accord with the spirit of the physics of immediate action we assume the Pythagorean theorem to be strictly valid only in the limit when the distances are infinitely small.

The special theory of relativity led to the discovery that *time* is associated as a fourth co-ordinate ( $x_4$ ) on an equal footing with the three co-ordinates of space, and that the scene of material events, *the world*, is therefore a *four-dimensional, metrical continuum*. And so the quadratic form (2), which defines the metrical properties of the world, is not necessarily positive as in the case of the geometry of three-dimensional space, but has the index of inertia 3.<sup>‡</sup> Riemann

\* The footnotes in square brackets are later additions by the author.

+ Math. Werke (2nd ed., Leipzig, 1892), No. XII, p. 282.

‡ That is to say that if the co-ordinates are chosen so that at one particular point of the continuum  $ds^2 = \pm dx_1^2 \pm dx_2^2 \pm dx_3^2 \pm dx_4^2$ , then in every case three of the signs will be + and one — (TRANS.).

## 202 GRAVITATION AND ELECTRICITY

himself did not fail to point out that this quadratic form was to be regarded as a physical reality, since it reveals itself, e.g. in centrifugal forces, as the origin of real effects upon matter, and that matter therefore presumably reacts upon it. Until then all geometricians and philosophers had looked upon the metrical properties of space as pertaining to space itself, independently of the matter which it contained. It is upon this idea, which it was quite impossible for Riemann in his day to carry through, that Einstein in our own time, independently of Riemann, has raised the imposing edifice of his general theory of relativity. According to Einstein the phenomena of *gravitation* must also be placed to the account of geometry, and the laws by which matter affects measurements are no other than the laws of gravitation: the  $g_{\mu\nu}$  in (2) form the components of the gravitational potential. While the gravitational potential thus consists of an invariant *quadratic* differential form, *electromagnetic phenomena* are governed by a four-potential of which the components  $\phi_\mu$  together compose an invariant *linear* differential form  $\Sigma \phi_\mu dx_\mu$ . But so far the two classes of phenomena, gravitation and electricity, stand side by side, the one separate from the other.

The later work of Levi-Civita,\* Hessenberg,† and the author‡ shows quite plainly that the fundamental conception on which the development of Riemann's geometry must be based if it is to be in agreement with nature, is that of the infinitesimal parallel displacement of a vector. If P and P\* are any two points connected by a curve, a given vector at P can be moved parallel to itself along this curve from P to P\*. But, generally speaking, this conveyance of a vector from P to P\* is not integrable, that is to say, the vector at P\* at which we arrive depends upon the path along which the displacement travels. It is only in Euclidean "gravitationless" geometry that integrability obtains. The Riemannian geometry referred to above still contains a residual element of finite geometry—without any substantial reason, as far as I can see.

\* "Nozione di parallelismo . . .", Rend. del Circ. Matem. di Palermo, Vol. 42 (1917).

† "Vektorielle Begründung der Differentialgeometrie," Math. Ann., Vol. 78 (1917).

‡ "Space, Time, and Matter" (1st ed., Berlin, 1918), § 14.

It seems to be due to the accidental origin of this geometry in the theory of surfaces. The quadratic form (2) enables us to compare, with respect to their length, not only two vectors at the same point, but also the vectors at any two points. *But a truly infinitesimal geometry must recognize only the principle of the transference of a length from one point to another point infinitely near to the first.* This forbids us to assume that the problem of the transference of length from one point to another at a finite distance is integrable, more particularly as the problem of the transference of direction has proved to be non-integrable. Such an assumption being recognized as false, a geometry comes into being, which, when applied to the world, explains in a surprising manner *not only the phenomena of gravitation, but also those of the electromagnetic field.* According to the theory which now takes shape, both classes of phenomena spring from the same source, and in fact *we cannot in general make any arbitrary separation of electricity from gravitation.* In this theory *all physical quantities have a meaning in world geometry.* In particular the quantities denoting physical effects appear at once as pure numbers. The theory leads to a world-law which in its essentials is defined without ambiguity. It even permits us in a certain sense to comprehend why the world has four dimensions. I shall now first of all give a sketch of the structure of the amended geometry of Riemann without any thought of its physical interpretation. Its application to physics will then follow of its own accord.

In a given system of co-ordinates the relative co-ordinates  $dx_\mu$  of a point P infinitely near to P—see (1)—are the components of the infinitesimal displacement PP'. The transition from one system of co-ordinates to another is expressed by definite formulæ of transformation,

$$x_\mu = x_\mu(x_1^*, x_2^* \dots x_n^*) \quad \mu = 1, 2, \dots n,$$

which determine the connexion between the co-ordinates of the same point in the two systems. Then between the components  $dx_\mu$  and the components  $dx_\mu^*$  of the same infinitesimal displacement of the point P we have the linear formulæ of transformation

$$dx_\mu = \sum_\nu a_{\mu\nu} dx_\nu^* \quad . \quad . \quad . \quad (3)$$

## 204 GRAVITATION AND ELECTRICITY

in which  $\alpha_{\mu\nu}$  are the values of the derivatives  $\frac{\partial x_\mu}{\partial x_\nu^*}$  at the point

P. A contravariant vector  $x$  at the point P referred to either system of co-ordinates has  $n$  known numbers  $\xi^\mu$  for its components, which in the transition to another system are transformed in exactly the same way (3) as the components of an infinitesimal displacement. I denote the totality of vectors at the point P as the vector-space at P. It is, firstly, linear or affine, i.e. by multiplication of a vector at P by a number, and by addition of two such vectors, there always arises a vector at P; and, secondly, it is metrical, i.e. by the symmetrical bilinear form belonging to (2) a scalar product

$$x \cdot y = y \cdot x = \sum_{\mu\nu} g_{\mu\nu} \xi^\mu \eta^\nu$$

is invariantly assigned to each pair of vectors  $x$  and  $y$  with components  $\xi^\mu, \eta^\mu$ . We take it, however, that this form is determined only as far as to a positive factor of proportionality, which remains arbitrary. If the manifold of points of space is represented by co-ordinates  $x_\mu$ , the  $g_{\mu\nu}$  are determined by the metrical properties at the point P only to the extent of their proportionality. In the physical sense, too, it is only the ratios of the  $g_{\mu\nu}$  that has an immediate tangible meaning. For the equation

$$\sum_{\mu\nu} g_{\mu\nu} dx_\mu dx_\nu = 0$$

is satisfied, when P is a given origin, by those infinitely proximate world-points which are reached by a light signal emitted at P. For the purpose of analytical presentation we have firstly to choose a definite system of co-ordinates, and secondly at each point P to determine the arbitrary factor of proportionality with which the  $g_{\mu\nu}$  are endowed. Accordingly the formulæ which emerge must possess a double property of invariance: they must be invariant with respect to any continuous transformations of co-ordinates, and they must remain unaltered if  $\lambda g_{\mu\nu}$ , where  $\lambda$  is an arbitrary continuous function of position, is substituted for the  $g_{\mu\nu}$ . The supervision of this second property of invariance is characteristic of our theory.

If P, P\* are any two points, and if to each vector  $x$  at P a vector  $x^*$  at P\* is assigned in such a way that in general  $\alpha x$

becomes  $\alpha x^*$ , and  $x + y$  becomes  $x^* + y^*$  ( $\alpha$  being any assigned number), and the vector zero at  $P$  is the only one to which the vector zero at  $P^*$  corresponds, we then have made an affine or linear replica of the vector-space at  $P$  on the vector-space at  $P^*$ . This replica has a particularly close resemblance when the scalar product of the vectors  $x^*$ ,  $y^*$  at  $P^*$  is proportional to that of  $x$  and  $y$  at  $P$  for all pairs of vectors  $x$ ,  $y$ . (In our view it is only this idea of a similar replica that has an objective sense; the previous theory permitted the more definite conception of a congruent replica.) The meaning of the parallel displacement of a vector at the point  $P$  to a neighbouring point  $P'$  is settled by the two axiomatic postulates.

1. By the parallel displacement of the vectors at the point  $P$  to the neighbouring point  $P'$  a similar image of the vector-space at  $P$  is made upon the vector-space at  $P'$ .

2. If  $P_1, P_2$  are two points in the neighbourhood of  $P$ , and the infinitesimal vector  $PP_2$  at  $P$  is transformed into  $P_1P_{12}$  by a parallel displacement to the point  $P_1$ , while  $PP_1$  at  $P$  is transformed into  $P_2P_{21}$  by parallel displacement to  $P_2$ , then  $P_{12}, P_{21}$  coincide, i.e. infinitesimal parallel displacements are commutative.

That part of postulate 1 which says that the parallel displacement is an affine transposition of the vector-space from  $P$  to  $P'$ , is expressed analytically as follows: the vector  $\xi^\mu$  at  $P = (x_1, x_2, \dots, x_n)$  is by displacement transformed into a vector  $\xi^\mu + d\xi^\mu$  at  $P' = (x_1 + dx_1, x_2 + dx_2, \dots, x_n + dx_n)$  the components of which are in a linear relation to  $\xi^\mu$ ,—

$$d\xi^\mu = - \sum_{\nu} d\gamma_{\nu}^{\mu} \xi^{\nu} \quad . \quad . \quad . \quad (4)$$

The second postulate teaches that the  $d\gamma_{\nu}^{\mu}$  are linear differential forms

$$d\gamma_{\nu}^{\mu} = \sum_{\rho} \Gamma_{\nu\rho}^{\mu} dx_{\rho},$$

the coefficients of which possess the symmetrical property

$$\Gamma_{\nu\rho}^{\mu} = \Gamma_{\rho\nu}^{\mu} \quad . \quad . \quad . \quad (5)$$

If two vectors  $\xi^\mu, \eta^\mu$  at  $P$  are transformed by parallel displacement at  $P'$  into  $\xi^\mu + d\xi^\mu, \eta^\mu + d\eta^\mu$ , then the postulate

## 206 GRAVITATION AND ELECTRICITY

of similarity stated under 1 above, which goes beyond affinity, tells us that

$$\sum_{\mu\nu} (g_{\mu\nu} + dg_{\mu\nu})(\xi^\mu + d\xi^\mu)(\eta^\nu + d\eta^\nu)$$

must be proportional to

$$\sum_{\mu\nu} g_{\mu\nu} \xi^\mu \eta^\nu.$$

If we call the factor of proportionality, which differs infinitesimally from 1,  $1 + d\phi$ , and define the reduction of an index in the usual way by the formula

$$a_\mu = \sum_\nu g_{\mu\nu} a^\nu,$$

we obtain

$$dg_{\mu\nu} - (d\gamma_{\nu\mu} + d\gamma_{\mu\nu}) = g_{\mu\nu} d\phi \quad . \quad . \quad (6)$$

From this it follows that  $d\phi$  is a linear differential form

$$d\phi = \sum_\mu \phi_\mu dx_\mu \quad . \quad . \quad . \quad (7)$$

If this is known, the equation (6) or

$$\Gamma_{\mu, \nu\rho} + \Gamma_{\nu, \mu\rho} = \frac{\partial g_{\mu\nu}}{\partial x_\rho} - g_{\mu\nu} \phi_\rho,$$

together with the condition for symmetry (5), gives unequivocally the quantities  $\Gamma$ . *The internal metrical connexion of space thus depends on a linear form (7) besides the quadratic form (2)—which is determined except as to an arbitrary factor of proportionality.\** If we substitute  $\lambda g_{\mu\nu}$  for  $g_{\mu\nu}$  with-

\* [I have now modified this structure in the following points (cf. the final presentation in ed. 4 of "Raum, Zeit, Materie," 1921, §§ 18, 18). (a) In place of postulates 1 and 2, which the parallel displacement has to fulfil, there is now one postulate: Let there be a system of co-ordinates at the point P, by the employment of which the components of every vector at P are not altered by parallel displacement to any point in infinite proximity to P. This postulate characterizes the essence of the parallel displacement as that of a transposition, concerning which it may be correctly asserted that it leaves the vectors "unaltered." (b) To the metrics at the single point P, according to which there is attached to every vector  $x = \xi^\mu$  at P a tract of such a kind that two vectors define the same tract when, and only when, they possess the same measure-number  $l = \sum g_{\mu\nu} \xi^\mu \xi^\nu$ , there must now be added the metrical connexion of P with the points in its neighbourhood: by congruent transposition to the infinitely near point P' a tract at P passes over into a definite tract at P'. If we make a requirement of this concept of congruent transposition of tracts analogous to that which has just been postulated, under (a), of the concept of parallel displacement of vectors, we see that this process (in which the measure-number  $l$  of the tract is increased by  $dl$ ) is expressed in the equations

$$dl = l d\phi; \quad d\phi = \sum \phi_\mu dx_\mu.$$

out changing the system of co-ordinates, the quantities  $d\gamma_\nu^\mu$  do not change,  $d\gamma_{\mu\nu}$  assumes the factor  $\lambda$ , and  $dg_{\mu\nu}$  becomes  $\lambda dg_{\mu\nu} + g_{\mu\nu}d\lambda$ . Equation (6) then shows that  $d\phi$  becomes

$$d\phi + \frac{d\lambda}{\lambda} = d\phi + d(\log \lambda).$$

What remains undetermined, therefore, in the linear form  $\Sigma \phi_\mu dx_\mu$  is not a factor of proportionality which would have to be settled by an arbitrary choice of a unit of measurement, but, rather, the arbitrary element inherent in it consists in an additive total differential. For the analytical representation of geometry the forms

$$g_{\mu\nu} dx_\mu dx_\nu, \quad \phi_\mu dx_\mu \quad . \quad . \quad . \quad (8)$$

are on an equal footing with

$$\lambda \cdot g_{\mu\nu} dx_\mu dx_\nu \text{ and } \phi_\mu dx_\mu + d(\log \lambda) \quad . \quad . \quad (9)$$

where  $\lambda$  is any positive function of position. Hence there is invariant significance in the anti-symmetrical tensor with the components

$$F_{\mu\nu} = \frac{\partial \phi_\mu}{\partial x_\nu} - \frac{\partial \phi_\nu}{\partial x_\mu} \quad . \quad . \quad . \quad (10)$$

i.e. the form

$$F_{\mu\nu} = dx_\mu \delta x_\nu - dx_\nu \delta x_\mu$$

which depends bilinearly on two arbitrary displacements  $dx$  and  $\delta x$  at the point P—or, rather, depends linearly on the surface element with the components  $\Delta x_{\mu\nu} = dx_\mu \delta x_\nu - dx_\nu \delta x_\mu$  which is defined by these two displacements. The special case of the theory as hitherto developed, in which the arbitrarily chosen unit of length at the origin allows itself to be transferred by parallel displacement to all points of space in a manner which is independent of the path traversed—this special case occurs when the  $g_{\mu\nu}$  can be absolutely determined in such a way that the  $\phi_\mu$  vanish. The  $\Gamma_{\nu\rho}^\mu$  are

In these circumstances the metrics and the metrical connexion determine the "affine" connexion (parallel displacement) without ambiguity—and indeed, according to my present view of the problem of space this is the most fundamental fact of geometry—whereas according to the presentation given in the text it is the linear form  $d\phi$  that remains arbitrary in the given metrics at the parallel displacement.]



## 208 GRAVITATION AND ELECTRICITY

then nothing else than the Christoffel three-indices symbols. The necessary and sufficient invariant condition for the occurrence of this case consists in the identical vanishing of the tensor  $F_{\mu\nu}$ .

This naturally suggests interpreting  $\phi_\mu$  in world-geometry as the four-potential, and the tensor  $F$  consequently as electromagnetic field. For the absence of an electromagnetic field is the necessary condition for the validity of Einstein's theory, which, up to the present, accounts for the phenomena of gravitation only. If this view is accepted, it will be seen that the electric quantities are of such a nature that their characterization by numbers in a definite system of co-ordinates does not depend on the arbitrary choice of a unit of measurement. In fact, in the question of the unit of measurement and of dimension there must be a new orientation of the theory. Hitherto a quantity has been spoken of as, e.g., a tensor of the second rank, if a single value of the quantity determines a matrix of numbers  $a_{\mu\nu}$  in each system of co-ordinates after an arbitrary unit of measurement has been selected, these numbers forming the coefficients of an invariant bilinear form of two arbitrary, infinitesimal displacements

$$a_{\mu\nu}dx_\mu dx_\nu \quad . \quad . \quad . \quad . \quad . \quad (11)$$

But here we speak of a tensor, if, with a system of co-ordinates taken as a base, and after definite selection of the factor of proportionality contained in the  $g_{\mu\nu}$ , the components  $a_{\mu\nu}$  are determined without ambiguity and in such a way that on transforming the co-ordinates the form (11) remains invariant, but on replacing  $g_{\mu\nu}$  by  $\lambda g_{\mu\nu}$  the  $a_{\mu\nu}$  become  $\lambda^e a_{\mu\nu}$ . We then say that the tensor has the weight  $e$ , or, ascribing to the linear element  $ds$  the dimension "length =  $l$ ," that it is of dimension  $l^{2e}$ . Only those tensors of weight 0 are absolutely invariant. The field tensor with the components  $F_{\mu\nu}$  is of this kind. By (10) it satisfies the first system of the Maxwell equations

$$\frac{\partial F_{\nu\rho}}{\partial x_\mu} + \frac{\partial F_{\rho\mu}}{\partial x_\nu} + \frac{\partial F_{\mu\nu}}{\partial x_\rho} = 0.$$

When once the idea of parallel displacement is clear, geometry and the tensor calculus can be established without difficulty.

(a) *Geodesic Lines*.—Given a point P and at that point a vector, the geodesic line from P in the direction of this vector is given by continuously moving the vector parallel to itself in its own direction. Employing a suitable parameter  $\tau$  the differential equation of the geodesic line is

$$\frac{d^2 x_\mu}{d\tau^2} + \Gamma_{\nu\rho}^\mu \frac{dx_\nu}{d\tau} \frac{dx_\rho}{d\tau} = 0.$$

(Of course it cannot be characterized as the line of smallest length, because the notion of curve-length has no meaning.)

(b) *Tensor Calculus*.—To deduce, for example, a tensor field of rank 2 by differentiation from a covariant tensor field of rank 1 and weight 0 with components  $f_\mu$ , we call in the help of an arbitrary vector  $\xi^\mu$  at the point P, form the invariant  $f_\mu \xi^\mu$  and its infinitely small alteration on transition from the point P with the co-ordinates  $x_\mu$  to the neighbouring point P' with the co-ordinates  $x_\mu + dx_\mu$  by shifting the vector along a parallel to itself during this transition. For this alteration we have

$$\frac{\partial f_\mu}{\partial x_\nu} \xi^\mu dx_\nu + f_\rho d\xi^\rho = \left( \frac{\partial f_\mu}{\partial x_\nu} - \Gamma_{\mu\nu}^\rho f_\rho \right) \xi^\mu dx_\nu.$$

The quantities in brackets on the right are therefore the components of a tensor field of rank 2 and weight 0, which is formed from the field  $f$  in a perfectly invariant manner.

(c) *Curvature*.—To construct the analogue to Riemann's tensor of curvature, let us begin with the figure employed above, of an infinitely small parallelogram, consisting of the points P,  $P_1$ ,  $P_2$ , and  $P_{12} = P_{21}$ .\* If we displace a vector  $x = \xi^\mu$  at P parallel to itself, to  $P_1$  and from there to  $P_{12}$ , and a second time first to  $P_2$  and thence to  $P_{21}$ , then, since  $P_{12}$  and  $P_{21}$  coincide, there is a meaning in forming the difference  $\Delta x$  of the two vectors obtained at this point. For their components we have

$$\Delta \xi^\mu = \Delta R_{\nu\rho}^\mu \xi^\nu \quad . \quad . \quad . \quad (12)$$

where the  $\Delta R_{\nu\rho}^\mu$  are independent of the displaced vector  $x$ , but

\* [Here it is not essential that opposite sides of the infinitely small "parallelogram" are produced by parallel displacement one from the other; we are concerned only with the coincidence of the points  $P_{12}$  and  $P_{21}$ .]

## 210 GRAVITATION AND ELECTRICITY

on the other hand depend linearly on the surface-element defined by the two displacements  $PP_1 = dx_\mu$ ,  $PP_2 = \delta x_\mu$ . Thus

$$\Delta R_\nu^\mu = R_{\nu\rho\sigma}^\mu dx_\rho \delta x_\sigma = \frac{1}{2} R_{\nu\rho\sigma}^\mu \Delta x_{\rho\sigma}.$$

The components of curvature  $R_{\nu\rho\sigma}^\mu$ , depending solely on the place P, possess the two properties of symmetry that (1) they change sign on the interchange of the last two indices  $\rho$  and  $\sigma$ , and (2), if we perform the three cyclic interchanges  $\nu\rho\sigma$ , and add up the appropriate components, the result is 0. Reducing the index  $\mu$ , we obtain at  $R_{\mu\nu\rho\sigma}$  the components of a covariant tensor of rank 4 and weight 1. Even without calculation we see that R divides in a natural, invariant manner into two parts,

$$R_{\nu\rho\sigma}^\mu = P_{\nu\rho\sigma}^\mu - \frac{1}{2} \delta_\nu^\mu F_{\rho\sigma} \quad (\delta_\nu^\mu = 1 \text{ if } \mu = \nu; = 0 \text{ if } \mu \neq \nu), \quad (13)$$

of which the first,  $P_{\nu\rho\sigma}^\mu$ , is anti-symmetrical, not only in the indices  $\rho\sigma$ , but also in  $\mu$  and  $\nu$ . Whereas the equations  $F_{\mu\nu} = 0$  characterize our space as one without an electromagnetic field, i.e. as one in which the problem of the conveyance of length is integrable, the equations  $P_{\nu\rho\sigma}^\mu = 0$  are, as (13) shows, the invariant conditions for the absence of a gravitational field, i.e. for the problem of the conveyance of direction to be integrable. The Euclidean space alone is one which at the same time is free of electricity and of gravitation.

The simplest invariant of a linear copy like (12), which to each vector  $x$  assigns a vector  $\Delta x$ , is its "spur"

$$\frac{1}{n} \Delta R_\mu^\mu.$$

For this, by (13), we obtain in the present case the form

$$- \frac{1}{2} F_{\rho\sigma} dx_\rho \delta x_\sigma$$

which we have already encountered above. The simplest invariant of a tensor like  $-\frac{1}{2} F_{\rho\sigma}$  is the "square of its magnitude"

$$L = \frac{1}{4} F_{\rho\sigma} F^{\rho\sigma} \quad . \quad . \quad . \quad (14)$$

L is evidently an invariant of weight - 2, because the tensor F has weight 0. If  $g$  is the negative determinant of the  $g_{\mu\nu}$ , and

$$d\omega = \sqrt{g} dx_0 dx_1 dx_2 dx_3 = \sqrt{g} dx$$

the volume of an infinitely small element of volume, it is known that the Maxwell theory is governed by the quantity of electrical action, which is equal to the integral  $\int L d\omega$  of this simplest invariant, extended over any chosen territory, and indeed is governed in the sense that, with any variations of the  $g_{\mu\nu}$  and  $\phi_\mu$ , which vanish at the limits of world-territory, we have

$$\delta \int L d\omega = \int (S^\mu d\phi_\mu + T^{\mu\nu} \delta g_{\mu\nu}) d\omega,$$

where

$$S^\mu = \frac{1}{\sqrt{g}} \frac{\partial(\sqrt{g} F^{\mu\nu})}{\partial x_\nu}$$

are the left-hand sides of the generalized Maxwellian equations (the right-hand sides of which are the components of the four-current), and the  $T^{\mu\nu}$  form the energy-momentum tensor of the electromagnetic field. As  $L$  is an invariant of weight  $-2$ , whereas the volume-element in  $n$ -dimensional geometry is an invariant of weight  $\frac{1}{2}n$ , the integral has significance only when the number of dimensions  $n = 4$ . Thus on our interpretation the possibility of the Maxwell theory is restricted to the case of four dimensions. In the four-dimensional world, however, the quantity of electromagnetic action becomes a pure number. Nevertheless, the magnitude of the quantity 1 cannot be ascertained in the traditional units of the c.g.s. system until a physical problem, to be tested by observation (as for example the electron), has been calculated on the basis of our theory.

Passing now from geometry to physics, we have to assume, following the precedent of Mie's theory,\* that all the laws of nature rest upon a definite integral invariant, the action-quantity

$$\int W d\omega = \int \mathfrak{B} dx, \quad \mathfrak{B} = W \sqrt{g},$$

in such a way that the real world is distinguished from all other possible four-dimensional metrical spaces by the characteristic that for it the action-quantity contained in any part of its domain assumes a stationary value in relation to such variations of the potentials  $g_{\mu\nu}$ ,  $\phi_\mu$  as vanish at the limits of

\* Ann. d. Physik, 87, 89, 40, 1912-13.

## 212 GRAVITATION AND ELECTRICITY

the territory in question.  $W$ , the world-density of the action, must be an invariant of weight - 2. The action-quantity is in any case a pure number; thus our theory at once accounts for that atomistic structure of the world to which current views attach the most fundamental importance—the action-quantum. The simplest and most natural conjecture which we can make for  $W$ , is

$$W = R_{\nu\rho\sigma}^{\mu} R_{\mu}^{\nu\rho\sigma} = |R|^2.$$

For this we also have, by (13),

$$W = |P|^2 + 4L.$$

(There could be no doubt about anything here except perhaps the factor 4, with which the electric term  $L$  is added to the first.) But even without particularizing the action-quantity we can draw some general conclusions from the principle of action. For we shall show that as, according to investigations by Hilbert, Lorentz, Einstein, Klein, and the author,\* the four laws of the conservation of matter (the energy-momentum tensor) are connected with the invariance of the action quantity (containing four arbitrary functions) with respect to transformations of co-ordinates, so in the same way the law of the conservation of electricity is connected with the "measure-invariance" [transition from (8) to (9)] which here makes its appearance for the first time, introducing a fifth arbitrary function. The manner in which the latter associates itself with the principles of energy and momentum seems to me one of the strongest general arguments in favour of the theory here set out—so far as there can be any question at all of confirmation in purely speculative matters.

For any variation which vanishes at the limits of the world-territory under consideration we have

$$\delta \int \mathfrak{M} dx = \int (\mathfrak{M}^{\mu\nu} \delta g_{\mu\nu} + w^{\mu} \delta \phi_{\mu}) dx \quad (\mathfrak{M}^{\mu\nu} = \mathfrak{M}^{\nu\mu}) \quad (15)$$

\* Hilbert, "Die Grundlagen der Physik," Göttinger Nachrichten, 20 Nov., 1915; H. A. Lorentz in four papers in the Versl. K. Ak. van Wetensch., Amsterdam, 1915-16; A. Einstein, Berl. Ber., 1916, pp. 1111-6; F. Klein, Gött. Nachr., 25 Jan., 1918; H. Weyl, Ann. d. Physik, 54, 1917, pp. 121-5.

The laws of nature then take the form

$$\mathfrak{B}^{\mu\nu} = 0, w^\mu = 0 \quad (16)$$

The former may be regarded as the laws of the gravitational field, the latter as those of the electromagnetic field. The quantities  $W_\nu^\mu, w^\mu$  defined by

$$\mathfrak{B}_\nu^\mu = \sqrt{g} W_\nu^\mu, w^\mu = \sqrt{g} w^\mu$$

are the mixed or, respectively, the contravariant components of a tensor of rank 2 or 1 respectively, and of weight - 2. In the system of equations (16) there are five which are redundant, in accordance with the properties of invariance. This is expressed in the following five invariant identities, which subsist between their left-hand sides:—

$$\frac{\partial w^\mu}{\partial x_\mu} \equiv \mathfrak{B}_\mu^\mu \quad (17)$$

$$\frac{\partial \mathfrak{B}_\nu^\mu}{\partial x_\mu} - \Gamma_{\nu\beta}^\alpha \mathfrak{B}_\alpha^\beta \equiv \frac{1}{2} F_{\mu\nu} w^\mu \quad (18)$$

The first results from the measure-invariance. For if in the transition from (8) to (9) we assume for  $\log \lambda$  an infinitely small function of position  $\delta\rho$ , we obtain the variation

$$\delta g_{\mu\nu} = g_{\mu\nu} \delta\rho, \quad \delta\phi_\mu = \frac{\partial(\delta\rho)}{\partial x_\mu}.$$

For this variation (15) must vanish. In the second place if we utilize the invariance of the action-quantity with respect to transformations of co-ordinates by means of an infinitely small deformation of the world-continuum,\* we obtain the identities

$$\frac{\partial \mathfrak{B}_\nu^\mu}{\partial x_\mu} - \frac{1}{2} \frac{\partial g_{\alpha\beta}}{\partial x_\nu} \mathfrak{B}^{\alpha\beta} + \frac{1}{2} \left( \frac{\partial w^\mu}{\partial x_\mu} \phi_\nu - \Gamma_{\alpha\nu}^\mu w^\alpha \right) \equiv 0$$

which change into (18) when, by (17)  $\partial w^\mu / \partial x_\mu$  is replaced by  $g_{\alpha\beta} \mathfrak{B}^{\alpha\beta}$

From the gravitational laws alone therefore we already obtain

$$\frac{\partial w^\mu}{\partial x_\mu} = 0, \quad (19)$$

\* Weyl, Ann. d. Physik, 54, 1917, pp. 121-5; F. Klein, Gött. Nachr., 25 Jan., 1918.

## 214 GRAVITATION AND ELECTRICITY

and from the laws of the electromagnetic field alone

$$\frac{\partial}{\partial x_\mu} \mathfrak{M}^\mu_\nu - \Gamma^\alpha_{\nu\beta} \mathfrak{M}^\beta_\alpha = 0 \quad . \quad . \quad . \quad (20)$$

In Maxwell's theory  $w^\mu$  has the form

$$w^\mu \equiv \frac{\partial(\sqrt{g} F^{\mu\nu})}{\partial x_\nu} - \mathfrak{s}^\mu, \quad \mathfrak{s}^\mu = \sqrt{g} s^\mu$$

where  $s^\mu$  denotes the four-current. Since the first part here satisfies the equation (19) identically, this equation gives us the law of conservation of electricity

$$\frac{1}{\sqrt{g}} \frac{\partial(\sqrt{g} s^\mu)}{\partial x_\mu} = 0.$$

Similarly in Einstein's theory of gravitation  $\mathfrak{M}^\mu_\nu$  consists of two terms, the first of which satisfies equation (20) identically, and the second is equal to the mixed components of the energy-momentum tensor  $T^\mu_\nu$  multiplied by  $\sqrt{g}$ . Thus equations (20) lead to the four laws of the conservation of matter. Quite analogous circumstances hold good in our theory if we choose the form (14) for the action-quantity. The five principles of conservation are "eliminants" of the field laws, i.e. they follow from them in a twofold manner, and thus demonstrate that among them there are five which are redundant.

With the form (14) for the action-quantity the Maxwell equations run, for example:—

$$\frac{1}{\sqrt{g}} \frac{\partial(\sqrt{g} F^{\mu\nu})}{\partial x_\nu} = s^\mu, \quad . \quad . \quad . \quad (21)$$

and the current is

$$s_\mu = \frac{1}{4} \left( R \phi_\mu + \frac{\partial R}{\partial x_\mu} \right),$$

where  $R$  denotes that invariant of weight  $-1$  which arises from  $R^\mu_{\nu\rho\sigma}$  if we first contract with respect to  $\mu, \rho$  and then with respect to  $\nu$  and  $\sigma$ . If  $R^*$  denotes Riemann's invariant of curvature constructed solely from the  $g_{\mu\nu}$ , calculation

gives

$$R = R^* - \frac{3}{\sqrt{g}} \frac{\partial(\sqrt{g}\phi^\mu)}{\partial x_\mu} + \frac{3}{2}\phi_\mu\phi^\mu.$$

In the static case, where the space components of the electromagnetic potential disappear, and all quantities are independent of the time  $x_0$ , by (21) we must have

$$R = R^* + \frac{3}{2}\phi_0\phi^0 = \text{const.}$$

But in a world-territory in which  $R \neq 0$  we may make  $R = \text{const.} = \pm 1$  everywhere, by appropriate determination of the unit of length. Only we have to expect, under conditions which are variable with time, surfaces  $R = 0$ , which evidently will play some singular part.  $R$  cannot be used as density of action (represented by  $R^*$  in Einstein's theory of gravitation) because it has not the weight  $-2$ . The consequence is that though our theory leads to Maxwell's electromagnetic equations, it does not lead to Einstein's gravitation equations. In their place appear differential equations of order 4. But indeed it is very improbable that Einstein's equations of gravitation are strictly correct, because, above all things, the gravitation constant occurring in them is not at all in the picture with the other constants of nature, the gravitation radius of the charge and mass of an electron, for example, being of an entirely different order of magnitude ( $10^{20}$  or  $10^{40}$  times as small) from that of the radius of the electron itself.\*

It was my intention here merely to develop briefly the general principles of the theory.† The problem naturally

\* Cf. Weyl, Ann. d. Physik, 54, 1917, p. 183.

† [The problem of defining all  $W$  invariants allowable as action-quantities, under the requirement that they should contain the derivatives of the  $g_{\mu\nu}$  only to the second order at most, and those of the  $\phi_\mu$  only to the first order, was solved by R. Weitzenböck (Sitzungsber. d. Akad. d. Wissensch. in Wien, 129, 1920; 130, 1921). If we omit the invariants  $W$  for which the variation  $\delta W d\omega$  vanishes identically, there remain according to a later calculation by R. Bach (Math. Zeitschrift, 9, 1921, pp. 125 and 189) only three combinations. The real  $W$  seems to be a linear combination of Maxwell's  $L$  and the square of  $R$ . This conjecture has been tested more carefully by W. Pauli (Physik. Zeitschrift, 20, 1919, pp. 457-67) and myself; in particular we succeeded in advancing so far on this basis as to deduce the equations of



## 216 GRAVITATION AND ELECTRICITY

presents itself of deducing the physical consequences of the theory on the basis of the special form for the action-quantity given in (14), and of comparing these with experience, examining particularly whether the existence of the electron and the peculiarities of the hitherto unexplained processes in the atom can be deduced from the theory.\* The task is extraordinarily complicated from the mathematical point of view, because it is impossible to obtain approximate solutions if we restrict ourselves to the linear terms; for since it is certainly not permissible to neglect terms of higher order in the interior of the electron, the linear equations obtained by neglecting these may have, in general, only the solution 0. I propose to return to all these matters in greater detail in another place.

motion of a material particle. The invariant (14) selected above, at hazard in the first place, seems on the contrary to play no part in nature. Cf. Raum, "Zeit, Materie," ed. 4, §§ 35, 36, or Weyl, Physik. Zeitschr., 22, 1921, pp. 478-80.]

\* [Meanwhile I have quite abandoned these hopes, raised by Mie's theory; I do not believe that the problem of matter is to be solved by a mere field theory. Cf. on this subject my article "Feld und Materie," Ann. d. Physik, 65, 1921, pp. 541-63.]





# CATALOGUE OF DOVER BOOKS

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### General

**WHAT IS SCIENCE?**, Norman Campbell. This excellent introduction explains scientific method, role of mathematics, types of scientific laws. Contents: 2 aspects of science, science & nature, laws of science, discovery of laws, explanation of laws, measurement & numerical laws, applications of science. 192pp. 5½ x 8. S43 Paperbound \$1.25

**THE COMMON SENSE OF THE EXACT SCIENCES**, W. K. Clifford. Introduction by James Newman, edited by Karl Pearson. For 70 years this has been a guide to classical scientific and mathematical thought. Explains with unusual clarity basic concepts, such as extension of meaning of symbols, characteristics of surface boundaries, properties of plane figures, vectors, Cartesian method of determining position, etc. Long preface by Bertrand Russell. Bibliography of Clifford. Corrected, 130 diagrams redrawn. 249pp. 5½ x 8. T61 Paperbound \$1.60

**SCIENCE THEORY AND MAN**, Erwin Schrödinger. This is a complete and unabridged reprint of *SCIENCE AND THE HUMAN TEMPERAMENT* plus an additional essay: "What is an Elementary Particle?" Nobel laureate Schrödinger discusses such topics as nature of scientific method, the nature of science, chance and determinism, science and society, conceptual models for physical entities, elementary particles and wave mechanics. Presentation is popular and may be followed by most people with little or no scientific training. "Fine practical preparation for a time when laws of nature, human institutions . . . are undergoing a critical examination without parallel," Waldemar Kaempffert, N. Y. TIMES. 192pp. 5½ x 8. T428 Paperbound \$1.35

**FADS AND FALLACIES IN THE NAME OF SCIENCE**, Martin Gardner. Examines various cults, quack systems, frauds, delusions which at various times have masqueraded as science. Accounts of hollow-earth fanatics like Symmes; Velikovsky and wandering planets; Hoerbiger; Bellamy and the theory of multiple moons; Charles Fort; dowsing, pseudoscientific methods for finding water, ores, oil. Sections on naturopathy, iridagnosis, zone therapy, food fads, etc. Analytical accounts of Wilhelm Reich and orgone sex energy; L. Ron Hubbard and Dianetics; A. Korzybski and General Semantics; many others. Brought up to date to include Bridey Murphy, others. Not just a collection of anecdotes, but a fair, reasoned appraisal of eccentric theory. Formerly titled *IN THE NAME OF SCIENCE*. Preface. Index. x + 384pp. 5½ x 8. T394 Paperbound \$1.50

**A DOVER SCIENCE SAMPLER**, edited by George Barkin. 64-page book, sturdily bound, containing excerpts from over 20 Dover books, explaining science. Edwin Hubble, George Sarton, Ernst Mach, A. d'Abro, Galileo, Newton, others, discussing island universes, scientific truth, biological phenomena, stability in bridges, etc. Copies limited; no more than 1 to a customer, FREE

**POPULAR SCIENTIFIC LECTURES**, Hermann von Helmholtz. Helmholtz was a superb expositor as well as a scientist of genius in many areas. The seven essays in this volume are models of clarity, and even today they rank among the best general descriptions of their subjects ever written. "The Physiological Causes of Harmony in Music" was the first significant physiological explanation of musical consonance and dissonance. Two essays, "On the Interaction of Natural Forces" and "On the Conservation of Force," were of great importance in the history of science, for they firmly established the principle of the conservation of energy. Other lectures include "On the Relation of Optics to Painting," "On Recent Progress in the Theory of Vision," "On Goethe's Scientific Researches," and "On the Origin and Significance of Geometrical Axioms." Selected and edited with an introduction by Professor Morris Kline. xii + 286pp. 5½ x 8½. T799 Paperbound \$1.45

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### Physics

**CONCERNING THE NATURE OF THINGS**, Sir William Bragg. Christmas lectures delivered at the Royal Society by Nobel laureate. Why a spinning ball travels in a curved track; how uranium is transmuted to lead, etc. Partial contents: atoms, gases, liquids, crystals, metals, etc. No scientific background needed; wonderful for intelligent child. 32pp. of photos, 57 figures. xii + 232pp. 5½ x 8. T31 Paperbound \$1.50

**THE RESTLESS UNIVERSE**, Max Born. New enlarged version of this remarkably readable account by a Nobel laureate. Moving from sub-atomic particles to universe, the author explains in very simple terms the latest theories of wave mechanics. Partial contents: air and its relatives, electrons & ions, waves & particles, electronic structure of the atom, nuclear physics. Nearly 1000 illustrations, including 7 animated sequences. 325pp. 6 x 8. T412 Paperbound \$2.00

## Catalogue of Dover Books

**FROM EUCLID TO EDDINGTON: A STUDY OF THE CONCEPTIONS OF THE EXTERNAL WORLD.** Sir Edmund Whittaker. A foremost British scientist traces the development of theories of natural philosophy from the western rediscovery of Euclid to Eddington, Einstein, Dirac, etc. The inadequacy of classical physics is contrasted with present day attempts to understand the physical world through relativity, non-Euclidean geometry, space curvature, wave mechanics, etc. 5 major divisions of examination: Space; Time and Movement; the Concepts of Classical Physics; the Concepts of Quantum Mechanics; the Eddington Universe. 212pp. 5½ x 8. T491 Paperbound \$1.35

**PHYSICS, THE PIONEER SCIENCE.** L. W. Taylor. First thorough text to place all important physical phenomena in cultural-historical framework; remains best work of its kind. Exposition of physical laws, theories-developed chronologically, with great historical, illustrative experiments diagrammed, described, worked out mathematically. Excellent physics text for self-study as well as class work. Vol. 1: Heat, Sound, motion, acceleration, gravitation, conservation of energy, heat engines, rotation, heat, mechanical energy, etc. 211 illus. 407pp. 5½ x 8. Vol. 2: Light, Electricity: Images, lenses, prisms, magnetism, Ohm's law, dynamos, telegraph, quantum theory, decline of mechanical view of nature, etc. Bibliography. 13 table appendix. Index. 551 illus. 2 color plates. 508pp. 5½ x 8.

Vol. 1 S565 Paperbound \$2.25  
Vol. 2 S566 Paperbound \$2.25  
The set \$4.50

**A SURVEY OF PHYSICAL THEORY.** Max Planck. One of the greatest scientists of all time, creator of the quantum revolution in physics, writes in non-technical terms of his own discoveries and those of other outstanding creators of modern physics. Planck wrote this book when science had just crossed the threshold of the new physics, and he communicates the excitement felt then as he discusses electromagnetic theories, statistical methods, evolution of the concept of light, a step-by-step description of how he developed his own momentous theory, and many more of the basic ideas behind modern physics. Formerly "A Survey of Physics." Bibliography. Index. 128pp. 5½ x 8. S650 Paperbound \$1.15

**THE ATOMIC NUCLEUS.** M. Korsunsky. The only non-technical comprehensive account of the atomic nucleus in English. For college physics students, etc. Chapters cover: Radioactivity, the Nuclear Model of the Atom, the Mass of Atomic Nuclei, the Disintegration of Atomic Nuclei, the Discovery of the Positron, the Artificial Transformation of Atomic Nuclei, Artificial Radioactivity, Mesons, the Neutrino, the Structure of Atomic Nuclei and Forces Acting Between Nuclear Particles, Nuclear Fission, Chain Reaction, Peaceful Uses, Thermocuclear Reactions. Slightly abridged edition. Translated by G. Yankovsky. 65 figures. Appendix includes 45 photographic illustrations. 413 pp. 5½ x 8. S1052 Paperbound \$2.00

**PRINCIPLES OF MECHANICS SIMPLY EXPLAINED.** Morton Mott-Smith. Excellent, highly readable introduction to the theories and discoveries of classical physics. Ideal for the layman who desires a foundation which will enable him to understand and appreciate contemporary developments in the physical sciences. Discusses: Density, The Law of Gravitation, Mass and Weight, Action and Reaction, Kinetic and Potential Energy, The Law of Inertia, Effects of Acceleration, The Independence of Motions, Galileo and the New Science of Dynamics, Newton and the New Cosmos, The Conservation of Momentum, and other topics. Revised edition of "This Mechanical World." Illustrated by E. Kosa, Jr. Bibliography and Chronology. Index. xiv + 171pp. 5½ x 8½. T1067 Paperbound \$1.35

**THE CONCEPT OF ENERGY SIMPLY EXPLAINED.** Morton Mott-Smith. Elementary, non-technical exposition which traces the story of man's conquest of energy, with particular emphasis on the developments during the nineteenth century and the first three decades of our own century. Discusses man's earlier efforts to harness energy, more recent experiments and discoveries relating to the steam engine, the engine indicator, the motive power of heat, the principle of excluded perpetual motion, the bases of the conservation of energy, the concept of entropy, the internal combustion engine, mechanical refrigeration, and many other related topics. Also much biographical material. Index. Bibliography. 33 illustrations. ix + 215pp. 5½ x 8½. T1071 Paperbound \$1.25

**HEAT AND ITS WORKINGS.** Morton Mott-Smith. One of the best elementary introductions to the theory and attributes of heat, covering such matters as the laws governing the effect of heat on solids, liquids and gases, the methods by which heat is measured, the conversion of a substance from one form to another through heating and cooling, evaporation, the effects of pressure on boiling and freezing points, and the three ways in which heat is transmitted (conduction, convection, radiation). Also brief notes on major experiments and discoveries. Concise, but complete, it presents all the essential facts about the subject in readable style. Will give the layman and beginning student a first-rate background in this major topic in physics. Index. Bibliography. 50 illustrations. x + 165pp. 5½ x 8½. T978 Paperbound \$1.15

**THE STORY OF ATOMIC THEORY AND ATOMIC ENERGY.** J. G. Feinberg. Wider range of facts on physical theory, cultural implications, than any other similar source. Completely non-technical. Begins with first atomic theory, 600 B.C., goes through A-bomb, developments to 1959. Avogadro, Rutherford, Bohr, Einstein, radioactive decay, binding energy, radiation danger, future benefits of nuclear power, dozens of other topics, told in lively, related, informal manner. Particular stress on European atomic research. "Deserves special mention . . . authoritative," Saturday Review. Formerly "The Atom Story." New chapter to 1959. Index. 34 illustrations. 251pp. 5½ x 8. T625 Paperbound \$1.80

## Catalogue of Dover Books

**THE STRANGE STORY OF THE QUANTUM, AN ACCOUNT FOR THE GENERAL READER OF THE GROWTH OF IDEAS UNDERLYING OUR PRESENT ATOMIC KNOWLEDGE.** B. Hoffmann. Presents lucidly and expertly, with barest amount of mathematics, the problems and theories which led to modern quantum physics. Dr. Hoffmann begins with the closing years of the 19th century, when certain trifling discrepancies were noticed, and with illuminating analogies and examples takes you through the brilliant concepts of Planck, Einstein, Pauli, de Broglie, Bohr, Schroedinger, Heisenberg, Dirac, Sommerfeld, Feynman, etc. This edition includes a new, long postscript carrying the story through 1958. "Of the books attempting an account of the history and contents of our modern atomic physics which have come to my attention, this is the best," H. Margenau, Yale University, in "American Journal of Physics," 32 tables and line illustrations. Index. 275pp. 5½ x 8. T518 Paperbound \$1.75

**THE EVOLUTION OF SCIENTIFIC THOUGHT FROM NEWTON TO EINSTEIN.** A. d'Abro. Einstein's special and general theories of relativity, with their historical implications, are analyzed in non-technical terms. Excellent accounts of the contributions of Newton, Riemann, Weyl, Planck, Eddington, Maxwell, Lorentz and others are treated in terms of space and time, equations of electromagnetics, finiteness of the universe, methodology of science. 21 diagrams. 482pp. 5½ x 8. T2 Paperbound \$2.25

**THE RISE OF THE NEW PHYSICS, A. d'Abro.** A half-million word exposition, formerly titled **THE DECLINE OF MECHANISM**, for readers not versed in higher mathematics. The only thorough explanation, in everyday language, of the central core of modern mathematical physical theory, treating both classical and modern theoretical physics, and presenting in terms almost anyone can understand the equivalent of 5 years of study of mathematical physics. Scientifically impeccable coverage of mathematical-physical thought from the Newtonian system up through the electronic theories of Dirac and Heisenberg and Fermi's statistics. Combines both history and exposition; provides a broad yet unified and detailed view, with constant comparison of classical and modern views on phenomena and theories. "A must for anyone doing serious study in the physical sciences," JOURNAL OF THE FRANKLIN INSTITUTE. "Extraordinary faculty . . . to explain ideas and theories of theoretical physics in the language of daily life," *ISIS*. First part of set covers philosophy of science, drawing upon the practice of Newton, Maxwell, Poincaré, Einstein, others, discussing modes of thought, experiment, interpretations of causality, etc. In the second part, 100 pages explain grammar and vocabulary of mathematics, with discussions of functions, groups, series, Fourier series, etc. The remainder is devoted to concrete, detailed coverage of both classical and quantum physics, explaining such topics as analytic mechanics, Hamilton's principle, wave theory of light, electromagnetic waves, groups of transformations, thermodynamics, phase rule, Brownian movement, kinetics, special relativity, Planck's original quantum theory, Bohr's atom, Zeeman effect, Broglie's wave mechanics, Heisenberg's uncertainty, Eigen-values, matrices, scores of other important topics. Discoveries and theories are covered for such men as Alembert, Born, Cantor, Debye, Euler, Foucault, Galois, Gauss, Hadamard, Kelvin, Kepler, Laplace, Maxwell, Pauli, Rayleigh, Volterra, Weyl, Young, more than 180 others. Indexed. 97 illustrations. ix + 982pp. 5½ x 8. T3 Volume 1, Paperbound \$2.25  
T4 Volume 2, Paperbound \$2.25

**SPINNING TOPS AND GYROSCOPIC MOTION,** John Perry. Well-known classic of science still unsurpassed for lucid, accurate, delightful exposition. How quasi-rigidity is induced in flexible and fluid bodies by rapid motions; why gyrostap fails, top rises; nature and effect on climatic conditions of earth's precessional movement; effect of internal fluidity on rotating bodies, etc. Appendixes describe practical uses to which gyroscopes have been put in ships, compasses, monorail transportation. 62 figures. 128pp. 5½ x 8. T416 Paperbound \$1.25

**THE UNIVERSE OF LIGHT,** Sir William Bragg. No scientific training needed to read Nobel Prize winner's expansion of his Royal Institute Christmas Lectures. Insight into nature of light, methods and philosophy of science. Explains lenses, reflection, color, resonance, polarization, x-rays, the spectrum, Newton's work with prisms, Huygens' with polarization, Crookes' with cathode ray, etc. Leads into clear statement of 2 major historical theories of light, corpuscle and wave. Dozens of experiments you can do. 199 illus., including 2 full-page color plates. 293pp. 5½ x 8. S538 Paperbound \$1.85

**THE STORY OF X-RAYS FROM RÖNTGEN TO ISOTOPES,** A. R. Bleich. Non-technical history of x-rays, their scientific explanation, their applications in medicine, industry, research, and art, and their effect on the individual and his descendants. Includes amusing early reactions to Röntgen's discovery, cancer therapy, detections of art and stamp forgeries, potential risks to patient and operator, etc. Illustrations show x-rays of flower structure, the gall bladder, gears with hidden defects, etc. Original Dover publication. Glossary. Bibliography. Index. 55 photos and figures. xiv + 186pp. 5½ x 8. T662 Paperbound \$1.50

**ELECTRONS, ATOMS, METALS AND ALLOYS,** Wm. Hume-Rothery. An introductory-level explanation of the application of the electronic theory to the structure and properties of metals and alloys, taking into account the new theoretical work done by mathematical physicists. Material presented in dialogue-form between an "Old Metallurgist" and a "Young Scientist." Their discussion falls into 4 main parts: the nature of an atom, the nature of a metal, the nature of an alloy, and the structure of the nucleus. They cover such topics as the hydrogen atom, electron waves, wave mechanics, Brillouin zones, co-valent bonds, radioactivity and natural disintegration, fundamental particles, structure and fission of the nucleus, etc. Revised, enlarged edition. 177 illustrations. Subject and name indexes. 407pp. 5½ x 8½. S1046 Paperbound \$2.25

## Catalogue of Dover Books

**TEACH YOURSELF MECHANICS, P. Abbott.** The lever, centre of gravity, parallelogram of force, friction, acceleration, Newton's laws of motion, machines, specific gravity, gas, liquid pressure, much more. 280 problems, solutions. Tables. 163 illus. 271pp. 6½ x 4¼.  
Clothbound \$2.00

**MATTER & MOTION, James Clerk Maxwell.** This excellent exposition begins with simple particles and proceeds gradually to physical systems beyond complete analysis: motion, force, properties of centre of mass of material system, work, energy, gravitation, etc. Written with all Maxwell's original insights and clarity. Notes by E. Larmor. 17 diagrams. 178pp. 5½ x 8.  
\$188 Paperbound \$1.35

**SOAP BUBBLES, THEIR COLOURS AND THE FORCES WHICH MOULD THEM, C. V. Boys.** Only complete edition, half again as much material as any other. Includes Boys' hints on performing his experiments, sources of supply. Dozens of lucid experiments show complexities of liquid films, surface tension, etc. Best treatment ever written. Introduction. 83 illustrations. Color plate. 202pp. 5½ x 8.  
T542 Paperbound 95¢

**MATTER & LIGHT, THE NEW PHYSICS, L. de Broglie.** Non-technical papers by a Nobel laureate explain electromagnetic theory, relativity, matter, light and radiation, wave mechanics, quantum physics, philosophy of science. Einstein, Planck, Bohr, others explained so easily that no mathematical training is needed for all but 2 of the 21 chapters. Unabridged. Index. 300pp. 5½ x 8.  
T35 Paperbound \$1.85

**SPACE AND TIME, Emile Borel.** An entirely non-technical introduction to relativity, by world-renowned mathematician, Sorbonne professor. (Notes on basic mathematics are included separately.) This book has never been surpassed for insight, and extraordinary clarity of thought, as it presents scores of examples, analogies, arguments, illustrations, which explain such topics as: difficulties due to motion; gravitation a force of inertia; geodesic lines; wave-length and difference of phase; x-rays and crystal structure; the special theory of relativity; and much more. Indexes. 4 appendixes. 15 figures. xvi + 243pp. 5½ x 8.  
T592 Paperbound \$1.45

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### Astronomy

**THE FRIENDLY STARS, Martha Evans Martin.** This engaging survey of stellar lore and science is a well-known classic, which has introduced thousands to the fascinating world of stars and other celestial bodies. Descriptions of Capella, Sirius, Arcturus, Vega, Polaris, etc.—all the important stars, with informative discussions of rising and setting of stars, their number, names, brightness, distances, etc. in a non-technical, highly readable style. Also: double stars, constellations, clusters—concentrating on stars and formations visible to the naked eye. New edition, revised (1963) by D. H. Menzel, Director Harvard Observatory. 23 diagrams by Prof. Ching-Sung Yu. Foreword by D. H. Menzel and W. W. Morgan. 2 Star Charts. Index. xii + 147pp. 5¾ x 8½.  
T1099 Paperbound \$1.00

**AN ELEMENTARY SURVEY OF CELESTIAL MECHANICS, Y. Ryabov.** Elementary exposition of gravitational theory and celestial mechanics. Historical introduction and coverage of basic principles, including: the elliptic, the orbital plane, the 2- and 3-body problems, the discovery of Neptune, planetary rotation, the length of the day, the shapes of galaxies, satellites (detailed treatment of Sputnik I), etc. First American reprinting of successful Russian popular exposition. Elementary algebra and trigonometry helpful, but not necessary; presentation chiefly verbal. Appendix of theorem proofs. 58 figures. 165pp. 5½ x 8.  
T756 Paperbound \$1.25

**THE SKY AND ITS MYSTERIES, E. A. Beut.** One of most lucid books on mysteries of universe; deals with astronomy from earliest observations to latest theories of expansion of universe, source of stellar energy, birth of planets, origin of moon craters, possibility of life on other planets. Discusses effects of sunspots on weather; distances, ages of several stars; master plan of universe; methods and tools of astronomers; much more. "Eminently readable book," London Times. Extensive bibliography. Over 50 diagrams. 12 full-page plates, fold-out star map. Introduction. Index. 5¼ x 7½.  
T627 Clothbound \$3.50

**THE REALM OF THE NEBULAE, E. Hubble.** One of the great astronomers of our time records his formulation of the concept of "island universes," and its impact on astronomy. Such topics are covered as the velocity-distance relation; classification, nature, distances, general field of nebulae; cosmological theories; nebulae in the neighborhood of the Milky Way. 39 photos of nebulae, nebulae clusters, spectra of nebulae, and velocity distance relations shown by spectrum comparison. "One of the most progressive lines of astronomical research," The Times (London). New introduction by A. Sandage. 55 illustrations. Index. iv + 201pp. 5½ x 8.  
\$455 Paperbound \$1.50



## Catalogue of Dover Books

**OUT OF THE SKY, H. H. Nininger.** A non-technical but comprehensive introduction to "meteoritics", the young science concerned with all aspects of the arrival of matter from outer space. Written by one of the world's experts on meteorites, this work shows how, despite difficulties of observation and sparseness of data, a considerable body of knowledge has arisen. It defines meteors and meteorites; studies fireball clusters and processions, meteorite composition, size, distribution, showers, explosions, origins, craters, and much more. A true connecting link between astronomy and geology. More than 175 photos, 22 other illustrations. References. Bibliography of author's publications on meteorites. Index. viii + 336pp. 5 3/8 x 8. T519 Paperbound \$1.85

**SATELLITES AND SCIENTIFIC RESEARCH, D. King-Hele.** Non-technical account of the manmade satellites and the discoveries they have yielded up to the autumn of 1961. Brings together information hitherto published only in hard-to-get scientific journals. Includes the life history of a typical satellite, methods of tracking, new information on the shape of the earth, zones of radiation, etc. Over 60 diagrams and 6 photographs. Mathematical appendix. Bibliography of over 100 items. Index. xii + 180pp. 5 3/8 x 8 1/2. T703 Paperbound \$2.00

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### Mathematics

**CHANCE, LUCK AND STATISTICS: THE SCIENCE OF CHANCE, Horace C. Levinson.** Theory of probability and science of statistics in simple, non-technical language. Part I deals with theory of probability, covering odd superstitions in regard to "luck," the meaning of betting odds, the law of mathematical expectation, gambling, and applications in poker, roulette, lotteries, dice, bridge, and other games of chance. Part II discusses the misuse of statistics, the concept of statistical probabilities, normal and skew frequency distributions, and statistics applied to various fields—birth rates, stock speculation, insurance rates, advertising, etc. "Presented in an easy humorous style which I consider the best kind of expository writing," Prof. A. C. Cohen, Industry Quality Control. Enlarged revised edition. Formerly titled "The Science of Chance." Preface and two new appendices by the author. Index. xiv + 365pp. 5 3/8 x 8. T1007 Paperbound \$1.85

**PROBABILITIES AND LIFE, Emile Borel.** Translated by M. Baudin. Non-technical, highly readable introduction to the results of probability as applied to everyday situations. Partial contents: Fallacies About Probabilities Concerning Life After Death; Negligible Probabilities and the Probabilities of Everyday Life; Events of Small Probability; Application of Probabilities to Certain Problems of Heredity; Probabilities of Deaths, Diseases, and Accidents; On Poisson's Formula. Index. 3 Appendices of statistical studies and tables. vi + 87pp. 5 3/8 x 8 1/2. T121 Paperbound \$1.00

**GREAT IDEAS OF MODERN MATHEMATICS: THEIR NATURE AND USE, Jagjit Singh.** Reader with only high school math will understand main mathematical ideas of modern physics, astronomy, genetics, psychology, evolution, etc., better than many who use them as tools, but comprehend little of their basic structure. Author uses his wide knowledge of non-mathematical fields in brilliant exposition of differential equations, matrices, group theory, logic, statistics, problems of mathematical foundations, imaginary numbers, vectors, etc. Original publication. 2 appendices. 2 Indexes. 65 illustr. 322pp. 5 3/8 x 8. S587 Paperbound \$1.75

**MATHEMATICS IN ACTION, O. G. Sutton.** Everyone with a command of high school algebra will find this book one of the finest possible introductions to the application of mathematics to physical theory. Ballistics, numerical analysis, waves and wavelike phenomena, Fourier series, group concepts, fluid flow and aerodynamics, statistical measures, and meteorology are discussed with unusual clarity. Some calculus and differential equations theory is developed by the author for the reader's help in the more difficult sections. 88 figures. Index. viii + 236pp. 5 3/8 x 8. T440 Clothbound \$3.50

**THE FOURTH DIMENSION SIMPLY EXPLAINED, edited by H. P. Manning.** 22 essays, originally Scientific American contest entries, that use a minimum of mathematics to explain aspects of 4-dimensional geometry: analogues to 3-dimensional space, 4-dimensional absurdities and curiosities (such as removing the contents of an egg without puncturing its shell), possible measurements and forms, etc. Introduction by the editor. Only book of its sort on a truly elementary level, excellent introduction to advanced works. 82 figures. 251pp. 5 3/8 x 8. T711 Paperbound \$1.35

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### Engineering, technology, applied science etc.

**TEACH YOURSELF ELECTRICITY**, C. W. Wilman. Electrical resistance, inductance, capacitance, magnets, chemical effects of current, alternating currents, generators and motors, transformers, rectifiers, much more. 230 questions, answers, worked examples. List of units. 115 illus. 194pp. 6½ x 4¼. Clothbound \$2.00

**ELEMENTARY METALLURGY AND METALLOGRAPHY**, A. M. Shrager. Basic theory and descriptions of most of the fundamental manufacturing processes involved in metallurgy. Partial contents: the structure of metals; slip, plastic deformation, and recrystallization; iron ore and production of pig iron; chemistry involved in the metallurgy of iron and steel; basic processes such as the Bessemer treatment, open-hearth process, the electric arc furnace—with advantages and disadvantages of each; annealing, hardening, and tempering steel; copper, aluminum, magnesium, and their alloys. For freshman engineers, advanced students in technical high schools, etc. Index. Bibliography. 177 diagrams. 17 tables. 284 questions and problems. 27-page glossary. ix + 389pp. 5½ x 8. S138 Paperbound \$2.25

**BASIC ELECTRICITY**, Prepared by the Bureau of Naval Personnel. Originally a training course text for U.S. Navy personnel, this book provides thorough coverage of the basic theory of electricity and its applications. Best book of its kind for either broad or more limited studies of electrical fundamentals . . . for classroom use or home study. Part 1 provides a more limited coverage of theory: fundamental concepts, batteries, the simple circuit, D.C. series and parallel circuits, conductors and wiring techniques, A.C. electricity, inductance and capacitance, etc. Part 2 applies theory to the structure of electrical machines—generators, motors, transformers, magnetic amplifiers. Also deals with more complicated instruments, synchros, servo-mechanisms. The concluding chapters cover electrical drawings and blueprints, wiring diagrams, technical manuals, and safety education. The book contains numerous questions for the student, with answers. Index and six appendices. 345 illustrations. x + 448pp. 6½ x 9¼. S973 Paperbound \$3.00

**BASIC ELECTRONICS**, prepared by the U.S. Navy Training Publications Center. A thorough and comprehensive manual on the fundamentals of electronics. Written clearly, it is equally useful for self-study or course work for those with a knowledge of the principles of basic electricity. Partial contents: Operating Principles of the Electron Tube; Introduction to Transistors; Power Supplies for Electronic Equipment; Tuned Circuits; Electron-Tube Amplifiers; Audio Power Amplifiers; Oscillators; Transmitters; Transmission Lines; Antennas and Propagation; Introduction to Computers; and related topics. Appendix. Index. Hundreds of illustrations and diagrams. vi + 471pp. 6½ x 9¼. S1076 Paperbound \$2.75

**BASIC THEORY AND APPLICATION OF TRANSISTORS**, Prepared by the U.S. Department of the Army. An introductory manual prepared for an army training program. One of the finest available surveys of theory and application of transistor design and operation. Minimal knowledge of physics and theory of electron tubes required. Suitable for textbook use, course supplement, or home study. Chapters: Introduction; fundamental theory of transistors; transistor amplifier fundamentals; parameters, equivalent circuits, and characteristic curves; bias stabilization; transistor analysis and comparison using characteristic curves and charts; audio amplifiers; tuned amplifiers; wide-band amplifiers; oscillators; pulse and switching circuits; modulation, mixing, and demodulation; and additional semiconductor devices. Unabridged, corrected edition. 240 schematic drawings, photographs, wiring diagrams, etc. 2 Appendices. Glossary. Index. 263pp. 6½ x 9¼. S380 Paperbound \$1.25

**TEACH YOURSELF HEAT ENGINES**, E. De Ville. Measurement of heat, development of steam and internal combustion engines, efficiency of an engine, compression-ignition engines, production of steam, the ideal engine, much more. 318 exercises, answers, worked examples. Tables. 76 illus. 220pp. 6½ x 4¼. Clothbound \$2.25

## BOOKS EXPLAINING SCIENCE AND MATHEMATICS

### Miscellaneous

**ON THE SENSATIONS OF TONE**, Hermann Helmholtz. This is an unmatched coordination of such fields as acoustical physics, physiology, experiment, history of music. It covers the entire gamut of musical tone. Partial contents: relation of musical science to acoustics, physical vs. physiological acoustics, composition of vibration, resonance, analysis of tones by sympathetic resonance, beats, chords, tonality, consonant chords, dissonances, progression of parts, etc. 33 appendices discuss various aspects of sound, physics, acoustics, music, etc. Translated by A. J. Ellis. New introduction by Prof. Henry Margenau of Yale. 68 figures. 43 musical passages analyzed. Over 100 tables. Index. xix + 576pp. 6½ x 9¼. S114 Paperbound \$3.50

## Catalogue of Dover Books

**THE NATURE OF LIGHT AND COLOUR IN THE OPEN AIR**, M. Minnaert. Why is falling snow sometimes black? What causes mirages, the fata morgana, multiple suns and moons in the sky? How are shadows formed? Prof. Minnaert of the University of Utrecht answers these and similar questions in optics, light, colour, for non-specialists. Particularly valuable to nature, science students, painters, photographers. Translated by H. M. Kremer-Priest, K. Jay. 202 illustrations, including 42 photos. xvi + 362pp. 5½ x 8. T196 Paperbound \$2.00

**THE PHYSICS OF MUSIC**, Alexander Wood. Introduction for musicians to the physical aspect of sound. No scientific training necessary to understand concepts, etc. Wealth of material on origin and development of instruments, physical principles involved in the production of their sounds, pitch, intensity and loudness, mechanism of the ear, dissonance and consonance, sound reproduction and recordings, concert halls, etc. Extensively revised by Dr. J. M. Bowsher. Indices. Bibliography. 16 plates. 114 illustrations. 270pp. 5½ x 8½. T322 Paperbound \$2.25

**GREAT IDEAS AND THEORIES OF MODERN COSMOLOGY**, Jagjit Singh. The theories of Jeans, Eddington, Milne, Kant, Bondi, Gold, Newton, Einstein, Gamow, Hoyle, Dirac, Kuiper, Hubble, Weizsäcker and many others on such cosmological questions as the origin of the universe, space and time, planet formation, "continuous creation," the birth, life, and death of the stars, the origin of the galaxies, etc. By the author of the popular "Great Ideas of Modern Mathematics." A gifted popularizer of science, he makes the most difficult abstractions crystal-clear even to the most non-mathematical reader. Index. xii + 276 pp. 5½ x 8½. T925 Paperbound \$1.85

**PIONEERS OF SCIENCE**, O. Lodge. Eminent scientist-expositor's authoritative, yet elementary survey of great scientific theories. Concentrating on individuals—Copernicus, Brahe, Kepler, Galileo, Descartes, Newton, Laplace, Herschel, Lord Kelvin, and other scientists—the author presents their discoveries in historical order adding biographical material on each man and full, specific explanations of their achievements. The clear and complete treatment of the post-Newtonian astronomers is a feature seldom found in other books on the subject. Index. 120 illustrations. xv + 404pp. 5½ x 8. T716 Paperbound \$1.85

## BIOGRAPHY OF SCIENTISTS

**ISAAC NEWTON: A BIOGRAPHY**, Louis Trenchard More. The definitive biography of Newton, his life and work. Presents Newton as a living man, with a critical, objective analysis of his character as well as a careful survey of his manifold accomplishments, scientific, theological, etc. The author, himself a professor of physics, has made full use of all of Newton's published works and all material in the Portsmouth Collection of Newton's personal and unpublished papers. The text includes numerous letters by Newton and his acquaintances, and many other of his papers—some translated from Latin to English by the author. A universally-esteemed work. Unabridged republication. 1 full-page plate. Index. xiii + 675pp. 5½ x 8½. T579 Paperbound \$2.50

**PIERRE CURIE**, Marie Curie. Mme. Curie, Nobel Prize winner, creates a memorable portrait of her equally famous husband and his lifelong scientific researches. She brings to life the determined personality of a great scientist at work. Her own autobiographical notes, included in this volume, reconstruct her own work on radiation which resulted in the isolation of radium. "A delightful book. It marks one of the few instances in which the proverbially humdrum life of the student of physical science, together with the austere ideals, has been made intelligible," New York Times. Unabridged reprint. Translated by Charlotte and Vernon Kellogg. Introduction by Mrs. Wm. Brown Meloney. 8 halftones. viii + 120pp. 5½ x 8½. T199 Paperbound \$1.00

**THE BOOK OF MY LIFE (DE VITA PROPRIA LIBER)**, Jerome Cardan. The remarkable autobiography of an important Renaissance mathematician, physician, and scientist, who at the same time was a paranoid, morbid, superstitious man, consumed with ambition and self-love (and self-pity). These chronicles of his fortunes and misfortunes make absorbing reading, giving us an extremely insightful view of a man's reactions and sensations—the first psychological autobiography. Through his eyes we can also see the superstitions and beliefs of an age, Renaissance medical practices, and the problems that concerned a trained mind in the 16th century. Unabridged republication of original English edition, translated by Jean Stoner. Introduction. Notes. Bibliography. xviii + 331pp. 5½ x 8½. T345 Paperbound \$1.60

**THE AUTOBIOGRAPHY OF CHARLES DARWIN, AND SELECTED LETTERS**, edited by Francis Darwin. Darwin's own record of his early life; the historic voyage aboard the "Beagle"; the furor surrounding evolution, and his replies; reminiscences of his son. Letters to Henslow, Lyell, Hooker, Huxley, Wallace, Kingsley, etc., and thoughts on religion and vivisection. We see how he revolutionized geology with his concept of ocean subsidence; how his great books on variation of plants and animals, primitive man, the expression of emotion among primates, plant fertilization, carnivorous plants, protective coloration, etc., came into being. Appendix. Index. 365pp. 5½ x 8. T479 Paperbound \$2.00

## PHILOSOPHY OF SCIENCE AND MATHEMATICS

**FOUNDATIONS OF SCIENCE: THE PHILOSOPHY OF THEORY AND EXPERIMENT**, N. R. Campbell. A critique of the most fundamental concepts of science in general and physics in particular. Examines why certain propositions are accepted without question, demarcates science from philosophy, clarifies the understanding of the tools of science. Part One analyzes the pre-suppositions of scientific thought: existence of the material world, nature of scientific laws, multiplication of probabilities, etc.; Part Two covers the nature of experiment and the application of mathematics: conditions for measurement, relations between numerical laws and theories, laws of error, etc. An appendix covers problems arising from relativity, force, motion, space, and time. A classic in its field. Index. xiii + 565pp. 5% x 8%.

S372 Paperbound \$2.95

**THE NATURE OF PHYSICAL THEORY**, P. W. Bridgman. Here is how modern physics looks to a highly unorthodox physicist—a Nobel laureate. Pointing out many absurdities of science, and demonstrating the inadequacies of various physical theories, Dr. Bridgman weighs and analyzes the contributions of Einstein, Bohr, Newton, Heisenberg, and many others. This is a non-technical consideration of the correlation of science and reality. Index. xi + 138pp. 5% x 8.

S33 Paperbound \$1.25

**THE VALUE OF SCIENCE**, Henri Poincaré. Many of the most mature ideas of the "last scientific universalist" covered with charm and vigor for both the beginning student and the advanced worker. Discusses the nature of scientific truth, whether order is innate in the universe or imposed upon it by man, logical thought versus intuition (relating to math, through the works of Weierstrass, Lie, Klein, Riemann), time and space (relativity, psychological time, simultaneity), Hertz's concept of force, interrelationship of mathematical physics to pure math, values within disciplines of Maxwell, Carnot, Mayer, Newton, Lorentz, etc. Index. iii + 147pp. 5% x 8.

S469 Paperbound \$1.35

**SCIENCE AND HYPOTHESIS**, Henri Poincaré. Creative psychology in science. How such concepts as number, magnitude, space, force, classical mechanics were developed, and how the modern scientist uses them in his thought. Hypothesis in physics, theories of modern physics. Introduction by Sir James Larmor. "Few mathematicians have had the breadth of vision of Poincaré, and none is his superior in the gift of clear exposition," E. T. Bell. Index. 272pp. 5% x 8.

S221 Paperbound \$1.35

**PHILOSOPHY AND THE PHYSICISTS**, L. S. Stebbing. The philosophical aspects of modern science examined in terms of a lively critical attack on the ideas of Jeans and Eddington. Discusses the task of science, causality, determinism, probability, consciousness, the relation of the world of physics to that of everyday experience. Probes the philosophical significance of the Planck-Bohr concept of discontinuous energy levels, the inferences to be drawn from Heisenberg's Uncertainty Principle, the implications of "becoming" involved in the 2nd law of thermodynamics, and other problems posed by the discarding of Laplacean determinism. 285pp. 5% x 8.

T480 Paperbound \$1.65

**THE PHILOSOPHICAL WRITINGS OF PEIRCE**, edited by Justus Buchler. (Formerly published as **THE PHILOSOPHY OF PEIRCE**.) This is a carefully balanced exposition of Peirce's complete system, written by Peirce himself. It covers such matters as scientific method, pure chance vs. law, symbolic logic, theory of signs, pragmatism, experiment, and other topics. Introduction by Justus Buchler, Columbia University. xvi + 368pp. 5% x 8.

T217 Paperbound \$2.00

**LANGUAGE, TRUTH AND LOGIC**, A. Ayer. A clear introduction to the Vienna and Cambridge schools of Logical Positivism. It sets up specific tests by which you can evaluate validity of ideas, etc. Contents: Function of philosophy, elimination of metaphysics, nature of analysis, a priori, truth and probability, etc. 10th printing. "I should like to have written it myself," Bertrand Russell. Index. 160pp. 5% x 8.

T10 Paperbound \$1.25

**MATHEMATICS AND SCIENCE: LAST ESSAYS (DERNIÈRES PENSÉES)**, Henri Poincaré. Translated by J. W. Bolduc. A posthumous volume of articles and lectures by the great French mathematician, philosopher, scientist. Here are nine pieces, never before translated into English, on such subjects as The Evolution of Laws, Space and Time, Space and 3 Dimensions, The Logic of Infinity in Mathematics (discussing Russell's theory of types), Mathematics and Logic, The Quantum Theory and its Modern Applications, Relationship Between Matter and Ether, Ethics and Science and The Moral Alliance. First English translation of *Dernières Pensées*. New index. viii + 128pp. 5% x 8%.

S1101 Paperbound \$1.25

**THE PSYCHOLOGY OF INVENTION IN THE MATHEMATICAL FIELD**, J. Hadamard. Where do ideas come from? What role does the unconscious play? Are ideas best developed by mathematical reasoning, word reasoning, visualization? What are the methods used by Einstein, Poincaré, Galton, Riemann? How can these techniques be applied by others? Hadamard, one of the world's leading mathematicians, discusses these and other questions. xiii + 145pp. 5% x 8.

T107 Paperbound \$1.25

## *Catalogue of Dover Books*

**EXPERIMENT AND THEORY IN PHYSICS**, Max Born. A Nobel laureate examines the nature and value of the counterclaims of experiment and theory in physics. Synthetic versus analytical scientific advances are analyzed in the work of Einstein, Bohr, Heisenberg, Planck, Eddington, Milne, and others by a fellow participant. 44pp. 5½ x 8. S308 Paperbound 75¢

**THE PHILOSOPHY OF SPACE AND TIME**, H. Reichenbach. An Important landmark in the development of the empiricist conception of geometry, covering the problem of the foundations of geometry, the theory of time, the consequences of Einstein's relativity, including: relations between theory and observations; coordinate and metrical properties of space; the psychological problem of visual intuition of non-Euclidean structures; and many other important topics in modern science and philosophy. The majority of ideas require only a knowledge of intermediate math. Introduction by R. Carnap. 49 figures. Index. xviii + 296pp. 5½ x 8. S443 Paperbound \$2.00

**OBSERVATION AND INTERPRETATION IN THE PHILOSOPHY OF PHYSICS: WITH SPECIAL REFERENCE TO QUANTUM MECHANICS**, Edited by S. Körner. A collection of papers by philosophers and physicists arising out of a symposium held at Bristol, England in 1957 under the auspices of the Colston Research Society. One of the most important contributions to the philosophy of science in recent years. The discussions center around the adequacy or inadequacy of quantum mechanics in its orthodox formulations. Among the contributors are A. J. Ayer, D. Bohm, K. Popper, F. Bopp, S. Körner, J. P. Vigiér, M. Polanyi, P. K. Feyerabend, W. C. Kneale, W. B. Gallie, G. Ryle, Sir Charles Darwin, and R. B. Braithwaite. xiv + 218pp. 5½ x 8½. S131 Paperbound \$1.60

**SPACE AND TIME IN CONTEMPORARY PHYSICS: AN INTRODUCTION TO THE THEORY OF RELATIVITY AND GRAVITATION**, Moritz Schlick. Exposition of the theory of relativity by the leader of the famed "Vienna Circle." Its essential purpose is to describe the physical doctrines of special and general relativity with particular reference to their philosophical significance. Explanations of such topics as the geometrical relativity of space, the connection with inertia and gravitation, the measure-determination of the space-time continuum, the finite universe, etc., with their philosophical ramifications. Index. xli + 89pp. 5½ x 8½. T1008 Paperbound \$1.00

**SUBSTANCE AND FUNCTION, & EINSTEIN'S THEORY OF RELATIVITY**, Ernst Cassirer. Two books bound as one. Cassirer establishes a philosophy of the exact sciences that takes into consideration newer developments in mathematics, and also shows historical connections. Partial contents: Aristotelian logic, Mill's analysis, Helmholtz & Kronecker, Russell & cardinal numbers, Euclidean vs. non-Euclidean geometry, Einstein's relativity. Bibliography. Index. xxi + 465pp. 5½ x 8. T50 Paperbound \$2.25

**PRINCIPLES OF MECHANICS**, Heinrich Hertz. This last work by the great 19th century physicist is not only a classic, but of great interest in the logic of science. Creating a new system of mechanics based upon space, time, and mass, it returns to axiomatic analysis, to understanding of the formal or structural aspects of science, taking into account logic, observation, and a priori elements. Of great historical importance to Poincaré, Carnap, Einstein, Milne. A 20-page introduction by R. S. Cohen, Wesleyan University, analyzes the implications of Hertz's thought and the logic of science. Bibliography. 13-page introduction by Helmholtz. xlii + 274pp. 5½ x 8. S316 Clothbound \$3.50  
S317 Paperbound \$1.85

**THE ANALYSIS OF MATTER**, Bertrand Russell. How do our senses concord with the new physics? This volume covers such topics as logical analysis of physics, pre-relativity physics, causality, scientific inference, physics and perception, special and general relativity, Weyl's theory, tensors, invariants and their physical interpretation, periodicity and qualitative series. "The most thorough treatment of the subject that has yet been published." THE NATION. Introduction by L. E. Denonn. 422pp. 5½ x 8. T231 Paperbound \$1.95

**FOUNDATIONS OF GEOMETRY**, Bertrand Russell. Analyzing basic problems in the overlap area between mathematics and philosophy, Nobel laureate Russell examines the nature of geometrical knowledge, the nature of geometry, and the application of geometry to space. It covers the history of non-Euclidean geometry, philosophic interpretations of geometry—especially Kant—projective and metrical geometry. This is most interesting as the solution offered in 1897 by a great mind to a problem still current. New introduction by Prof. Morris Kline of N. Y. University. xli + 201pp. 5½ x 8. S232 Clothbound \$3.25  
S233 Paperbound \$1.75

**IDENTITY AND REALITY**, Emile Meyerson. Called by Einstein a "brilliant study in the theory of knowledge," this book by the renowned Franco-German thinker is a major treatise in the philosophy of science and epistemology. Thorough, critical inquiries into causality, scientific laws, conservation of matter and energy, the unity of matter, Carnot's principle, the irrational, the elimination of time. Searches out the solutions of epistemological questions that form the bases of the scientific method. Authorized translation by Kate Loewenberg. Author's prefaces. Editor's preface. Appendices. Index. 495pp. 5½ x 8½. T65 Paperbound \$2.25

**ESSAYS IN EXPERIMENTAL LOGIC**, John Dewey. This stimulating series of essays touches upon the relationship between inquiry and experience, dependence of knowledge upon thought, character of logic; judgments of practice, data and meanings, stimuli of thought, etc. Index. viii + 444pp. 5½ x 8. T73 Paperbound \$2.25

# PHYSICS

## General physics

**FOUNDATIONS OF PHYSICS**, R. B. Lindsay & H. Margenau. Excellent bridge between semi-popular works & technical treatises. A discussion of methods of physical description, construction of theory; valuable for physicist with elementary calculus who is interested in ideas that give meaning to data, tools of modern physics. Contents include symbolism, mathematical equations; space & time foundations of mechanics; probability; physics & continua; electron theory; special & general relativity; quantum mechanics; causality. "Thorough and yet not overdetailed. Unreservedly recommended," NATURE (London). Unabridged, corrected edition. List of recommended readings. 35 illustrations. xi + 537pp. 5% x 8. S377 Paperbound \$3.00

**FUNDAMENTAL FORMULAS OF PHYSICS**, ed. by D. H. Menzel. Highly useful, fully inexpensive reference and study text, ranging from simple to highly sophisticated operations. Mathematics integrated into text—each chapter stands as short textbook of field represented. Vol. 1: Statistics, Physical Constants, Special Theory of Relativity, Hydrodynamics, Aerodynamics, Boundary Value Problems in Math. Physics; Viscosity, Electromagnetic Theory, etc. Vol. 2: Sound, Acoustics, Geometrical Optics, Electron Optics, High-Energy Phenomena, Magnetism, Biophysics, much more. Index. Total of 800pp. 5% x 8. Vol. 1 S595 Paperbound \$2.25  
Vol. 2 S596 Paperbound \$2.25

**MATHEMATICAL PHYSICS**, D. H. Menzel. Thorough one-volume treatment of the mathematical techniques vital for classic mechanics, electromagnetic theory, quantum theory, and relativity. Written by the Harvard Professor of Astrophysics for junior, senior, and graduate courses, it gives clear explanations of all those aspects of function theory, vectors, matrices, dyadics, tensors, partial differential equations, etc., necessary for the understanding of the various physical theories. Electron theory, relativity, and other topics seldom presented appear here in considerable detail. Scores of definitions, conversion factors, dimensional constants, etc. "More detailed than normal for an advanced text . . . excellent set of sections on Dyadics, Matrices, and Tensors," JOURNAL OF THE FRANKLIN INSTITUTE. Index. 193 problems, with answers. x + 412pp. 5% x 8. S56 Paperbound \$2.00

**THE SCIENTIFIC PAPERS OF J. WILLARD GIBBS**. All the published papers of America's outstanding theoretical scientist (except for "Statistical Mechanics" and "Vector Analysis"). Vol. I (thermodynamics) contains one of the most brilliant of all 19th-century scientific papers—the 300-page "On the Equilibrium of Heterogeneous Substances," which founded the science of physical chemistry, and clearly stated a number of highly important natural laws for the first time; 8 other papers complete the first volume. Vol. II includes 2 papers on dynamics, 8 on vector analysis and multiple algebra, 5 on the electromagnetic theory of light, and 6 miscellaneous papers. Biographical sketch by H. A. Bumstead. Total of xxxvi + 718pp. 5% x 8%. S721 Vol I Paperbound \$2.50  
S722 Vol II Paperbound \$2.00  
The set \$4.50

**BASIC THEORIES OF PHYSICS**, Peter Gabriel Bergmann. Two-volume set which presents a critical examination of important topics in the major subdivisions of classical and modern physics. The first volume is concerned with classical mechanics and electrodynamics: mechanics of mass points, analytical mechanics, matter in bulk, electrostatics and magnetostatics, electromagnetic interaction, the field waves, special relativity, and waves. The second volume (Heat and Quanta) contains discussions of the kinetic hypothesis, physics and statistics, stationary ensembles, laws of thermodynamics, early quantum theories, atomic spectra, probability waves, quantization in wave mechanics, approximation methods, and abstract quantum theory. A valuable supplement to any thorough course or text. Heat and Quanta: Index. 8 figures. x + 300pp. 5% x 8½. S968 Paperbound \$2.00  
Mechanics and Electrodynamics: Index. 14 figures. vii + 280pp. 5% x 8½. S969 Paperbound \$1.85

**THEORETICAL PHYSICS**, A. S. Kompaneys. One of the very few thorough studies of the subject in this price range. Provides advanced students with a comprehensive theoretical background. Especially strong on recent experimentation and developments in quantum theory. Contents: Mechanics (Generalized Coordinates, Lagrange's Equation, Collision of Particles, etc.), Electrodynamics (Vector Analysis, Maxwell's equations, Transmission of Signals, Theory of Relativity, etc.), Quantum Mechanics (the inadequacy of Classical Mechanics, the Wave Equation, Motion in a Central Field, Quantum Theory of Radiation, Quantum Theories of Dispersion and Scattering, etc.), and Statistical Physics (Equilibrium Distribution of Molecules in an Ideal Gas, Boltzmann statistics, Bose and Fermi Distribution, Thermodynamic Quantities, etc.). Revised to 1961. Translated by George Yankovsky, authorized by Kompaneys. 137 exercises. 56 figures. 529pp. 5% x 8½. S972 Paperbound \$2.50

**ANALYTICAL AND CANONICAL FORMALISM IN PHYSICS**, André Mercier. A survey, in one volume, of the variational principles (the key principles—in mathematical form—from which the basic laws of any one branch of physics can be derived) of the several branches of physical theory, together with an examination of the relationships among them. Contents: the Lagrangian Formalism, Lagrangian Densities, Canonical Formalism, Canonical Form of Electrodynamics, Hamiltonian Densities, Transformations, and Canonical Form with Vanishing Jacobian Determinant. Numerous examples and exercises. For advanced students, teachers, etc. 6 figures. Index. viii + 222pp. 5% x 8½. S1077 Paperbound \$1.75

## Catalogue of Dover Books

### Acoustics, optics, electricity and magnetism, electromagnetics, magneto-hydrodynamics

**THE THEORY OF SOUND, Lord Rayleigh.** Most vibrating systems likely to be encountered in practice can be tackled successfully by the methods set forth by the great Nobel laureate, Lord Rayleigh. Complete coverage of experimental, mathematical aspects of sound theory. Partial contents: Harmonic motions, vibrating systems in general, lateral vibrations of bars, curved plates or shells, applications of Laplace's functions to acoustical problems, fluid friction, plane vortex-sheet, vibrations of solid bodies, etc. This is the first inexpensive edition of this great reference and study work. Bibliography. Historical introduction by R. B. Lindsay. Total of 1040pp. 97 figures. 5½ x 8.

\$292, \$293, Two volume set, paperbound, \$4.70

**THE DYNAMICAL THEORY OF SOUND, H. Lamb.** Comprehensive mathematical treatment of the physical aspects of sound, covering the theory of vibrations, the general theory of sound, and the equations of motion of strings, bars, membranes, pipes, and resonators. Includes chapters on plane, spherical, and simple harmonic waves, and the Helmholtz Theory of Audition. Complete and self-contained development for student and specialist; all fundamental differential equations solved completely. Specific mathematical details for such important phenomena as harmonics, normal modes, forced vibrations of strings, theory of reed pipes, etc. Index. Bibliography. 86 diagrams. viii + 307pp. 5½ x 8.

\$655 Paperbound \$2.00

**WAVE PROPAGATION IN PERIODIC STRUCTURES, L. Brillouin.** A general method and application to different problems: pure physics, such as scattering of X-rays of crystals, thermal vibration in crystal lattices, electronic motion in metals; and also problems of electrical engineering. Partial contents: elastic waves in 1-dimensional lattices of point masses. Propagation of waves along 1-dimensional lattices. Energy flow. 2 dimensional, 3 dimensional lattices. Mathieu's equation. Matrices and propagation of waves along an electric line. Continuous electric lines. 131 illustrations. Bibliography. Index. xii + 253pp. 5½ x 8.

\$34 Paperbound \$2.00

**THEORY OF VIBRATIONS, N. W. McLachlan.** Based on an exceptionally successful graduate course given at Brown University, this discusses linear systems having 1 degree of freedom, forced vibrations of simple linear systems, vibration of flexible strings, transverse vibrations of bars and tubes, transverse vibration of circular plate, sound waves of finite amplitude, etc. Index. 99 diagrams. 160pp. 5½ x 8.

\$190 Paperbound \$1.50

**LIGHT: PRINCIPLES AND EXPERIMENTS, George S. Monk.** Covers theory, experimentation, and research. Intended for students with some background in general physics and elementary calculus. Three main divisions: 1) Eight chapters on geometrical optics—fundamental concepts (the ray and its optical length, Fermat's principle, etc.); laws of image formation, apertures in optical systems, photometry, optical instruments etc.; 2) 9 chapters on physical optics—interference, diffraction, polarization, spectra, the Rayleigh refractometer, the wave theory of light, etc.; 3) 23 instructive experiments based directly on the theoretical text. "Probably the best intermediate textbook on light in the English language. Certainly, it is the best book which includes both geometrical and physical optics," J. Rud Nielson, PHYSICS FORUM. Revised edition. 102 problems and answers. 12 appendices. 6 tables. Index. 270 illustrations. xi + 489pp. 5½ x 8½.

\$341 Paperbound \$2.50

**PHOTOMETRY, John W. T. Walsh.** The best treatment of both "bench" and "illumination" photometry in English by one of Britain's foremost experts in the field (President of the International Commission on Illumination). Limited to those matters, theoretical and practical, which affect the measurement of light flux, candlepower, illumination, etc., and excludes treatment of the use to which such measurements may be put after they have been made. Chapters on Radiation, The Eye and Vision, Photo-Electric Cells, The Principles of Photometry, The Measurement of Luminous Intensity, Colorimetry, Spectrophotometry, Stellar Photometry, The Photometric Laboratory, etc. Third revised (1958) edition. 281 illustrations. 10 appendices. xiv + 544pp. 5½ x 9¼.

\$319 Paperbound \$3.00

**EXPERIMENTAL SPECTROSCOPY, R. A. Sawyer.** Clear discussion of prism and grating spectrographs and the techniques of their use in research, with emphasis on those principles and techniques that are fundamental to practically all uses of spectroscopic equipment. Beginning with a brief history of spectroscopy, the author covers such topics as light sources, spectroscopic apparatus, prism spectroscopes and graphs, diffraction grating, the photographic process, determination of wave length, spectral intensity, infrared spectroscopy, spectrochemical analysis, etc. This revised edition contains new material on the production of replica gratings, solar spectroscopy from rockets, new standard of wave length, etc. Index. Bibliography. 111 illustrations. x + 358pp. 5½ x 8½.

\$1045 Paperbound \$2.25

**FUNDAMENTALS OF ELECTRICITY AND MAGNETISM, L. B. Loeb.** For students of physics, chemistry, or engineering who want an introduction to electricity and magnetism on a higher level and in more detail than general elementary physics texts provide. Only elementary differential and integral calculus is assumed. Physical laws developed logically, from magnetism to electric currents, Ohm's law, electrolysis, and on to static electricity, induction, etc. Covers an unusual amount of material; one third of book on modern material: solution of wave equation, photoelectric and thermionic effects, etc. Complete statement of the various electrical systems of units and interrelations. 2 indexes. 75 pages of problems with answers stated. Over 300 figures and diagrams. xix + 669pp. 5½ x 8.

\$745 Paperbound \$3.50

## Catalogue of Dover Books

**ITICAL ANALYSIS OF ELECTRICAL AND OPTICAL WAVE-MOTION**, Harry Bateman. Written f this century's most distinguished mathematical physicists, this is a practical on to those developments of Maxwell's electromagnetic theory which are directly with the solution of the partial differential equation of wave motion. Methods of ave-equation, polar-cylindrical coordinates, diffraction, transformation of coordinates, ous solutions, electromagnetic fields with moving singularities, etc. Index. 168pp. \$14 Paperbound **\$1.75**

**IS OF PHYSICAL OPTICS**, Ernst Mach. This classical examination of the propagation color, polarization, etc. offers an historical and philosophical treatment that has n surpassed for breadth and easy readability. Contents: Rectilinear propagation of flection, refraction. Early knowledge of vision. Dioptrics. Composition of light. color and dispersion. Periodicity. Theory of Interference. Polarization. Mathematical ation of properties of light. Propagation of waves, etc. 279 Illustrations, 10 por-pendix. Indexes. 324pp. 5½ x 8. \$178 Paperbound **\$2.00**

**JRY OF OPTICS**, Paul Drude. One of finest fundamental texts in physical optics, ffers thorough coverage, complete mathematical treatment of basic ideas. Includes eatment of application of thermodynamics to optics; sine law in formation of transparent crystals, magnetically active substances, velocity of light, apertures, epending upon them, polarization, optical instruments, etc. Introduction by A. A. l. Index. 110 illus. 567pp. 5½ x 8. \$532 Paperbound **\$3.00**

**AL THEORY ON THE GIORGI SYSTEM**, P. Corneliuss. A new clarification of the funda- oncepts of electricity and magnetism, advocating the convenient m.k.s. system of t is steadily gaining followers in the sciences. Illustrating the use and effectiveness arminology with numerous applications to concrete technical problems, the author ounds the famous Giorgi system of electrical physics. His lucid presentation easoned, cogent argument for the universal adoption of this system form one of t pieces of scientific exposition in recent years. 28 figures. Index. Conversion tables ating earlier data into modern units. Translated from 3rd Dutch edition by L. J. + 187pp. 5½ x 8¾. \$909 Clothbound **\$6.00**

**WAVES: BEING RESEARCHES ON THE PROPAGATION OF ELECTRIC ACTION WITH ELOCITY THROUGH SPACE**, Heinrich Hertz. This classic work brings together the peters in which Hertz—Helmholtz's protégé and one of the most brilliant figures entury research—probed the existence of electromagnetic waves and showed experi- that their velocity equalled that of light, research that helped lay the groundwork evelopment of radio, television, telephone, telegraph, and other modern technological Unabridged republication of original edition. Authorized translation by D. E. Jones. y Lord Kelvin. Index of names. 40 illustrations. xvii + 278pp. 5½ x 8½. \$57 Paperbound **\$1.75**

**ELECTRICITY: AN INTRODUCTION TO THE THEORY AND APPLICATIONS OF ELECTRO- ICAL PHENOMENA IN CRYSTALS**, Walter G. Cady. This is the most complete and sys- everage of this important field in print—now regarded as something of scientific This republication, revised and corrected by Prof. Cady—one of the foremost con- In this area—contains a sketch of recent progress and new material on Ferro- Time Standards, etc. The first 7 chapters deal with fundamental theory of crystal ty. 5 important chapters cover basic concepts of piezoelectricity, including com- of various competing theories in the field. Also discussed: piezoelectric resonators methods of manufacture, influences of air-gaps, etc.); the piezo oscillator; the es, history, and observations relating to Rochelle salt; ferroelectric crystals; miscel- applications of piezoelectricity; pyroelectricity; etc. "A great work," W. A. Wooster, . Revised (1963) and corrected edition. New preface by Prof. Cady. 2 Appendices. Illustrations. 62 tables. Bibliography. Problems. Total of 1 + 822pp. 5½ x 8½. \$1094 Vol. I Paperbound **\$2.50**  
\$1095 Vol. II Paperbound **\$2.50**  
Two volume set Paperbound **\$5.00**

**ISM AND VERY LOW TEMPERATURES**, H. B. G. Casimir. A basic work in the literature emperature physics. Presents a concise survey of fundamental theoretical principles, o points out promising lines of investigation. Contents: Classical Theory and Experi- Methods, Quantum Theory of Paramagnetism, Experiments on Adiabatic Demagnetiza- eoretical Discussion of Paramagnetism at Very Low Temperatures, Some Experimental Relaxation Phenomena. Index. 89-item bibliography. ix + 95pp. 5½ x 8. \$943 Paperbound **\$1.35**

**ED PAPERS ON NEW TECHNIQUES FOR ENERGY CONVERSION: THERMOELECTRIC S; THERMIONIC; PHOTOVOLTAIC AND ELECTRICAL EFFECTS; FUSION**, Edited by Sumner ne. Brings together in one volume the most important papers (1954-1961) in modern echnology. Included among the 37 papers are general and qualitative descriptions field as a whole, indicating promising lines of research. Also: 15 papers on thermo- methods, 7 on thermionic, 5 on photovoltaic, 4 on electrochemical effect, and 2 on led fusion research. Among the contributors are: Joffe, Maria Telkes, Herold, Herring, , Jaumot, Post, Austin, Wilson, Pfann, Rappaport, Morehouse, Domenicali, Moss, , Harman, Von Doenhoeft. Preface and introduction by the editor. Bibliographies. + 451pp. 6½ x 9¾. \$37 Paperbound **\$3.00**



## Catalogue of Dover Books

**SUPERFLUIDS: MACROSCOPIC THEORY OF SUPERCONDUCTIVITY**, Vol. I, Fritz London. The major work by one of the founders and great theoreticians of modern quantum physics. Consolidates the researches that led to the present understanding of the nature of superconductivity. Prof. London here reveals that quantum mechanics is operative on the macroscopic plane as well as the submolecular level. Contents: Properties of Superconductors and Their Thermodynamical Correlation; Electrodynamics of the Pure Superconducting State; Relation between Current and Field; Measurements of the Penetration Depth, Non-Viscous Flow vs. Superconductivity; Micro-waves in Superconductors; Reality of the Domain Structure; and many other related topics. A new epilogue by M. J. Buckingham discusses developments in the field up to 1960. Corrected and expanded edition. An appreciation of the author's life and work by L. W. Nordheim. Biography by Edith London. Bibliography of his publications. 45 figures. 2 Indices. xviii + 173pp. 5½ x 8½. \$44 Paperbound \$1.75

**SELECTED PAPERS ON PHYSICAL PROCESSES IN IONIZED PLASMAS**, Edited by Donald H. Menzel, Director, Harvard College Observatory. 30 important papers relating to the study of highly ionized gases or plasmas selected by a foremost contributor in the field, with the assistance of Dr. L. H. Aller. The essays include 18 on the physical processes in gaseous nebulae, covering problems of radiation and radiative transfer, the Balmer decrement, electron temperatures, spectrophotometry, etc. 10 papers deal with the interpretation of nebular spectra, by Bohm, Van Vleck, Aller, Minkowski, etc. There is also a discussion of the intensities of "forbidden" spectral lines by George Shortley and a paper concerning the theory of hydrogenic spectra by Menzel and Pekeris. Other contributors: Goldberg, Hebb, Baker, Bowen, Ufford, Liller, etc. viii + 374pp. 6½ x 9¼. \$60 Paperbound \$2.95

**THE ELECTROMAGNETIC FIELD**, Max Mason & Warren Weaver. Used constantly by graduate engineers. Vector methods exclusively: detailed treatment of electrostatics, expansion methods, with tables converting any quantity into absolute electromagnetic, absolute electrostatic, practical units. Discrete charges, ponderable bodies, Maxwell field equations, etc. Introduction. Indexes. 416pp. 5½ x 8. \$185 Paperbound \$2.25

**THEORY OF ELECTRONS AND ITS APPLICATION TO THE PHENOMENA OF LIGHT AND RADIANT HEAT**, H. Lorentz. Lectures delivered at Columbia University by Nobel laureate Lorentz. Unabridged, they form a historical coverage of the theory of free electrons, motion, absorption of heat, Zeeman effect, propagation of light in molecular bodies, inverse Zeeman effect, optical phenomena in moving bodies, etc. 109 pages of notes explain the more advanced sections. Index. 9 figures. 352pp. 5½ x 8. \$173 Paperbound \$2.00

**FUNDAMENTAL ELECTROMAGNETIC THEORY**, Ronold P. King, Professor Applied Physics, Harvard University. Original and valuable introduction to electromagnetic theory and to circuit theory from the standpoint of electromagnetic theory. Contents: Mathematical Description of Matter—stationary and nonstationary states; Mathematical Description of Space and of Simple Media—Field Equations, Integral Forms of Field Equations, Electromagnetic Force, etc.; Transformation of Field and Force Equations; Electromagnetic Waves in Unbounded Regions; Skin Effect and Internal Impedance—in a solid cylindrical conductor, etc.; and Electrical Circuits—Analytical Foundations, Near-zone and quasi-near zone circuits, Balanced two-wire and four-wire transmission lines. Revised and enlarged version. New preface by the author. 5 appendices (Differential operators: Vector Formulas and Identities, etc.). Problems. Indexes. Bibliography. xvi + 580pp. 5½ x 8½. \$1023 Paperbound \$3.00

## Hydrodynamics

**A TREATISE ON HYDRODYNAMICS**, A. B. Basset. Favorite text on hydrodynamics for 2 generations of physicists, hydrodynamical engineers, oceanographers, ship designers, etc. Clear enough for the beginning student, and thorough source for graduate students and engineers on the work of d'Alembert, Euler, Laplace, Lagrange, Poisson, Green, Clebsch, Stokes, Cauchy, Helmholtz, J. J. Thomson, Love, Hicks, Greenhill, Besant, Lamb, etc. Great amount of documentation on entire theory of classical hydrodynamics. Vol I: theory of motion of frictionless liquids, vortex, and cyclic irrotational motion, etc. 132 exercises. Bibliography. 3 Appendixes. xli + 264pp. Vol II: motion in viscous liquids, harmonic analysis, theory of tides, etc. 112 exercises, Bibliography. 4 Appendixes. xv + 328pp. Two volume set. 5½ x 8.

\$724 Vol I Paperbound \$1.75

\$725 Vol II Paperbound \$1.75

The set \$3.50

**HYDRODYNAMICS**, Horace Lamb. Internationally famous complete coverage of standard reference work on dynamics of liquids & gases. Fundamental theorems, equations, methods, solutions, background, for classical hydrodynamics. Chapters Include Equations of Motion, Integration of Equations in Special Cases, Irrotational Motion, Motion of Liquid in 2 Dimensions, Motion of Solids through Liquid-Dynamical Theory, Vortex Motion, Tidal Waves, Surface Waves, Waves of Expansion, Viscosity, Rotating Masses of liquids. Excellently planned, arranged; clear, lucid presentation. 6th enlarged, revised edition. Index. Over 900 footnotes, mostly bibliographical. 119 figures. xv + 738pp. 6½ x 9¼. \$256 Paperbound \$3.75

## Catalogue of Dover Books

**HYDRODYNAMICS**, H. Dryden, F. Murnaghan, Harry Bateman. Published by the National Research Council in 1932 this enormous volume offers a complete coverage of classical hydrodynamics. Encyclopedic in quality. Partial contents: physics of fluids, motion, turbulent flow, compressible fluids, motion in 1, 2, 3 dimensions; viscous fluids rotating, laminar motion, resistance of motion through viscous fluid, eddy viscosity, hydraulic flow in channels of various shapes, discharge of gases, flow past obstacles, etc. Bibliography of over 2,900 items. Indexes. 23 figures. 634pp. 5½ x 8. S303 Paperbound \$2.75

## Mechanics, dynamics, thermodynamics, elasticity

**MECHANICS**, J. P. Den Hartog. Already a classic among introductory texts, the M.I.T. professor's lively and discursive presentation is equally valuable as a beginner's text, an engineering student's refresher, or a practicing engineer's reference. Emphasis in this highly readable text is on illuminating fundamental principles and showing how they are embodied in a great number of real engineering and design problems: trusses, loaded cables, beams, jacks, hoists, etc. Provides advanced material on relative motion and gyroscopes not usual in introductory texts. "Very thoroughly recommended to all those anxious to improve their real understanding of the principles of mechanics." *MECHANICAL WORLD*. Index. List of equations. 334 problems, all with answers. Over 550 diagrams and drawings. ix + 462pp. 5½ x 8.

S754 Paperbound \$2.00

**THEORETICAL MECHANICS: AN INTRODUCTION TO MATHEMATICAL PHYSICS**, J. S. Ames, F. D. Murnaghan. A mathematically rigorous development of theoretical mechanics for the advanced student, with constant practical applications. Used in hundreds of advanced courses. An unusually thorough coverage of gyroscopic and baryscopic material, detailed analyses of the Coriolis acceleration, applications of Lagrange's equations, motion of the double pendulum, Hamilton-Jacobi partial differential equations, group velocity and dispersion, etc. Special relativity is also included. 159 problems. 44 figures. ix + 462pp. 5½ x 8.

S461 Paperbound \$2.25

**THEORETICAL MECHANICS: STATICS AND THE DYNAMICS OF A PARTICLE**, W. D. MacMillan. Used for over 3 decades as a self-contained and extremely comprehensive advanced undergraduate text in mathematical physics, physics, astronomy, and deeper foundations of engineering. Early sections require only a knowledge of geometry; later, a working knowledge of calculus. Hundreds of basic problems, including projectiles to the moon, escape velocity, harmonic motion, ballistics, falling bodies, transmission of power, stress and strain, elasticity, astronomical problems. 340 practice problems plus many fully worked out examples make it possible to test and extend principles developed in the text. 200 figures. xvii + 430pp. 5½ x 8.

S467 Paperbound \$2.25

**THEORETICAL MECHANICS: THE THEORY OF THE POTENTIAL**, W. D. MacMillan. A comprehensive, well balanced presentation of potential theory, serving both as an introduction and a reference work with regard to specific problems, for physicists and mathematicians. No prior knowledge of integral relations is assumed, and all mathematical material is developed as it becomes necessary. Includes: Attraction of Finite Bodies; Newtonian Potential Function; Vector Fields, Green and Gauss Theorems; Attraction of Surfaces and Lines; Surface Distribution of Matter; Two-Layer Surfaces; Spherical Harmonics; Ellipsoidal Harmonics; etc. "The great number of particular cases . . . should make the book valuable to geophysicists and others actively engaged in practical applications of the potential theory." *Review of Scientific Instruments*. Index. Bibliography. xiii + 469pp. 5½ x 8.

S486 Paperbound \$2.50

**THEORETICAL MECHANICS: DYNAMICS OF RIGID BODIES**, W. D. MacMillan. Theory of dynamics of a rigid body is developed, using both the geometrical and analytical methods of instruction. Begins with exposition of algebra of vectors, it goes through momentum principles, motion in space, use of differential equations and infinite series to solve more sophisticated dynamics problems. Partial contents: moments of inertia, systems of free particles, motion parallel to a fixed plane, rolling motion, method of periodic solutions, much more. 82 figs. 199 problems. Bibliography. Indexes. xii + 476pp. 5½ x 8.

S641 Paperbound \$2.50

**MATHEMATICAL FOUNDATIONS OF STATISTICAL MECHANICS**, A. I. Khinchin. Offering a precise and rigorous formulation of problems, this book supplies a thorough and up-to-date exposition. It provides analytical tools needed to replace cumbersome concepts, and furnishes for the first time a logical step-by-step introduction to the subject. Partial contents: geometry & kinematics of the phase space, ergodic problem, reduction to theory of probability, application of central limit problem, ideal monatomic gas, foundation of thermo-dynamics, dispersion and distribution of sum functions. Key to notations. Index. viii + 179pp. 5½ x 8.

S147 Paperbound \$1.50

**ELEMENTARY PRINCIPLES IN STATISTICAL MECHANICS**, J. W. Gibbs. Last work of the great Yale mathematical physicist, still one of the most fundamental treatments available for advanced students and workers in the field. Covers the basic principle of conservation of probability of phase, theory of errors in the calculated phases of a system, the contributions of Clausius, Maxwell, Boltzmann, and Gibbs himself, and much more. Includes valuable comparison of statistical mechanics with thermodynamics: Carnot's cycle, mechanical definitions of entropy, etc. xvi + 208pp. 5½ x 8.

S707 Paperbound \$1.45

## Catalogue of Dover Books

**PRINCIPLES OF MECHANICS AND DYNAMICS**, Sir William Thomson (Lord Kelvin) and Peter Guthrie Tait. The principles and theories of fundamental branches of classical physics explained by two of the greatest physicists of all time. A broad survey of mechanics, with material on hydrodynamics, elasticity, potential theory, and what is now standard mechanics. Thorough and detailed coverage, with many examples, derivations, and topics not included in more recent studies. Only a knowledge of calculus is needed to work through this book. Vol. I (Preliminary): Kinematics; Dynamical Laws and Principles; Experience (observation, experimentation, formation of hypotheses, scientific method); Measures and Instruments; Continuous Calculating Machines. Vol. II (Abstract Dynamics): Statics of a Particle—Attraction; Statics of Solids and Fluids. Formerly Titled "Treatise on Natural Philosophy." Unabridged reprint of revised edition. Index. 168 diagrams. Total of xlii + 1035pp. 5½ x 8½.  
Vol. I: S966 Paperbound \$2.35  
Vol. II: S967 Paperbound \$2.35  
Two volume Set Paperbound \$4.70

**INVESTIGATIONS ON THE THEORY OF THE BROWNIAN MOVEMENT**, Albert Einstein. Reprints from rare European journals. 5 basic papers, including the Elementary Theory of the Brownian Movement, written at the request of Lorentz to provide a simple explanation. Translated by A. D. Cowper. Annotated, edited by R. Fürth. 33pp. of notes elucidate, give history of previous investigations. Author, subject indexes. 62 footnotes. 124pp. 5½ x 8.  
S304 Paperbound \$1.25

**MECHANICS VIA THE CALCULUS**, P. W. Norris, W. S. Legge. Covers almost everything, from linear motion to vector analysis: equations determining motion, linear methods, compounding of simple harmonic motions, Newton's laws of motion, Hooke's law, the simple pendulum, motion of a particle in 1 plane, centers of gravity, virtual work, friction, kinetic energy of rotating bodies, equilibrium of strings, hydrostatics, shearing stresses, elasticity, etc. 550 problems. 3rd revised edition. xli + 367pp. 6 x 9.  
S207 Clothbound \$4.95

**THE DYNAMICS OF PARTICLES AND OF RIGID, ELASTIC, AND FLUID BODIES; BEING LECTURES ON MATHEMATICAL PHYSICS**, A. G. Webster. The reissuing of this classic fills the need for a comprehensive work on dynamics. A wide range of topics is covered in unusually great depth, applying ordinary and partial differential equations. Part I considers laws of motion and methods applicable to systems of all sorts; oscillation, resonance, cyclic systems, etc. Part 2 is a detailed study of the dynamics of rigid bodies. Part 3 introduces the theory of potential; stress and strain, Newtonian potential functions, gyrostatics, wave and vortex motion, etc. Further contents: Kinematics of a point; Lagrange's equations; Hamilton's principle; Systems of vectors; Statics and dynamics of deformable bodies; much more, not easily found together in one volume. Unabridged reprinting of 2nd edition. 20 pages of notes on differential equations and the higher analysis. 203 illustrations. Selected bibliography. Index. xi + 588pp. 5½ x 8.  
S522 Paperbound \$2.45

**A TREATISE ON DYNAMICS OF A PARTICLE**, E. J. Routh. Elementary text on dynamics for beginning mathematics or physics student. Unusually detailed treatment from elementary definitions to motion in 3 dimensions, emphasizing concrete aspects. Much unique material important in recent applications. Covers impulsive forces, rectilinear and constrained motion in 2 dimensions, harmonic and parabolic motion, degrees of freedom, closed orbits, the conical pendulum, the principle of least action, Jacobi's method, and much more. Index. 559 problems, many fully worked out, incorporated into text. xlii + 418pp. 5½ x 8.  
S696 Paperbound \$2.25

**DYNAMICS OF A SYSTEM OF RIGID BODIES (Elementary Section)**, E. J. Routh. Revised 7th edition of this standard reference. This volume covers the dynamical principles of the subject, and its more elementary applications: finding moments of inertia by integration, foci of inertia, d'Alembert's principle, impulsive forces, motion in 2 and 3 dimensions, Lagrange's equations, relative indicatrix, Euler's theorem, large tautochronous motions, etc. Index. 55 figures. Scores of problems. xv + 443pp. 5½ x 8.  
S664 Paperbound \$2.50

**DYNAMICS OF A SYSTEM OF RIGID BODIES (Advanced Section)**, E. J. Routh. Revised 6th edition of a classic reference aid. Much of its material remains unique. Partial contents: moving axes, relative motion, oscillations about equilibrium, motion. Motion of a body under no forces, any forces. Nature of motion given by linear equations and conditions of stability. Free, forced vibrations, constants of integration, calculus of finite differences, variations, precession and nutation, motion of the moon, motion of string, chain, membranes. 64 figures. 498pp. 5½ x 8.  
S229 Paperbound \$2.45

**DYNAMICAL THEORY OF GASES**, James Jeans. Divided into mathematical and physical chapters for the convenience of those not expert in mathematics, this volume discusses the mathematical theory of gas in a steady state, thermodynamics, Boltzmann and Maxwell, kinetic theory, quantum theory, exponentials, etc. 4th enlarged edition, with new material on quantum theory, quantum dynamics, etc. Indexes. 28 figures. 444pp. 6½ x 9¾.  
S136 Paperbound \$2.75

**THE THEORY OF HEAT RADIATION**, Max Planck. A pioneering work in thermodynamics, providing basis for most later work, Nobel laureate Planck writes on Deductions from Electrodynamics and Thermodynamics, Entropy and Probability, Irreversible Radiation Processes, etc. Starts with simple experimental laws of optics, advances to problems of spectral distribution of energy and irreversibility. Bibliography. 7 illustrations, xiv + 224pp. 5½ x 8.  
S546 Paperbound \$1.75

## Catalogue of Dover Books

**FOUNDATIONS OF POTENTIAL THEORY, O. D. Kellogg.** Based on courses given at Harvard this is suitable for both advanced and beginning mathematicians. Proofs are rigorous, and much material not generally available elsewhere is included. Partial contents: forces of gravity, fields of force, divergence theorem, properties of Newtonian potentials at points of free space, potentials as solutions of Laplace's equations, harmonic functions, electrostatics, electric images, logarithmic potential, etc. One of Grundlehren Series. ix + 384pp. 5½ x 8. \$144 Paperbound \$2.00

**THERMODYNAMICS, Enrico Fermi.** Unabridged reproduction of 1937 edition. Elementary in treatment; remarkable for clarity, organization. Requires no knowledge of advanced math beyond calculus, only familiarity with fundamentals of thermometry, calorimetry. Partial Contents: Thermodynamic systems; First & Second laws of thermodynamics; Entropy; Thermodynamic potentials: phase rule, reversible electric cell; Gaseous reactions: van't Hoff reaction box, principle of LeChatelier; Thermodynamics of dilute solutions: osmotic & vapor pressures, boiling & freezing points; Entropy constant. Index. 25 problems. 24 illustrations. x + 160pp. 5½ x 8. \$361 Paperbound \$1.75

**THE THERMODYNAMICS OF ELECTRICAL PHENOMENA IN METALS and A CONDENSED COLLECTION OF THERMODYNAMIC FORMULAS, P. W. Bridgman.** Major work by the Nobel Prizewinner: stimulating conceptual introduction to aspects of the electron theory of metals, giving an intuitive understanding of fundamental relationships concealed by the formal systems of Onsager and others. Elementary mathematical formulations show clearly the fundamental thermodynamical relationships of the electric field, and a complete phenomenological theory of metals is created. This is the work in which Bridgman announced his famous "thermomotive force" and his distinction between "driving" and "working" electromotive force. We have added in this Dover edition the author's long unavailable tables of thermodynamic formulas, extremely valuable for the speed of reference they allow. Two works bound as one. Index. 33 figures. Bibliography. xviii + 256pp. 5½ x 8. \$723 Paperbound \$1.75

**TREATISE ON THERMODYNAMICS, Max Planck.** Based on Planck's original papers this offers a uniform point of view for the entire field and has been used as an introduction for students who have studied elementary chemistry, physics, and calculus. Rejecting the earlier approaches of Helmholtz and Maxwell, the author makes no assumptions regarding the nature of heat, but begins with a few empirical facts, and from these deduces new physical and chemical laws. 3rd English edition of this standard text by a Nobel laureate. xvi + 297pp. 5½ x 8. \$219 Paperbound \$1.85

**THE MATHEMATICAL THEORY OF ELASTICITY, A. E. H. Love.** A wealth of practical illustration combined with thorough discussion of fundamentals—theory, application, special problems and solutions. Partial Contents: Analysis of Strain & Stress, Elasticity of Solid Bodies, Elasticity of Crystals, Vibration of Spheres, Cylinders, Propagation of Waves in Elastic Solid Media, Torsion, Theory of Continuous Beams, Plates. Rigorous treatment of Volterra's theory of dislocations, 2-dimensional elastic systems, other topics of modern interest. "For years the standard treatise on elasticity." AMERICAN MATHEMATICAL MONTHLY. 4th revised edition. Index. 76 figures. xviii + 643pp. 6½ x 9¼. \$174 Paperbound \$3.25

**STRESS WAVES IN SOLIDS, H. Kolsky, Professor of Applied Physics, Brown University.** The most readable survey of the theoretical core of current knowledge about the propagation of waves in solids, fully correlated with experimental research. Contents: Part I—Elastic Waves: propagation in an extended plastic medium, propagation in bounded elastic media, experimental investigations with elastic materials. Part II—Stress Waves in Imperfectly Elastic Media: internal friction, experimental investigations of dynamic elastic properties, plastic waves and shock waves, fractures produced by stress waves. List of symbols. Appendix. Supplemented bibliography. 3 full-page plates. 46 figures. x + 213pp. 5½ x 8½. \$1098 Paperbound \$1.75

## Relativity, quantum theory, atomic and nuclear physics

**SPACE TIME MATTER, Hermann Weyl.** "The standard treatise on the general theory of relativity" (Nature), written by a world-renowned scientist, provides a deep clear discussion of the logical coherence of the general theory, with introduction to all the mathematical tools needed: Maxwell, analytical geometry, non-Euclidean geometry, tensor calculus, etc. Basis is classical space-time, before absorption of relativity. Partial contents: Euclidean space, mathematical form, metrical continuum, relativity of time and space, general theory. 15 diagrams. Bibliography. New preface for this edition. xviii + 330pp. 5½ x 8. \$267 Paperbound \$2.25

**ATOMIC SPECTRA AND ATOMIC STRUCTURE, G. Herzberg.** Excellent general survey for chemists, physicists specializing in other fields. Partial contents: simplest line spectra and elements of atomic theory, building-up principle and periodic system of elements, hyperfine structure of spectral lines, some experiments and applications. Bibliography. 80 figures. Index. xli + 257pp. 5½ x 8. \$115 Paperbound \$2.00

## Catalogue of Dover Books

**THE PRINCIPLE OF RELATIVITY**, A. Einstein, H. Lorentz, H. Minkowski, H. Weyl. These are the 11 basic papers that founded the general and special theories of relativity, all translated into English. Two papers by Lorentz on the Michelson experiment, electromagnetic phenomena. Minkowski's SPACE & TIME, and Weyl's GRAVITATION & ELECTRICITY. 7 epoch-making papers by Einstein: ELECTROMAGNETICS OF MOVING BODIES, INFLUENCE OF GRAVITATION IN PROPAGATION OF LIGHT, COSMOLOGICAL CONSIDERATIONS, GENERAL THEORY, and 3 others. 7 diagrams. Special notes by A. Sommerfeld. 224pp. 5½ x 8.

\$81 Paperbound \$2.00

**EINSTEIN'S THEORY OF RELATIVITY**, Max Born. Revised edition prepared with the collaboration of Gunther Leibfried and Walter Blem. Steering a middle course between superficial popularizations and complex analyses, a Nobel laureate explains Einstein's theories clearly and with special insight. Easily followed by the layman with a knowledge of high school mathematics, the book has been thoroughly revised and extended to modernize those sections of the well-known original edition which are now out of date. After a comprehensive review of classical physics, Born's discussion of special and general theories of relativity covers such topics as simultaneity, kinematics, Einstein's mechanics and dynamics, relativity of arbitrary motions, the geometry of curved surfaces, the space-time continuum, and many others. Index. Illustrations, vii + 376pp. 5½ x 8.

\$769 Paperbound \$2.00

**ATOMS, MOLECULES AND QUANTA**, Arthur E. Ruark and Harold C. Urey. Revised (1963) and corrected edition of a work that has been a favorite with physics students and teachers for more than 30 years. No other work offers the same combination of atomic structure and molecular physics and of experiment and theory. The first 14 chapters deal with the origins and major experimental data of quantum theory and with the development of conceptions of atomic and molecular structure prior to the new mechanics. These sections provide a thorough introduction to atomic and molecular theory, and are presented lucidly and as simply as possible. The six subsequent chapters are devoted to the laws and basic ideas of quantum mechanics: Wave Mechanics, Hydrogenic Atoms in Wave Mechanics, Matrix Mechanics, General Theory of Quantum Dynamics, etc. For advanced college and graduate students in physics. Revised, corrected republication of original edition, with supplementary notes by the authors. New preface by the authors. 9 appendices. General reference list. Indices. 228 figures. 71 tables. Bibliographical material in notes, etc. Total of xxiii + 810pp. 5½ x 8½.

\$1106 Vol. I Paperbound \$2.50

\$1107 Vol. II Paperbound \$2.50

Two volume set Paperbound \$5.00

**WAVE MECHANICS AND ITS APPLICATIONS**, N. F. Mott and I. N. Sneddon. A comprehensive introduction to the theory of quantum mechanics; not a rigorous mathematical exposition it progresses, instead, in accordance with the physical problems considered. Many topics difficult to find at the elementary level are discussed in this book. Includes such matters as: the wave nature of matter, the wave equation of Schrödinger, the concept of stationary states, properties of the wave functions, effect of a magnetic field on the energy levels of atoms, electronic spin, two-body problem, theory of solids, cohesive forces in ionic crystals, collision problems, interaction of radiation with matter, relativistic quantum mechanics, etc. All are treated both physically and mathematically. 68 illustrations. 11 tables. Indexes. xii + 393pp. 5½ x 8½.

\$1070 Paperbound \$2.35

**BASIC METHODS IN TRANSFER PROBLEMS**, V. Kourganoff, Professor of Astrophysics, U. of Paris. A coherent digest of all the known methods which can be used for approximate or exact solutions of transfer problems. All methods demonstrated on one particular problem—Milne's problem for a plane parallel medium. Three main sections: fundamental concepts (the radiation field and its interaction with matter, the absorption and emission coefficients, etc.); different methods by which transfer problems can be attacked; and a more general problem—the non-gray case of Milne's problem. Much new material, drawing upon declassified atomic energy reports and data from the USSR. Entirely understandable to the student with a reasonable knowledge of analysis. Unabridged, revised reprinting. New preface by the author. Index. Bibliography. 2 appendices. xv + 281pp. 5½ x 8½.

\$1074 Paperbound \$2.00

**PRINCIPLES OF QUANTUM MECHANICS**, W. V. Houston. Enables student with working knowledge of elementary mathematical physics to develop facility in use of quantum mechanics, understand published work in field. Formulates quantum mechanics in terms of Schrödinger's wave mechanics. Studies evidence for quantum theory, for inadequacy of classical mechanics, 2 postulates of quantum mechanics; numerous important, fruitful applications of quantum mechanics in spectroscopy, collision problems, electrons in solids; other topics. "One of the most rewarding features . . . is the interlacing of problems with text." Amer. J. of Physics. Corrected edition. 21 illus. Index. 296pp. 5½ x 8.

\$524 Paperbound \$2.00

**PHYSICAL PRINCIPLES OF THE QUANTUM THEORY**, Werner Heisenberg. A Nobel laureate discusses quantum theory; Heisenberg's own work, Compton, Schrödinger, Wilson, Einstein, many others. Written for physicists, chemists who are not specialists in quantum theory, only elementary formulae are considered in the text; there is a mathematical appendix for specialists. Profound without sacrifice of clarity. Translated by C. Eckart, F. Hoyt. 18 figures. 192pp. 5½ x 8.

\$113 Paperbound \$1.35

# PHYSICS, HISTORIES AND CLASSICS

**A HISTORY OF PHYSICS: IN ITS ELEMENTARY BRANCHES (THROUGH 1925), INCLUDING THE EVOLUTION OF PHYSICAL LABORATORIES**, Florian Cajori. Revised and enlarged edition. The only first-rate brief history of physics. Still the best entry for a student or teacher into the antecedents of modern theories of physics. A clear, non-mathematical, handy reference work which traces in critical fashion the developments of ideas, theories, techniques, and apparatus from the Greeks to the 1920's. Within each period he analyzes the basic topics of mechanics, light, electricity and magnetism, sound, atomic theory and structure of matter, radioactivity, etc. A chapter on modern research: Curie, Kelvin, Planck's quantum theory, thermodynamics, Fitzgerald and Lorentz, special and general relativity, J. J. Thomson's model of an atom, Bohr's discoveries and later results, wave mechanics, and many other matters. Much bibliographic detail in footnotes. Index. 16 figures. xv + 424pp. 5% x 8. T970 Paperbound \$2.00

**A HISTORY OF THE MATHEMATICAL THEORIES OF ATTRACTION AND THE FIGURE OF THE EARTH: FROM THE TIME OF NEWTON TO THAT OF LAPLACE**, I. Todhunter. A technical and detailed review of the theories concerning the shape of the earth and its gravitational pull, from the earliest investigations in the seventeenth century up to the middle of the nineteenth. Some of the greatest mathematicians and scientists in history applied themselves to these questions: Newton ("Principia Mathematica"), Huygens, Maupertuis, Simpson, d'Alembert, etc. Others discussed are Poisson, Gauss, Plana, Lagrange, Bolt, and many more. Particular emphasis is placed on the theories of Laplace and Legendre, several chapters being devoted to Laplace's "Mécanique Céleste" and his memoirs, and several others to the memoirs of Legendre. Important to historians of science and mathematics and to the specialist who desires background information in the field. 2 volumes bound as 1. Index. xxxvi + 984pp. 5% x 8.

\$148 Clothbound \$7.50

**OPTICKS**, Sir Isaac Newton. In its discussions of light, reflection, color, refraction, theories of wave and corpuscular theories of light, this work is packed with scores of insights and discoveries. In its precise and practical discussion of construction of optical apparatus, contemporary understandings of phenomena it is truly fascinating to modern physicists, astronomers, mathematicians. Foreword by Albert Einstein. Preface by I. B. Cohen of Harvard University. 7 pages of portraits, facsimile pages, letters, etc. cxvi + 414pp. 5% x 8.

\$205 Paperbound \$2.25

**TREATISE ON LIGHT**, Christiaan Huygens. The famous original formulation of the wave theory of light, this readable book is one of the two decisive and definitive works in the field of light (Newton's "Optics" is the other). A scientific giant whose researches ranged over mathematics, astronomy, and physics, Huygens, in this historic work, covers such topics as rays propagated in straight lines, reflection and refraction, the spreading and velocity of light, the nature of opaque bodies, the non-spherical nature of light in the atmosphere, properties of Iceland Crystal, and other related matters. Unabridged republication of original (1912) English edition. Translated and introduced by Silvanus P. Thompson. 52 illustrations. xii + 129pp. 5% x 8.

\$179 Paperbound \$1.50

**FARADAY'S EXPERIMENTAL RESEARCHES IN ELECTRICITY**. Faraday's historic series of papers containing the fruits of years of original experimentation in electrical theory and electrochemistry. Covers his findings in a variety of areas: Induction of electric currents, Evolution of electricity from magnetism, New electrical state or condition of matter, Explication of Arago's magnetic phenomena, New law of electric conduction, Electro-chemical decomposition, Electricity of the Voltaic Pile, Static Induction, Nature of the electric force or forces, Nature of electric current, The character and direction of the electric force of the Gymnotus, Magneto-electric spark, The magnetization of light and the illumination of magnetic lines of force, The possible relation of gravity to electricity, Sub-terraneous electro-telegraph wires, Some points of magnetic philosophy, The diamagnetic conditions of flame and gases, and many other matters. Complete and unabridged republication. 3 vols. bound as 2. Originally reprinted from the Philosophical Transactions of 1831-8. Indices. Illustrations. Total of 1463pp. 5% x 8.

\$783-4, Clothbound \$17.50 (tentative)

**REFLECTIONS ON THE MOTIVE POWER OF FIRE**, Sadi Carnot, and other papers on the 2nd law of thermodynamics by E. Clapeyron and R. Clausius. Carnot's "Reflections" laid the groundwork of modern thermodynamics. Its non-technical, mostly verbal statements examine the relations between heat and the work done by heat in engines, establishing conditions for the economical working of these engines. The papers by Clapeyron and Clausius here reprinted added further refinements to Carnot's work, and led to its final acceptance by physicists. Selections from posthumous manuscripts of Carnot are also included. All papers in English. New introduction by E. Mendoza. 12 illustrations. xxii + 152pp. 5% x 8.

\$661 Paperbound \$1.50

**DIALOGUES CONCERNING TWO NEW SCIENCES**, Galileo Galilei. This classic of experimental science, mechanics, engineering, is as enjoyable as it is important. A great historical document giving insights into one of the world's most original thinkers. It is based on 30 years' experimentation. It offers a lively exposition of dynamics, elasticity, sound, ballistics, strength of materials, the scientific method. "Superior to everything else of mine," Galileo. Trans. by H. Crew, A. Salvio. 126 diagrams. Index. xxi + 288pp. 5% x 8.

\$89 Paperbound \$1.75

## Catalogue of Dover Books

**TREATISE ON ELECTRICITY AND MAGNETISM**, James Clerk Maxwell. For more than 80 years a seemingly inexhaustible source of leads for physicists, mathematicians, engineers. Total of 1082pp. on such topics as Measurement of Quantities, Electrostatics, Elementary Mathematical Theory of Electricity, Electrical Work and Energy in a System of Conductors, General Theorems, Theory of Electrical Images, Electrolysis, Conduction, Polarization, Dielectrics, Resistance, etc. "The greatest mathematical physicist since Newton," Sir James Jeans. 3rd edition. 107 figures, 21 plates. 1082pp. 5½ x 8. S636-7, 2 volume set, paperbound \$4.00

**A HISTORY OF THE THEORY OF ELASTICITY AND THE STRENGTH OF MATERIALS**, I. Todhunter and K. Pearson. For over 60 years a basic reference, unsurpassed in scope or authority. Both a history of the mathematical theory of elasticity from Galileo, Hooke, and Mariotte to Saint Venant, Kirchhoff, Clebsch, and Lord Kelvin and a detailed presentation of every important mathematical contribution during this period. Presents proofs of thousands of theorems and laws, summarizes every relevant treatise, many unavailable elsewhere. Practically a book alone is devoted to modern founders: Saint Venant, Lamé, Boussinesq, Rankine, Lord Kelvin, F. Neumann, Kirchhoff, Clebsch. Hundreds of pages of technical and physical treatises on specific applications of elasticity to particular materials. Indispensable for the mathematician, physicist, or engineer working with elasticity. Unabridged, corrected reprint of original 3-volume 1886-1893 edition. Three volume set. Two indexes. Appendix to Vol. I. Total of 2344pp. 5½ x 8½. S914-916 The set, Clothbound \$15.00

**DE MAGNETE**, William Gilbert. This classic work on magnetism founded a new science. Gilbert was the first to use the word "electricity", to recognize mass as distinct from weight, to discover the effect of heat on magnetic bodies; invent an electroscope, differentiate between static electricity and magnetism, conceive of the earth as a magnet. Written by the first great experimental scientist, this lively work is valuable not only as an historical landmark, but as the delightfully easy to follow record of a perpetually searching, ingenious mind. Translated by P. F. Mottelay. 25-page biographical memoir. 90 figures. lix + 368pp. 5½ x 8. S470 Paperbound \$2.00

## ASTRONOMY

**THE INTERNAL CONSTITUTION OF THE STARS**, Sir A. S. Eddington. Influence of this has been enormous; first detailed exposition of theory of radiative equilibrium for stellar interiors, of all available evidence for existence of diffuse matter in interstellar space. Studies quantum theory, polytropic gas spheres, mass-luminosity relations, variable stars, etc. Discussions of equations paralleled with informal exposition of intimate relationship of astrophysics with great discoveries in atomic physics, radiation. Introduction. Appendix. Index. 421pp. 5½ x 8. S563 Paperbound \$2.75

**PLANETARY THEORY**, E. W. Brown and C. A. Shook. Provides a clear presentation of basic methods for calculating planetary orbits for today's astronomer. Begins with a careful exposition of specialized mathematical topics essential for handling perturbation theory and then goes on to indicate how most of the previous methods reduce ultimately to two general calculation methods: obtaining expressions either for the coordinates of planetary positions or for the elements which determine the perturbed paths. An example of each is given and worked in detail. Corrected edition. Preface. Appendix. Index. xli + 302pp. 5½ x 8½. S1133 Paperbound \$2.25

**CANON OF ECLIPSES (CANON DER FINSTERNISSE)**, Prof. Theodor Ritter von Oppolzer. Since its original publication in 1887, this has been the standard reference and the most extensive single volume of data on the calculation of solar and lunar eclipses, past and future. A comprehensive introduction gives a full explanation of the use of the tables for the calculations of the exact dates of eclipses, etc. Data furnished for the calculation of 8,000 solar and 5,200 lunar eclipses, going back as far as 1200 B.C. and giving predictions up to the year 2161. Information is also given for partial and ring eclipses. All calculations based on Universal (Greenwich) Time. An unsurpassed reference work for astronomers, scientists engaged in space research and developments, historians, etc. Unabridged republication, with corrections. Preface to this edition by Donald Menzel and Owen Gingerich of the Harvard College Observatory. Translated by Owen Gingerich. 160 charts. lxx + 538pp. 8¾ x 11¼. S114 Clothbound \$10.00

**THEORY OF THE MOTION OF THE HEAVENLY BODIES MOVING ABOUT THE SUN IN CONIC SECTIONS**, Karl Friedrich Gauss. A landmark of theoretical astronomy by the great German scientist. Still authoritative and invaluable to the practicing astronomer. Part I develops the relations between the quantities on which the motion about the sun of the heavenly bodies depends—relations pertaining simply to position in the orbit, simply to position in space, between several places in orbit, and between several places in space. The calculation methods of Part II based on the groundwork of Part I include: determination of an orbit from 3 complete observations, from 4 observations (of which only two are complete), determination of an orbit satisfying as nearly as possible any number of observations whatever, and determination of orbits, taking into account the perturbations. Translation of "Theoria Motus" and with an appendix by C. H. Davis. Unabridged republication. Appendices and tables. 13 figures. xviii + 376pp. 6½ x 9¼. S1056 Paperbound \$2.95

## Catalogue of Dover Books

**THE GALACTIC NOVAE**, C. Payne-Gaposchkin, Prof. of Astronomy, Harvard Univ. A work that will be the standard reference source for years to come. Gathers together all the pertinent data, results recorded by countless observers of galactic novae over the centuries, in order to formulate a valid starting point for an interpretation of the nova process. Covers information and statistics on known novae, their variations in luminosity, distribution in the sky, spectral changes, etc.; symbiotic novae; frequently-recurring variables of the U Geminorum and Z Camelopardis class; supernovae; comparison of spectral changes; theories and interpretations of these phenomena, etc. "A comprehensive summary of everything that is now known about these stars," SCIENCE. Bibliographical references. Preface. Indices. 49 figures. 6 plates. 101 tables. x + 336pp. 5½ x 8¾. S1170 Paperbound \$2.45

**BINARY STARS**, R. G. Aitken. Still the definitive work in the field of double star astronomy. Written by the director of the Lick Observatory (considered the father of the modern study of binary star systems), this book sums up the results of 40 years of experience in the field, plus the work of centuries of research. Includes historical survey of major discoveries and contributions of the past, observational methods for visual binary stars, the radial velocity of a star (by Dr. J. H. Moore), eclipsing binary stars, known orbits of binary stars, some binary systems of special interest, the origin of binary stars. Much information on methods of spectrum analysis, orbit plotting, use of the telescope, and other practical matters. Useful for classroom study and advanced hobbyists, etc. Revised edition, corrected and with additional notes by Prof. J. T. Kent. New preface. 50 tables, 13 figures, 4 full-page plates. Bibliographies. Appendix. Indices. xii + 309pp. 5½ x 8¾. S1102 Paperbound \$2.00

**THE NATURE OF COMETS**, N. B. Richter. An authority on comets presents a concise, but thorough survey of the state of our present-day knowledge of comets and cometary activity. Based on over 20 years of research, this is a middle-level account that even the layman can appreciate, providing a fund of information on historical theories (from 1700 to the present); statistical research on total number of comets, orbital forms, perturbations caused by Jupiter, comet groups, etc.; the structure of a comet; comets as processes of cosmic decay; origin and formation of comets; etc. Also: a lengthy introduction on modern theories by Dr. R. A. Lyttleton, much technical data and observational material of specific comets, supplementary tables, and the like. Revised (1963) edition. Translated and revised by Arthur Beer. 69 illustrations, including 54 photographs of comets, tails, spectra. 41 tables. Bibliography. Index. xii + 221pp. S1111 Clothbound \$10.00

**CELESTIAL OBJECTS FOR COMMON TELESCOPES**, Rev. T. W. Webb. Classic handbook for the use and pleasure of the amateur astronomer. Of inestimable aid in locating and identifying thousands of celestial objects. Vol. I, The Solar System: discussions of the principle and operation of the telescope, procedures of observations and telescope-photography, spectroscopy, etc., precise location information of sun, moon, planets, meteors. Vol. II, The Stars: alphabetical listing of constellations, information on double stars, clusters, stars with unusual spectra, variables, and nebulae, etc. Nearly 4,000 objects noted. Edited and extensively revised by Margaret W. Mayall, director of the American Assn. of Variable Star Observers. New index by Mrs. Mayall giving the location of all objects mentioned in the text for Epoch 2000. New Precession Table added. New appendices on the planetary satellites, constellation names and abbreviations, and solar system data. Total of 46 illustrations. Total of xxxix + 606pp. 5½ x 8. Vol. I: T917 Paperbound \$2.25  
Vol. II: T918 Paperbound \$2.25  
Two Volume Set Paperbound \$4.50

**ASTRONOMY AND COSMOGONY**, Sir James Jeans. A modern classic which is still of enormous value to everyone in astronomy, etc., this is Jean's last and most famous exposition. The summation of a lifetime's devotion to science, it presents his final conclusions on a host of problems ranging over the whole of descriptive astronomy, astrophysics, stellar dynamics, and cosmology. Contents: The Light from the Stars, Gaseous Stars, the Source of Stellar Energy, Liquid Stars, The Evolution of the Stars, The Configuration of Rotating Masses, The Evolution of Binary Systems, The Ages of the Stars, The Great Nebulae, The Galactic Systems, Variable Stars, etc. New preface by L. Motz, Columbia U. 16 full-page photographic illustrations. xv + 428pp. 5½ x 8¾. S923 Paperbound \$2.45

**ASTRONOMY OF STELLAR ENERGY AND DECAY**, Martin Johnson. Middle level treatment of astronomy as interpreted by modern atomic physics. Part One is non-technical, examines physical properties, source of energy, spectroscopy, fluctuating stars, various models and theories, etc. Part Two parallels these topics, providing their mathematical foundation. "Clear, concise, and readily understandable," American Library Assoc. Bibliography. 3 indexes. 29 illustrations. 216pp. 5½ x 8. S537 Paperbound \$1.50

**MATHEMATICAL THEORIES OF PLANETARY MOTIONS**, Otto Džibek. Translated by Mark W. Harrington and William J. Hussey. Lucid account of the principles of mathematical astronomy. It examines that part of celestial mechanics which deals with the motions of heavenly bodies considered as material points. Contents: Solution of the Problem of Two Bodies; Formation of the General Integrals for Problem of n Bodies . . . Including discussions of elliptic, parabolic, and hyperbolic orbits, the solution of Kepler's equation, etc.; and sections headed The General Properties of the Integrals and The Theory of Perturbations . . . which deals with the theory of absolute perturbations, analytical development of the perturbing function, the variation of the elements, the secular variation of the mean longitude, etc. vi + 294pp. 5½ x 8¾. S129 Paperbound \$2.00



## Catalogue of Dover Books

**A COMPENDIUM OF SPHERICAL ASTRONOMY**, S. Newcomb. Long a standard collection of basic methods and formulas most useful to the working astronomer, and clear full text for students. Includes the most important common approximations; 40 pages on the method of least squares; general theory of spherical coordinates; parallax; aberration; astronomical refraction; theory of precession; proper motion of the stars; methods of deriving positions of stars; and much more. Index. 9 Appendices of tables, formulas, etc. 36 figures. xviii + 444pp. 5% x 8.  
\$690 Paperbound \$2.25

**PRINCIPLES OF STELLAR DYNAMICS**, S. Chandrasekhar. A leading astrophysicist here presents the theory of stellar dynamics as a branch of classical dynamics, clarifying the fundamental issues and the underlying motivations of the theory. He analyzes the effects of stellar encounters in terms of the classical 2-body problem, and investigates problems centering about Liouville's theorem and the solutions of the equations of continuity. This edition also includes 4 important papers by the author published since "Stellar Dynamics," and equally indispensable for all workers in the field: "New Methods in Stellar Dynamics" and "Dynamical Friction," Parts I, II, and III. Index. 3 Appendixes. Bibliography. 50 illustrations. x + 313pp. 5% x 8.  
\$659 Paperbound \$2.25

**AN INTRODUCTION TO THE STUDY OF STELLAR STRUCTURE**, Subrahmanyan Chandrasekhar. Outstanding treatise on stellar dynamics by one of world's greatest astrophysicists. Uses classical & modern math methods to examine relationship between loss of energy, the mass, and radius of stars in a steady state. Discusses thermodynamic laws from Carathéodory's axiomatic standpoint; adiabatic, polytropic laws; work of Ritter, Emden, Kelvin, others; Stromgren envelopes as starter for theory of gaseous stars; Gibbs statistical mechanics (quantum); degenerate stellar configuration & theory of white dwarfs, etc. "Highest level of scientific merit," BULLETIN, AMER. MATH. SOC. Bibliography. Appendixes. Index. 33 figures. 509pp. 5% x 8.  
\$413 Paperbound \$2.75

**STATISTICAL ASTRONOMY**, Robert J. Trumpler and Harold F. Weaver, University of California. Standard introduction to the principles and techniques of statistical astronomy, a field of rapidly growing importance in this space age. An extensive section, "Elements of Statistical Theory," provides the astronomer with the tools for solving problems of descriptive astronomy, observational errors, constitution of extra-galactic nebulae, etc. Procedures used in statistical astronomy are related to basic mathematical principles of statistics such as univariate distribution, integral equations, general theory of samples, etc. Other sections deal with: Statistical Description of the Galactic System; Stellar Motions in the Vicinity of the Sun; Luminosity—Spectral Type Distribution; Space Distribution of Stars; and Galactic Rotation. List of symbols. Appendix (10 tables). 2 indexes. Extensive bibliography. 31 tables. 97 figures. xxi + 644pp. 5% x 8 1/2.  
\$301 Paperbound \$3.00

**AN INTRODUCTORY TREATISE ON DYNAMICAL ASTRONOMY**, H. C. Plummer. Unusually wide connected and concise coverage of nearly every significant branch of dynamical astronomy, stressing basic principles throughout: determination of orbits, planetary theory, lunar theory, precession and nutation, and many of their applications. Hundreds of formulas and theorems worked out completely, important methods thoroughly explained. Covers motion under a central attraction, orbits of double stars and spectroscopic binaries, the libration of the moon, and much more. Index. 8 diagrams. xxi + 343pp. 5% x 8 1/2.  
\$689 Paperbound \$2.35

**AN INTRODUCTORY TREATISE ON THE LUNAR THEORY**, E. W. Brown. Indispensable for all scientists and engineers interested in orbital calculation, satellites, or navigation of space. Only work in English to explain in detail 5 major mathematical approaches to the problem of 3 bodies, those of Laplace, de Pontécoulant, Hansen, Delaunay, and Hill. Covers expressions for mutual attraction, equations of motion, forms of solution, variations of the elements in disturbed motion, the constants and their interpretations, planetary and other disturbing influences, etc. Index. Bibliography. Tables. xvi + 292pp. 5% x 8 1/2.  
\$666 Paperbound \$2.00

**SPHERICAL AND PRACTICAL ASTRONOMY**, W. Chauvenet. First book in English to apply mathematical techniques to astronomical problems is still standard work. Covers almost entire field, rigorously, with over 300 examples worked out. Vol. 1, spherical astronomy, applications to nautical astronomy; determination of hour angles, parallactic angle for known stars; interpolation; parallax; laws of refraction; predicting eclipses; precession, nutation of fixed stars; etc. Vol. 2, theory, use, of instruments; telescope; measurement of arcs, angles in general; electro-chronograph; sextant, reflecting circles; zenith telescope; etc. 100-page appendix of detailed proof of Gauss' method of least squares. 5th revised edition. Index. 15 plates, 20 tables. 1340pp. 5% x 8.  
Vol. 1 \$618 Paperbound \$2.75  
Vol. 2 \$619 Paperbound \$2.75  
The set \$5.50

**RADIATIVE TRANSFER**, S. Chandrasekhar. Definitive work in field provides foundation for analysis of stellar atmospheres, planetary illumination, sky radiation; to physicists, a study of problems analogous to those in theory of diffusion of neutrons. Partial contents: equation of transfer, isotropic scattering, H-functions, diffuse reflection and transmission, Rayleigh scattering, X, Y functions, radiative equilibrium of stellar atmospheres. Extensive bibliography. 3 appendices. 35 tables. 35 figures. 407pp. 5% x 8 1/2.  
\$599 Paperbound \$2.25

## ENGINEERING AND TECHNOLOGY

### General and mathematical

**ENGINEERING MATHEMATICS**, Kenneth S. Miller. A text for graduate students of engineering to strengthen their mathematical background in differential equations, etc. Mathematical steps very explicitly indicated. Contents: Determinants and Matrices, Integrals, Linear Differential Equations, Fourier Series and Integrals, Laplace Transform, Network Theory, Random Function . . . all vital requisites for advanced modern engineering studies. Unabridged republication. Appendices: Borel Sets; Riemann-Stieltjes Integral; Fourier Series and Integrals. Index. References at Chapter Ends. xli + 417pp. 6 x 8½. \$1121 Paperbound \$2.00

**MATHEMATICAL ENGINEERING ANALYSIS**, Rufus Oldenburger. A book designed to assist the research engineer and scientist in making the transition from physical engineering situations to the corresponding mathematics. Scores of common practical situations found in all major fields of physics are supplied with their correct mathematical formulations—applications to automobile springs and shock absorbers, clocks, throttle torque of diesel engines, resistance networks, capacitors, transmission lines, microphones, neon tubes, gasoline engines, refrigeration cycles, etc. Each section reviews basic principles of underlying various fields: mechanics of rigid bodies, electricity and magnetism, heat, elasticity, fluid mechanics, and aerodynamics. Comprehensive and eminently useful. Index. 169 problems, answers. 200 photos and diagrams. xiv + 426pp. 5½ x 8½. \$919 Paperbound \$2.50

**MATHEMATICS OF MODERN ENGINEERING**, E. G. Keller and R. E. Doherty. Written for the Advanced Course in Engineering of the General Electric Corporation, deals with the engineering use of determinants, tensors, the Heaviside operational calculus, dyadics, the calculus of variations, etc. Presents underlying principles fully, but purpose is to teach engineers to deal with modern engineering problems, and emphasis is on the perennial engineering attack of set-up and solve. Indexes. Over 185 figures and tables. Hundreds of exercises, problems, and worked-out examples. References. Two volume set. Total of xxxiii + 623pp. 5½ x 8.

\$734 Vol I Paperbound \$1.85  
\$735 Vol II Paperbound \$1.85  
The set \$3.70

**MATHEMATICAL METHODS FOR SCIENTISTS AND ENGINEERS**, L. P. Smith. For scientists and engineers, as well as advanced math students. Full investigation of methods and practical description of conditions under which each should be used. Elements of real functions, differential and integral calculus, space geometry, theory of residues, vector and tensor analysis, series of Bessel functions, etc. Each method illustrated by completely-worked-out examples, mostly from scientific literature. 368 graded unsolved problems. 100 diagrams. x + 453pp. 5½ x 8½. \$220 Paperbound \$2.00

**THEORY OF FUNCTIONS AS APPLIED TO ENGINEERING PROBLEMS**, edited by R. Rothe, F. Ollendorf, and K. Pohlhausen. A series of lectures given at the Berlin Institute of Technology that shows the specific applications of function theory in electrical and allied fields of engineering. Six lectures provide the elements of function theory in a simple and practical form, covering complex quantities and variables, integration in the complex plane, residue theorems, etc. Then 5 lectures show the exact uses of this powerful mathematical tool, with full discussions of problem methods. Index. Bibliography. 108 figures. x + 189pp. 5½ x 8.

\$733 Paperbound \$1.35

### Aerodynamics and hydrodynamics

**AIRPLANE STRUCTURAL ANALYSIS AND DESIGN**, E. E. Sechler and L. G. Dunn. Systematic authoritative book which summarizes a large amount of theoretical and experimental work on structural analysis and design. Strong on classical subsonic material still basic to much aeronautic design . . . remains a highly useful source of information. Covers such areas as layout of the airplane, applied and design loads, stress-strain relationships for stable structures, truss and frame analysis, the problem of instability, the ultimate strength of stiffened flat sheet, analysis of cylindrical structures, wings and control surfaces, fuselage analysis, engine mounts, landing gears, etc. Originally published as part of the CALCUT Aeronautical Series. 256 illustrations. 47 study problems. Indexes. xi + 420pp. 5½ x 8½. \$1043 Paperbound \$2.25

**FUNDAMENTALS OF HYDRO- AND AEROMECHANICS**, L. Prandtl and O. G. Tietjens. The well-known standard work based upon Prandtl's lectures at Goettingen. Wherever possible hydrodynamics theory is referred to practical considerations in hydraulics, with the view of unifying theory and experience. Presentation is extremely clear and though primarily physical, mathematical proofs are rigorous and use vector analysis to a considerable extent. An Engineering Society Monograph, 1934. 186 figures. Index. xvi + 270pp. 5½ x 8.

\$374 Paperbound \$1.85

## Catalogue of Dover Books

**FLUID MECHANICS FOR HYDRAULIC ENGINEERS**, H. Rouse. Standard work that gives a coherent picture of fluid mechanics from the point of view of the hydraulic engineer. Based on courses given to civil and mechanical engineering students at Columbia and the California Institute of Technology, this work covers every basic principle, method, equation, or theory of interest to the hydraulic engineer. Much of the material, diagrams, charts, etc., in this self-contained text are not duplicated elsewhere. Covers irrotational motion, conformal mapping, problems in laminar motion, fluid turbulence, flow around immersed bodies, transportation of sediment, general characteristics of wave phenomena, gravity waves in open channels, etc. Index. Appendix of physical properties of common fluids. Frontispiece + 245 figures and photographs. xvi + 422pp. 5½ x 8. \$729 Paperbound \$2.25

**WATERHAMMER ANALYSIS**, John Parmakian. Valuable exposition of the graphical method of solving waterhammer problems by Assistant Chief Designing Engineer, U.S. Bureau of Reclamation. Discussions of rigid and elastic water column theory, velocity of waterhammer waves, theory of graphical waterhammer analysis for gate operation, closings, openings, rapid and slow movements, etc., waterhammer in pump discharge caused by power failure, waterhammer analysis for compound pipes, and numerous related problems. "With a concise and lucid style, clear printing, adequate bibliography and graphs for approximate solutions at the project stage, it fills a vacant place in waterhammer literature." **WATER POWER**. 43 problems. Bibliography. Index. 113 illustrations. xiv + 161pp. 5½ x 8½. \$1061 Paperbound \$1.65

**AERODYNAMIC THEORY: A GENERAL REVIEW OF PROGRESS**, William F. Durand, editor-in-chief. A monumental joint effort by the world's leading authorities prepared under a grant of the Guggenheim Fund for the Promotion of Aeronautics. Intended to provide the student and aeronautic designer with the theoretical and experimental background of aeronautics. Never equalled for breadth, depth, reliability. Contains discussions of special mathematical topics not usually taught in the engineering or technical courses. Also: an extended two-part treatise on Fluid Mechanics, discussions of aerodynamics of perfect fluids, analyses of experiments with wind tunnels, applied airfoil theory, the non-lifting system of the airplane, the air propeller, hydrodynamics of boats and floats, the aerodynamics of cooling, etc. Contributing experts include Munk, Giacomelli, Prandtl, Toussaint, Von Karman, Klemperer, among others. Unabridged republication. 6 volumes bound as 3. Total of 1,012 figures, 12 plates. Total of 2,186pp. Bibliographies. Notes. Indices. 5½ x 8. \$328-\$330 Paperbound The Set \$13.50

**APPLIED HYDRO- AND AEROMECHANICS**, L. Prandtl and O. G. Tietjens. Presents, for the most part, methods which will be valuable to engineers. Covers flow in pipes, boundary layers, airfoil theory, entry conditions, turbulent flow in pipes, and the boundary layer, determining drag from measurements of pressure and velocity, etc. "Will be welcomed by all students of aerodynamics." **NATURE**. Unabridged, unaltered. An Engineering Society Monograph, 1934. Index. 226 figures, 28 photographic plates illustrating flow patterns. xvi + 311pp. 5½ x 8. \$375 Paperbound \$2.00

**SUPERSONIC AERODYNAMICS**, E. R. C. Miles. Valuable theoretical introduction to the supersonic domain, with emphasis on mathematical tools and principles, for practicing aerodynamicists and advanced students in aeronautical engineering. Covers fundamental theory, divergence theorem and principles of circulation, compressible flow and Helmholtz laws, the Prandtl-Busemann graphic method for 2-dimensional flow, oblique shock waves, the Taylor-Maccoll method for cones in supersonic flow, the Chaplygin method for 2-dimensional flow, etc. Problems range from practical engineering problems to development of theoretical results. "Rendered outstanding by the unprecedented scope of its contents . . . has undoubtedly filled a vital gap." **AERONAUTICAL ENGINEERING REVIEW**. Index. 173 problems, answers. 106 diagrams. 7 tables. xii + 255pp. 5½ x 8. \$214 Paperbound \$1.45

**HYDRAULIC TRANSIENTS**, G. R. Rich. The best text in hydraulics ever printed in English . . . by one of America's foremost engineers (former Chief Design Engineer for T.V.A.). Provides a transition from the basic differential equations of hydraulic transient theory to the arithmetic integration computation required by practicing engineers. Sections cover Water Hammer, Turbine Speed Regulation, Stability of Governing, Water-Hammer Pressures in Pump Discharge Lines, The Differential and Restricted Orifice Surge Tanks, The Normalized Surge Tank Charts of Calame and Gaden, Navigation Locks, Surges in Power Canals—Tidal Harmonics, etc. Revised and enlarged. Author's prefaces. Index. xiv + 409pp. 5½ x 8½. \$116 Paperbound \$2.50

**HYDRAULICS AND ITS APPLICATIONS**, A. H. Gibson. Excellent comprehensive textbook for the student and thorough practical manual for the professional worker, a work of great stature in its area. Half the book is devoted to theory and half to applications and practical problems met in the field. Covers modes of motion of a fluid, critical velocity, viscous flow, eddy formation, Bernoulli's theorem, flow in converging passages, vortex motion, form of effluent streams, notches and weirs, skin friction, losses at valves and elbows, siphons, erosion of channels, jet propulsion, waves of oscillation, and over 100 similar topics. Final chapters (nearly 400 pages) cover more than 100 kinds of hydraulic machinery: Pelton wheel, speed regulators, the hydraulic ram, surge tanks, the scoop wheel, the Venturi meter, etc. A special chapter treats methods of testing theoretical hypotheses: scale models of rivers, tidal estuaries, siphon spillways, etc. 5th revised and enlarged (1952) edition. Index. Appendix. 427 photographs and diagrams. 95 examples, answers. xv + 813pp. 6 x 9. \$791 Clothbound \$8.00

# PUZZLES AND RECREATIONS

**ATHEMATICS**, Henry Ernest Dudeney. The foremost British originator of this always intriguing, witty, and paradoxical classic, one of the most mathematical amusements. More than 430 puzzles, problems, and games, problems on number manipulation, uncursal and other route measuring, weighing, packing, age, kinship, chessboards, joining, dissection, and many others. Solutions. More than 450 illustrations. 7473 Paperbound \$1.25

**THE GAME OF LOGIC**, Lewis Carroll. "Symbolic Logic" is not concerned with logic, but is instead a collection of over 380 problems posed with the syllogism, and a fascinating diagrammatic method of drawing "The Game of Logic," Carroll's whimsical imagination devises a logical diagrams and counters (included) to manipulate hundreds of tricky syllogisms. "Hit or Miss" is a lagniappe of 101 additional puzzles in the same manner. Until this reprint edition, both of these books were rarities costing \$15.00. Symbolic Logic: Index, xxxi + 199pp. The Game of Logic: 96pp. Two 5 1/2" x 8". T492 Paperbound \$1.50

**NTHS: A BOOK OF PUZZLES**, W. Shepherd. Mazes, formerly associated with are still among the most intriguing of intellectual puzzles. This is a collection of 50 amusements that embody the principle of the maze: a tradition; 3-dimensional, ribbon, and Möbius-strip mazes; hidden messages; etc.—almost all built on amusing story situations. 84 illustrations. Solutions. xv + 122pp. 5 3/8 x 8. 1731 Paperbound \$1.00

**CREATIONS, M. Kralchik.** Some 250 puzzles, problems, demonstrations of  
for beginners & advanced mathematicians. Unusual historical prob-  
lem for interval, Arabic, Hindu sources: modern problems based on "mathematics  
geometry, topology, arithmetic, etc. Pastimes derived from figurative  
numbers, Fermat numbers; fairy chess, latrunculus, reversal, many topics.  
ellent for insights into special fields of math. 181 illustrations. 330pp.  
1163 Paperbound \$1.75

**PZZLES OF SAM LOYD, Vol. I**, selected and edited by M. Gardner. Puzzles created and innovator. Selected from his famous "Cyclopedia of the Unique style and historical flavor of the originals. There are posers in the algebra, probability, game theory, routing, topology, counting, etc., research, geographical, and on. Includes his famous "15" variations, a national craze, and his "Horse of a Different Color" which sold millions. His most ingenious puzzles in all, 120 line drawings and diagrams. Solutions. xx + 167pp. 5 3/8 x 8. 1498 Paperbound \$1.00

**IN MATHEMATICS**, Hubert Phillips ("Calliban"). Calliban is generally considered the modern problemist. Here are 100 of his best and wittiest puzzles, **authored** by himself from such publications as the London Daily Telegraph, and **intended** to put even the sharpest puzzle detective through his paces. **Permeant** of clear thinking and a logical mind. Complete solutions are provided.  $x + 107pp.$  5%  $x 8\frac{1}{2}$ . T91 Paperbound \$1.00

**IN LOGIC AND REASONING**, H. Phillips ("Callban"). 100 choice, hitherto  
by England's best-known problemist. No special knowledge needed to  
or inferential problems, just an unclouded mind, nerves of steel, and  
presented are both necessary and just sufficient to allow one unambiguous  
n 30 different types of puzzles, all ingenious and varied, many one of a  
challenge the expert, please the beginner. Original publication. 100 puzzles,  
T 107pp. 5 1/2 x 8 1/2. T19 Paperbackbound \$1.00

**PUZZLES FOR BEGINNERS AND ENTHUSIASTS**, G. Mott-Smith. 188 mathematical puzzles. Mental agility. Inference, Interpretation, algebra, dissection of plane figures, use of numbers, decimation, permutations, probability, all enter these puzzles. Puzzles like the Odic Force, How to Draw an Ellipse, Spider's Cousin, etc. Detailed solutions. Appendix with square roots, triangular numbers, etc. Illustrations. 2nd revised edition. 248pp. 5 3/4" x 8". \$1.98 Paperbound \$1.00

**MAGIC AND MYSTERY**, Martin Gardner. Card tricks, feats of mental mathematics, other "magic" explained as applications of probability, sets, topology, various branches of mathematics. Creative examination of laws of probability with scores of new tricks and insights. 115 sections discuss tricks of cards, coins, geometrical vanishing tricks, dozens of others. No sleight of hand. Success guaranteed. 115 illustrations. xii + 174pp. 5% x 8. T335 Paperbound \$1.00

## Catalogue of Dover Books

**RECREATIONS IN THE THEORY OF NUMBERS: THE QUEEN OF MATHEMATICS ENTERTAINS**, Albert H. Belfer. The theory of numbers is often referred to as the "Queen of Mathematics." In this book Mr. Belfer has compiled the first English volume to deal exclusively with the recreational aspects of number theory, an inherently recreational branch of mathematics. The author's clear style makes for enjoyable reading as he deals with such topics as: perfect numbers, amicable numbers, Fermat's theorem, Wilson's theorem, polygonal and figurate numbers, methods of factoring, primitive roots, Euler's function, polygonal and figurate numbers, Mersenne numbers, congruence, repeating decimals, etc. Countless puzzle problems, with full answers and explanations. For mathematicians and mathematically-inclined laymen, etc. New publication. 28 figures. 9 illustrations. 103 tables. Bibliography at chapter ends. vi + 247pp. 5% x 8½. T1096 Paperbound \$1.85

**PAPER FOLDING FOR BEGINNERS**, W. D. Murray and F. J. Rigney. A delightful introduction to the varied and entertaining Japanese art of origami (paper folding), with a full crystal-clear text that anticipates every difficulty; over 275 clearly labeled diagrams of all important stages in creation. You get results at each stage, since complex figures are logically developed from simpler ones. 43 different pieces are explained: place mats, drinking cups, bonbon boxes, sailboats, frogs, roosters, etc. 6 photographic plates. 279 diagrams. 95pp. 5% x 8½. T713 Paperbound \$1.00

**1800 RIDDLES, ENIGMAS AND CONUNDRUMS**, Darwin A. Hindman. Entertaining collection ranging from hilarious gags to outrageous puns to sheer nonsense—a welcome respite from sophisticated humor. Children, toastmasters, and practically anyone with a funny bone will find these zany riddles tickling and eminently repeatable. Sample: "Why does Santa Claus always go down the chimney?" "Because it suits him." Some old, some new—covering a wide variety of subjects. New publication. iii + 154pp. 5% x 8½. T1059 Paperbound \$1.00

**EASY-TO-DO ENTERTAINMENTS AND DIVERSIONS WITH CARDS, STRING, COINS, PAPER AND MATCHES**, R. M. Abraham. Over 300 entertaining games, tricks, puzzles, and pastimes for children and adults. Invaluable to anyone in charge of groups of youngsters for party givers, etc. Contains sections on card tricks and games, making things by paperfolding—toys, decorations, and the like; tricks with coins, matches, and pieces of string; descriptions of games; toys that can be made from common household objects; mathematical recreations; word games; and 50 miscellaneous entertainments. Formerly "Winter Nights Entertainments." Introduction by Lord Baden Powell. 329 illustrations. v + 186pp. 5% x 8. T921 Paperbound \$1.00

**DIVERSIONS AND PASTIMES WITH CARDS, STRING, PAPER AND MATCHES**, R. M. Abraham. Another collection of amusements and diversion for game and puzzle fans of all ages. Many new paperfolding ideas and tricks, an extensive section on amusements with knots and splices, two chapters of easy and not-so-easy problems, coin and match tricks, and lots of other parlor pastimes from the agile mind of the late British problemist and gamester. Corrected and revised version. Illustrations. 160pp. 5% x 8½. T1127 Paperbound \$1.00

**STRING FIGURES AND HOW TO MAKE THEM: A STUDY OF CAT'S-CRADLE IN MANY LANDS**, Caroline Furness Jayne. In a simple and easy-to-follow manner, this book describes how to make 107 different string figures. Not only is looping and crossing string between the fingers a common youthful diversion, but it is an ancient form of amusement practiced in all parts of the globe, especially popular among primitive tribes. These games are fun for all ages and offer an excellent means for developing manual dexterity and coordination. Much insight also for the anthropological observer on games and diversions in many different cultures. Index. Bibliography. Introduction by A. C. Haddon, Cambridge University. 17 full-page plates. 950 illustrations. xxiii + 407pp. 5% x 8½. T152 Paperbound \$2.00

**CRYPTANALYSIS**, Helen F. Gaines. (Formerly **ELEMENTARY CRYPTANALYSIS**.) A standard elementary and intermediate text for serious students. It does not confine itself to old material, but contains much that is not generally known, except to experts. Concealment, Transposition, Substitution ciphers; Vigenere, Kasiski, Playfair, multifid, dozens of other techniques. Appendix with sequence charts, letter frequencies in English, 5 other languages, English word frequencies. Bibliography. 167 codes. New to this edition: solution to codes. vi + 230pp. 5% x 8. T97 Paperbound \$2.00

**MAGIC SQUARES AND CUBES**, W. S. Andrews. Only book-length treatment in English, a thorough non-technical description and analysis. Here are nasik, overlapping, pandiagonal, serated squares; magic circles, cubes, spheres, rhombuses. Try your hand at 4-dimensional magical figures! Much unusual folklore and tradition included. High school algebra is sufficient. 754 diagrams and illustrations. viii + 419pp. 5% x 8. T658 Paperbound \$1.85

**CALIBAN'S PROBLEM BOOK: MATHEMATICAL, INFERENTIAL, AND CRYPTOGRAPHIC PUZZLES**, H. Phillips ("Caliban"), S. T. Shovelton, G. S. Marshall. 105 ingenious problems by the greatest living creator of puzzles based on logic and inference. Rigorous, modern, pliant, and reflecting the author's unusual personality, these intermediate and advanced puzzles all involve the ability to reason clearly through complex situations; some call for mathematical knowledge, ranging from algebra to number theory. Solutions. xi + 180pp. 5% x 8. T736 Paperbound \$1.25

# FICTION

**THE LAND THAT TIME FORGOT and THE MOON MAID, Edgar Rice Burroughs.** In the opinion of many, Burroughs' best work. The first concerns a strange island where evolution is individual rather than phylogenetic. Speechless anthropoids develop into intelligent human beings within a single generation. The second projects the reader far into the future and describes the first voyage to the Moon (in the year 2025), the conquest of the Earth by the Moon, and years of violence and adventure as the enslaved Earthmen try to regain possession of their planet. "An imaginative tour de force that keeps the reader keyed up and expectant," *NEW YORK TIMES*. Complete, unabridged text of the original two novels (three parts in each). 5 illustrations by J. Allen St. John. vi + 552pp. 5½ x 8½.

T1020 Clothbound \$3.75  
T358 Paperbound \$2.00

**AT THE EARTH'S CORE, PELLUCIDAR, TANAR OF PELLUCIDAR: THREE SCIENCE FICTION NOVELS BY EDGAR RICE BURROUGHS.** Complete, unabridged texts of the first three Pellucidar novels. Tales of derring-do by the famous master of science fiction. The locale for these three related stories is the inner surface of the hollow Earth where we discover the world of Pellucidar, complete with all types of bizarre, menacing creatures, strange peoples, and alluring maidens—guaranteed to delight all Burroughs fans and a wide circle of adventure lovers. Illustrated by J. Allen St. John and P. F. Berdanier. vi + 433pp. 5½ x 8½.

T1051 Paperbound \$2.00

**THE PIRATES OF VENUS and LOST ON VENUS: TWO VENUS NOVELS BY EDGAR RICE BURROUGHS.** Two related novels, complete and unabridged. Exciting adventure on the planet Venus with Earthman Carson Napier broken-field running through one dangerous episode after another. All lovers of swashbuckling science fiction will enjoy these two stories set in a world of fascinating societies, fierce beasts, 5000-ft. trees, lush vegetation, and wide seas. Illustrations by Fortunino Matania. Total of vi + 340pp. 5½ x 8½.

T1053 Paperbound \$1.75

**A PRINCESS OF MARS and A FIGHTING MAN OF MARS: TWO MARTIAN NOVELS BY EDGAR RICE BURROUGHS.** "Princess of Mars" is the very first of the great Martian novels written by Burroughs, and it is probably the best of them all; it set the pattern for all of his later fantasy novels and contains a thrilling cast of strange peoples and creatures and the formula of Olympian heroism amidst ever-fluctuating fortunes which Burroughs carries off so successfully. "Fighting Man" returns to the same scenes and cities—many years later. A mad scientist, a degenerate dictator, and an indomitable defender of the right clash—with the fate of the Red Planet at stake! Complete, unabridged reprinting of original editions. Illustrations by F. E. Schoonover and Hugh Hutton. v + 356pp. 5½ x 8½.

T1140 Paperbound \$1.75

**THREE MARTIAN NOVELS, Edgar Rice Burroughs.** Contains: Thuvia, Maid of Mars; The Chessmen of Mars; and The Master Mind of Mars. High adventure set in an imaginative and intricate conception of the Red Planet. Mars is peopled with an intelligent, heroic human race which lives in densely populated cities and with fierce barbarians who inhabit dead sea bottoms. Other exciting creatures abound amidst an inventive framework of Martian history and geography. Complete unabridged reprintings of the first edition. 16 illustrations by J. Allen St. John. vi + 499pp. 5½ x 8½.

T39 Paperbound \$1.85

**THREE PROPHETIC NOVELS BY H. G. WELLS, edited by E. F. Bleiler.** Complete texts of "When the Sleeper Wakes" (1st book printing in 50 years), "A Story of the Days to Come," "The Time Machine" (1st complete printing in book form). Exciting adventures in the future are as enjoyable today as 50 years ago when first printed. Predict TV, movies, intercontinental airplanes, prefabricated houses, air-conditioned cities, etc. First important author to foresee problems of mind control, technological dictatorships. "Absolute best of imaginative fiction," *N. Y. Times*. Introduction. 335pp. 5½ x 8.

T605 Paperbound \$1.50

**28 SCIENCE FICTION STORIES OF H. G. WELLS.** Two full unabridged novels, *MEN LIKE GODS* and *STAR BEGOTTEN*, plus 26 short stories by the master science-fiction writer of all time. Stories of space, time, invention, exploration, future adventure—an indispensable part of the library of everyone interested in science and adventure. **PARTIAL CONTENTS:** Men Like Gods, The Country of the Blind, In the Abyss, The Crystal Egg, The Man Who Could Work Miracles, A Story of the Days to Come, The Valley of Spiders, and 21 more! 928pp. 5½ x 8.

T265 Clothbound \$4.50

**THE WAR IN THE AIR, IN THE DAYS OF THE COMET, THE FOOD OF THE GODS: THREE SCIENCE FICTION NOVELS BY H. G. WELLS.** Three exciting Wells offerings bearing on vital social and philosophical issues of his and our own day. Here are tales of air power, strategic bombing, East vs. West, the potential miracles of science, the potential disasters from outer space, the relationship between scientific advancement and moral progress, etc. First reprinting of "War in the Air" in almost 50 years. An excellent sampling of Wells at his storytelling best. Complete, unabridged reprintings. 16 illustrations. 645pp. 5½ x 8½.

T1135 Paperbound \$2.00

## *Catalogue of Dover Books*

**SEVEN SCIENCE FICTION NOVELS, H. G. Wells.** Full unabridged texts of 7 science-fiction novels of the master. Ranging from biology, physics, chemistry, astronomy to sociology and other studies, Mr. Wells extrapolates whole worlds of strange and intriguing character. "One will have to go far to match this for entertainment, excitement, and sheer pleasure . . ." *NEW YORK TIMES*. Contents: *The Time Machine*, *The Island of Dr. Moreau*, *The First Men in the Moon*, *The Invisible Man*, *The War of the Worlds*, *The Food of the Gods*, *In the Days of the Comet*. 1015pp. 5½ x 8. T264 Clothbound \$4.50

**BEST GHOST STORIES OF J. S. LE FANU**, Selected and Introduced by E. F. Bleiler. LeFanu is deemed the greatest name in Victorian supernatural fiction. Here are 16 of his best horror stories, including 2 nouvelles: "Carmilla," a classic vampire tale couched in a perverse eroticism, and "The Haunted Baronet." Also: "Sir Toby's Will," "Green Tea," "Schalken the Painter," "Ullor de Lacy," "The Familiar," etc. The first American publication of about half of this material: a long-overdue opportunity to get a choice sampling of LeFanu's work. New selection (1964). 8 illustrations. 5¾ x 8¾. T415 Paperbound \$1.85

**THE WONDERFUL WIZARD OF OZ, L. F. Baum.** Only edition in print with all the original W. W. Denslow illustrations in full color—as much a part of "The Wizard" as Tenniel's drawings are for "Alice in Wonderland." "The Wizard" is still America's best-loved fairy tale, in which, as the author expresses it, "The wonderment and joy are retained and the heartaches and nightmares left out." Now today's young readers can enjoy every word and wonderful picture of the original book. New introduction by Martin Gardner. A Baum bibliography. 23 full-page color plates. viii + 268pp. 5¾ x 8. T691 Paperbound \$1.50

**GHOST AND HORROR STORIES OF AMBROSE BIERCE**, Selected and Introduced by E. F. Bleiler. 24 morbid, eerie tales—the cream of Bierce's fiction output. Contains such memorable pieces as "The Moonlit Road," "The Damned Thing," "An Inhabitant of Carcosa," "The Eyes of the Panther," "The Famous Gilson Bequest," "The Middle Toe of the Right Foot," and other chilling stories, plus the essay, "Visions of the Night" in which Bierce gives us a kind of rationale for his aesthetic of horror. New collection (1964). xxii + 199pp. 5¾ x 8¾. T767 Paperbound \$1.00

## **HUMOR**

**MR. DOOLEY ON IVRYTHING AND IVRYBODY**, Finley Peter Dunne. Since the time of his appearance in 1893, "Mr. Dooley," the fictitious Chicago bartender, has been recognized as America's most humorous social and political commentator. Collected in this volume are 102 of the best Dooley pieces—all written around the turn of the century, the height of his popularity. Mr. Dooley's Irish brogue is employed wittily and penetratingly on subjects which are just as fresh and relevant today as they were then: corruption and hypocrisy of politicians, war preparations and chauvinism, automation, Latin American affairs, superbombs, etc. Other articles range from Rudyard Kipling to football. Selected with an introduction by Robert Hutchinson. xii + 244pp. 5¾ x 8¾. T626 Paperbound \$1.00

**RUTHLESS RHYMES FOR HEARTLESS HOMES and MORE RUTHLESS RHYMES FOR HEARTLESS HOMES**, Harry Graham ("Col. D. Streamer"). A collection of Little Willy and 48 other poetic "disasters." Graham's funniest and most disrespectful verse, accompanied by original illustrations. Nonsensical, wry humor which employs stern parents, careless nurses, uninhibited children, practical jokers, single-minded golfers, Scottish lairds, etc. in the leading roles. A precursor of the "sick joke" school of today. This volume contains, bound together for the first time, two of the most perennially popular books of humor in England and America. T930 Paperbound 75¢

**A WHIMSEY ANTHOLOGY**, Collected by Carolyn Wells. 250 of the most amusing rhymes ever written. Acrostics, anagrams, palindromes, alphabetical jingles, tongue twisters, echo verses, alliterative verses, riddles, mnemonic rhymes, interior rhymes, over 40 limericks, etc. by Lewis Carroll, Edward Lear, Joseph Addison, W. S. Gilbert, Christina Rossetti, Chas. Lamb, James Boswell, Hood, Dickens, Swinburne, Leigh Hunt, Harry Graham, Poe, Eugene Field, and many others. xiv + 221pp. 5¾ x 8¾. T195 Paperbound \$1.25

**MY PIOUS FRIENDS AND DRUNKEN COMPANIONS and MORE PIOUS FRIENDS AND DRUNKEN COMPANIONS**, Songs and ballads of Conviviality Collected by Frank Shay. Magnificently illuminated by John Held, Jr. 132 ballads, blues, vaudeville numbers, drinking songs, cowboy songs, sea chanties, comedy songs, etc. of the Naughty Nineties and early 20th century. Over a third are reprinted with music. Many perennial favorites such as: *The Band Played On*, *Frankie and Johnnie*, *The Old Gray Mare*, *The Face on the Bar-room Floor*, etc. Many others unlocatable elsewhere: *The Dog-Catcher's Child*, *The Cannibal Maiden*, *Don't Go in the Lion's Cage Tonight*, *Mother*, etc. Complete verses and introductions to songs. Unabridged republication of first editions, 2 indexes (song titles and first lines and choruses). Introduction by Frank Shay. 2 volumes bounds as 1. Total of xvi + 235pp. 5¾ x 8¾. T946 Paperbound \$1.25

## Catalogue of Dover Books

**MAX AND MORITZ.** Wilhelm Busch. Edited and annotated by H. Arthur Klein. Translated by H. Arthur Klein, M. C. Klein, and others. The mischievous high jinks of Max and Moritz, Peter and Paul, Ker and Plunk, etc. are delightfully captured in sketch and rhyme. (Companion volume to "Hypocritical Helena.") In addition to the title piece, it contains: Ker and Plunk; Two Dogs and Two Boys; The Egghead and the Two Cut-ups of Corinth; Deceitful Henry; The Boys and the Pipe; Cat and Mouse; and others. (Original German text with accompanying English translations.) Afterword by H. A. Klein. vi + 216pp. 5½ x 8½.  
T181 Paperbound \$1.15

**THROUGH THE ALIMENTARY CANAL WITH GUN AND CAMERA: A FASCINATING TRIP TO THE INTERIOR.** Personally Conducted by George S. Chappell. In mock-travelogue style, the amusing account of an imaginative journey down the alimentary canal. The "explorers" enter the esophagus, round the Adam's Apple, narrowly escape from a fierce Amoeba, struggle through the Impenetrable Nerve Forests of the Lumbar Region, etc. Illustrated by the famous cartoonist, Otto Soglow, the book is as much a brilliant satire of academic pomposity and professional travel literature as it is a clever use of the facts of physiology for supremely comic purposes. Preface by Robert Benchley. Author's Foreword. 1 Photograph. 17 Illustrations by O. Soglow. xli + 114pp. 5½ x 8½.  
T376 Paperbound \$1.00

**THE BAD CHILD'S BOOK OF BEASTS, MORE BEASTS FOR WORSE CHILDREN, and A MORAL ALPHABET.** H. Belloc. Hardly an anthology of humorous verse has appeared in the last 50 years without at least a couple of these famous nonsense verses. But one must see the entire volumes—with all the delightful original illustrations by Sir Basil Blackwood—to appreciate fully Belloc's charming and witty verses that play so subacutely on the platitudes of life and morals that beset his day—and ours. A great humor classic. Three books in one. Total of 157pp. 5½ x 8.  
T749 Paperbound \$1.00

**THE DEVIL'S DICTIONARY.** Ambrose Bierce. Sardonic and irreverent barbs puncturing the pomposities and absurdities of American politics, business, religion, literature, and arts, by the country's greatest satirist in the classic tradition. Epigrammatic as Shaw, piercing as Swift, American as Mark Twain, Will Rogers, and Fred Allen. Bierce will always remain the favorite of a small coterie of enthusiasts, and of writers and speakers whom he supplies with "some of the most gorgeous witticisms of the English language." (H. L. Mencken) Over 1000 entries in alphabetical order. 144pp. 5½ x 8.  
T487 Paperbound \$1.00

**THE COMPLETE NONSENSE OF EDWARD LEAR.** This is the only complete edition of this master of gentle madness available at a popular price. A BOOK OF NONSENSE, NONSENSE SONGS, MORE NONSENSE SONGS AND STORIES in their entirety with all the old favorites that have delighted children and adults for years. The Dong With A Luminous Nose, The Jumbles, The Owl and the Pussycat, and hundreds of other bits of wonderful nonsense. 214 limericks, 3 sets of Nonsense Botany, 5 Nonsense Alphabets. 546 drawings by Lear himself, and much more. 320pp. 5½ x 8.  
T167 Paperbound \$1.00

**SINGULAR TRAVELS, CAMPAIGNS, AND ADVENTURES OF BARON MUNCHAUSEN.** R. E. Raspe, with 90 illustrations by Gustave Doré. The first edition in over 150 years to reestablish the deeds of the Prince of Liars exactly as Raspe first recorded them in 1785—the genuine Baron Munchausen, one of the most popular personalities in English literature. Included also are the best of the many sequels, written by other hands. Introduction on Raspe by J. Carswell. Bibliography of early editions. xiv + 192pp. 5½ x 8.  
T698 Paperbound \$1.00

**HOW TO TELL THE BIRDS FROM THE FLOWERS.** R. W. Wood. How not to confuse a carrot with a parrot, a grape with an ape, a puffin with nuffin. Delightful drawings, clever puns, absurd little poems point out farfetched resemblances in nature. The author was a leading physicist. Introduction by Margaret Wood White. 106 illus. 60pp. 5½ x 8.  
T523 Paperbound 75¢

**JOE MILLER'S JESTS OR, THE WITS VADE-MECUM.** The original Joe Miller jest book. Gives a keen and pungent impression of life in 18th-century England. Many are somewhat on the bawdy side and they are still capable of provoking amusement and good fun. This volume is a facsimile of the original "Joe Miller" first published in 1739. It remains the most popular and influential humor book of all time. New introduction by Robert Hutchinson. xxi + 70pp. 5½ x 8½.  
T423 Paperbound \$1.00

*Prices subject to change without notice.*

*Dover publishes books on art, music, philosophy, literature, languages, history, social sciences, psychology, handicrafts, orientalia, puzzles and entertainments, chess, pets and gardens, books explaining science, intermediate and higher mathematics, mathematical physics, engineering, biological sciences, earth sciences, classics of science, etc. Write to:*

Dept. catrr.  
Dover Publications, Inc.  
180 Varick Street, N.Y. 14, N.Y.



The equation of motion for the pendulum is of the form :

$$\frac{d^2\theta}{dt^2} + \frac{mgl_0}{I} (\sin \theta - \sin \theta_e) = 0 \quad (2-34)$$

where the term  $mgl_0 \sin \theta_e$  represents the constant perturbing torque,  $I$  is the moment of inertia, and oscillations occur about the equilibrium angle  $\theta_e$ . If the perturbation is small relative to the maximum displacement, the angular frequency of oscillation is given, to a high degree of approximation, by:

$$\omega = \left( \frac{mgl_0}{I} \right)^{1/2} \quad (2-35)$$

The equation for phase oscillation is similar, with the phase angle in particle motion substituting for the angular displacement of the pendulum, the average accelerating energy per turn taking the place of the perturbing torque due to the weight, and the equilibrium phase substituting for equilibrium angle. To complete the analogy, however, the mass and length of the pendulum (its moment of inertia) should vary slowly to be equivalent to the increasing mass and energy of the particles.

Derivation of the equations of phase oscillation from the equations of motion goes beyond the scope of this monograph. However, the method will be described briefly, and the results given. The first two of the equations of motion are used, Eqs. 2-1 and 2-2. It is assumed that the phase oscillations are so slow and the amplitude of the associated radial oscillations so small, that we can neglect the radial acceleration term in Eq. 2-1. This is the basic assumption used by all writers utilizing the quasi-stationary theory; it is accomplished by letting  $dr/dt = 0$  in Eq. 2-1. We also neglect the radiation loss term  $dL/d\theta$ , which is significant only for electrons at very high energy. The betatron induction term  $d\Phi/dt$  is retained because it affects the oscillation energy associated with the changing orbit radius.

With these simplifications, when the two equations of